A 90-year-old man was referred to the clinic with a history of progressive shortness of breath and dry cough. Eighteen months ago, he was able to complete a full round of golf. He has now stopped playing as he is breathless after walking 200 m. He had been a heavy smoker all his life. He has no known asbestos exposure.

On examination, he is breathless at rest. He has a photosensitive rash on his face and neck. He is not clubbed. His pulse is irregularly irregular. Auscultation of his chest reveals fine inspiratory crackles bi-basally.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	121 g/l	135 - 175 g/l
White cell count (WCC)	9.2 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	6.1 × 10^9/l	1.5 - 7.0 × 10^9/l
Platelets (PLT)	203 × 10^9/l	150 - 400 × 10^9/l
Thyroid-stimulating hormone (TSH)	7.1 mu/l	0.4 - 5.0 mu/l

Blood gases on 30% oxygen:

pH	7.48
pCO_2	4.68 kPa
pO_2	8.8 kPa
Bicarbonate	25.4 mmol/l

His chest X-ray is shown below:



Created by Mahbub Alam Chhondo
From Bangladesh

Spirometry:

FEV_1	1.7 l (predicted 2.0 – 2.4 l)
FVC	2.0 l (predicted 3.2 – 3.8 l)
KCO	74% predicted

What is the likely cause of the suspected diagnosis?

Your answer was correct

A Rheumatoid arthritis

B Amiodarone

C Pesticides

D Sarcoidosis

E Smoking

Explanation

B Amiodarone

This man has interstitial lung disease – gradual onset of shortness breath and dry cough. He is not clubbed (but this could well be a feature) and has fine inspiratory crackles. Lung function tests are restrictive (reduced FEV1 and FVC with normal FEV1/FVC ratio) and he has type I respiratory failure. The chest radiograph shows small lung fields and reticular nodular shadowing.

Next you have to look for a cause. He has a photosensitive rash on his face, atrial fibrillation and abnormal thyroid function, suggesting that he is on amiodarone. Interstitial lung disease is an important side effect of long-term amiodarone administration.

Other side effects of amiodarone include:

- Hyper- and hypothyroidism (responsible for deranged TSH seen in this patient)
- Peripheral sensory neuropathy
- Hepatitis
- Photosensitivity
- Ataxia
- Corneal micro-deposits (reversible)
- Metallic taste
- Arrhythmias (torsade de pointes): a number of drugs cause pulmonary fibrosis – see below.

Diagnosis is made by a combination of clinical findings, lung function tests, CXR, BAL and high-resolution CT. It is confirmed with a VATS or an open lung biopsy in patients fit to undergo the procedure.

If there was no obvious cause for the interstitial lung disease and radiology and biopsy histology were consistent with usual interstitial pneumonia, then idiopathic pulmonary fibrosis would be the most likely diagnosis.

Drugs causing interstitial lung disease:

- Antibiotics – nitrofurantoin
- Cardiovascular – amiodarone, beta-blockers
- Chemotherapeutic – bleomycin, busulfan, cyclophosphamide, carmustine, lomustine, methotrexate, melphalan, mitomycin C, nitrosureas, sulfasalazine, 5-fluorouracil (5FU)
- Drugs used to treat rheumatoid arthritis – sulfasalazine, penicillamine, cyclophosphamide, methotrexate, gold
- Drugs used to treat migraine – ergotamine, ergots, methysergide
- Others:
  bromocriptine
  radiation
  dothiepin
  pesticides
  high-flow oxygen.

Causes of lower interstitial lung disease:

- Drugs
- Connective tissue disease
- Asbestosis
- Idiopathic pulmonary fibrosis.

A Rheumatoid arthritis

There is nothing in the history to suggest RA. Rheumatoid arthritis resulting in interstitial lung disease can cause shortness of breath, chronic dry cough, fatigue and weakness. There may be lung nodules.

C Pesticides

It is more likely that the ILD has been caused by amiodarone, given the patient’s sun-exposed rash.

D Sarcoidosis

This is not sarcoidosis, the presentation of which would be very unusual at this age. When sarcoidosis affects the lungs there is a collection of inflammatory cells forming lumps known as granulomas. This may present with wheezing, cough, shortness of breath or chest pain. Lofgren syndrome is where there is fever, lymphadenopathy, arthritis and erythema nodosum.

E Smoking

Amiodarone is more likely to cause ILD than smoking. Smoking is associated with desquamative interstitial pneumonia, but the history and age of the patient are against this.

70117

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 54-year-old man presented with increasing shortness of breath and orthopnoea. He had noticed these symptoms over the last few years but only went to his GP after he had had a severe episode of breathlessness when he went wading in the sea on holiday. On further enquiry, he had also become aware of morning headaches and an increasing tendency to fall asleep during the day.

On examination he looked well. Auscultation of his chest revealed reduced breath sounds and percussion note bi-basally.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	177 g/l	135 - 175 g/l
White cell count (WCC)	12.3 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	289 × 10^9/l	150 - 400 × 10^9/l
Packed cell volume (PCV)	0.51	0.4 - 0.53
Residual volume Erect (% predicted)	105	
Total lung capacity Erect (% predicted)	80	
Vital capacity Erect (% predicted)/Supine (% predicted)	75/50	
TLCO Erect (% predicted)	70	
KCO Erect (% predicted)	95	

What is the most likely diagnosis?

Your answer was incorrect

A Obstructive sleep apnoea

B Nocturnal asthma

C Bilateral diaphragmatic weakness

D Left ventricular failure

E Multiple sclerosis

Explanation

C Bilateral diaphragmatic weakness

In bilateral diaphragmatic weakness, patients commonly present with breathlessness on exertion and when lying flat. It can be associated with sleep apnoea, resulting in daytime somnolence and headaches. The symptoms are due to the paradoxical movement of the diaphragm during inspiration.

In the supine position, expansion of the ribs results in movement of abdominal contents into the chest, aided by gravity. Symptoms can be worse when standing up to the waist in water, as this counteracts the effects of gravity and prevents outward movement of the abdomen during inspiration and therefore produces a similar situation to being in the supine position.

A chest X-ray may show relatively small lung fields and basal linear shadowing due to sub-segmental collapse. A SNIF test (diaphragmatic screening on ultrasound) will show paradoxical movement of the diaphragm, particularly on sniffing. Blood gases may show a type II respiratory failure, particularly at night. Lung function tests classically show a low vital capacity, which falls further in the supine position. All lung volumes except the residual volume are reduced. The gas transfer tends to mildly impaired with a normal KCO. Treatment is with non-invasive ventilation.

B Nocturnal asthma

The patient has no symptoms to suggest that asthma is the diagnosis. There is no wheeze or cough. The breathlessness is progressive and there are several other features in this history to suggest diaphragmatic weakness.

A Obstructive sleep apnoea

The 15% change in vital capacity when supine alongside breathlessness when submerged in water and orthopnoea make diaphragmatic weakness more likely.

D Left ventricular failure

Orthopnoea and dull percussion notes bilaterally (due to pleural effusions) in isolation could be in keeping with LVF, but there is more going on here. The patient has a change in sitting and supine VC, there is a normal transfer factor when corrected for lung volumes and haemoglobin is high (you would expect a lower Hb in LVF due to fluid overload and haemodilution).

E Multiple sclerosis

Patients with MS can experience diaphragmatic weakness, but there are no other clinical features to suggest an underlying diagnosis of MS.

70150

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 41-year-old woman presented with fatigue, joint aches and dry, painful, red eyes.

Routine blood testing reveals a calcium level just above the upper end of the normal range. A biopsy of one of her skin lesions was taken.



Which enzyme is activated by macrophages in this condition?

Your answer was correct

A	Lactate dehydrogenase
B	1-alpha hydroxylase
C	Lysozyme
D	Glucose-6-phosphate dehydrogenase
E	Acid phosphatase

Explanation

B 1-alpha hydroxylase

1-alpha hydroxylase is activated by macrophages in sarcoid granulomas (the diagnosis here), and converts 25-hydroxyvitamin D3 to 1,25-dihydroxyvitamine D3, the active form of the vitamin. Absorption of calcium from the gut increases as a consequence as does renal calcium reabsorption from the distal tubule.

A Lactate dehydrogenase

Lactate dehydrogenase is raised in myonecrosis, haematological malignancies and in patients with haemolysis.

C Lysozyme

Lysozyme is an enzyme which drives bacterial cell wall lysis, it is not involved in the pathogenesis of sarcoidosis.

D Glucose-6-phosphate dehydrogenase

Deficiency of G-6PD leads to red cell haemolysis in the presence of drugs which act as oxidising agents.

E Acid phosphatase

Acid phosphatase is used as a marker for prostate cancer.

39099

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

A 54-year-old man presents to the Respiratory Medicine Clinic due to cough and shortness of breath on exertion for the past six months. He has a past medical history of gastro-oesophageal reflux disease, for which he takes omeprazole 10 mg per day, and he smokes ten cigarettes per day. Pulmonary function tests reveal a forced expiratory volume in one second (FEV_1) to forced vital capacity (FVC) ratio of 1, with FEV_1 85% of predicted and FVC 84% of predicted. Chest X-ray reveals increased interstitial lung markings, and high-resolution computed tomography (HRCT) of the chest reveals a pattern in keeping with usual interstitial pneumonia.

Given the likely diagnosis, which of the following intervention options for this patient has been shown to have an impact on survival time?

Your answer was incorrect

A Administration of proton pump inhibitor (PPI)

B Nintedanib

C Oxygen therapy

D Pulmonary rehabilitation

E Smoking cessation

Explanation

A Administration of proton pump inhibitor (PPI)

This patient has idiopathic pulmonary fibrosis (IPF), as evidenced by his restrictive pattern of lung disease on pulmonary function testing and the pattern of usual interstitial pneumonia on high-resolution computed tomography. Minimal treatment options are currently available for IPF. Pulmonary rehabilitation is recommended to improve quality of life; smoking cessation is recommended to prevent the occurrence of co-morbid chronic obstructive pulmonary disease (COPD), and nintedanib and pirfenidone have been shown to reduce the functional decline in IPF but are currently only recommended, according to the National Institute for Health and Care Excellence guidelines, to be started in individuals when the person has a forced vital capacity (FVC) of between 50% and 80% predicted, with the caveat that medication is stopped if the disease progresses (a confirmed decline in percentage predicted FVC of 10% or more) in any 12-month period. The only intervention that has been shown to be independently associated with increased survival in patients with IPF and gastro-oesophageal reflux disease is PPIs.

B Nintedanib

Both pirfenidone and nintedanib have been trialled in patients with IPF and have shown beneficial outcomes in reducing the functional decline in IPF. However, there is currently no evidence that these medications are able to improve survival.

C Oxygen therapy

Currently, there is no evidence that oxygen therapy improves survival in patients with IPF. Its beneficial effects may be an improvement in exercise tolerance and a reduction in the risk of developing pulmonary hypertension and cor pulmonale.

D Pulmonary rehabilitation

Pulmonary rehabilitation has been proven to improve quality of life and functional status. It does not show any impact on survival time and the effects seen are reduced six months after the programme. This may be related to the finding that patients do not keep up with exercises at home following a rehabilitation programme.

E Smoking cessation

Smoking cessation is recommended in patients with IPF to reduce the risk of developing concomitant COPD. However, it is not independently associated with improvement in survival time.

72998

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 24-year-old man presents to the Emergency Admission for being unwell over the last couple of days. He tells you that he has had flu for the last four days. He complains of a dry cough, headache, fever, malaise and muscle pains.

On examination, he is unwell with a blood pressure of 110/60 mmHg, pulse 92/min and a respiratory rate of 22/min. His oxygen saturation is 95% on air. On auscultation, he has scattered crackles in both lung fields. His cardiovascular and abdominal examinations are unremarkable.

Investigations reveal the following:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	98 g/l	135–175 g/l
White cell count (WCC)	12.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	278 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	130 mmol/l	135–145 mmol/l
Potassium (K^+)	3.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	112 μmol/l	50–120 μmol/l
Urea	8.1 mmol/l	2.5–6.5 mmol/l
Mean corpuscular volume (MCV)	89 fl	76–98 fl
Blood film	Red cell agglutination reticulocytosis, occasional spherocytes	

Chest X-ray shown below



What is the likely diagnosis?

Your answer was correct

A *Streptococcus pneumoniae*

B *Mycoplasma pneumoniae*

C *Chlamydia pneumoniae*

D *Legionella pneumophila*

E Viral pneumonia

Explanation

B *Mycoplasma pneumoniae*

Mycoplasma pneumoniae is a disease of young adults. It presents with flu-like symptoms with low-grade fever, dry cough, headache and myalgia. Cold agglutinins may cause an autoimmune haemolytic anaemia. Coombs' test is positive. The low sodium is related to SIADH. The chest x-ray here shows perihilar inflammation with increased interstitial markings in the lower zones. Treatment is with clarithromycin or doxycycline. It is important to note that clarithromycin should be given orally unless contraindicated, as it has good bio-availability when given orally and can often cause thrombophlebitis when given intravenously.

Image source: https://radiopaedia.org/cases/atypical-pneumonia-mycoplasma?lang=gb

A *Streptococcus pneumoniae*

Streptococcal pneumonia is the most common type of pneumonia but has higher incidence in the elderly, post-splenectomy and in those with pre-existing heart or lung disease. Patients usually present with fever, pleurisy and can develop sepsis. Chest X-ray demonstrates lobar consolidation, and diagnosis can be confirmed with urinary antigen. The dry cough and myalgia features, as well as electrolyte disturbances, make this a less likely diagnosis.

C *Chlamydia pneumoniae*

Spread occurs between people and follows a biphasic course, first with pharyngitis, hoarseness and otitis, then leading to pneumonia. Diagnosis is with chlamydophila fixation test and PCR. Treatment is with clarithromycin. The absence of upper airway symptoms makes this a less likely diagnosis.

D *Legionella pneumophila*

Legionella has an incubation period of 2–10 days, and presents with fever, rigors, headache, myalgia, dry cough, progressively increasing respiratory distress and confusion. Diarrhoea and vomiting are common, as are non-specific liver function test abnormalities. Relative bradycardia may be seen in some patients, in contrast to other acute pneumonias. Pleural effusion may be seen in up to 50%, and chest X-ray appearance may take up to 4 months to return to normal. Diagnosis is made with serology, or with PCR on urinary *Legionella* antigen. Treatment is with quinolones as first-line therapy, with macrolides as a potential alternative. Complications include pericarditis, encephalitis and kidney disease. Hyponatremia is frequently seen in legionella pneumonia. This is a strong differential diagnosis, but the absence of travel history makes this a less likely diagnosis.

E Viral pneumonia

Viral pneumonia can occur with influenza, measles, CMV and varicella zoster. Whilst the symptoms of myalgia would match with a viral infection, the presence of haemolysis would not be explained, making it an unlikely diagnosis.

7513

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 45-year-old non-smoking man remains breathless at night with a polyphonic wheeze, despite already being prescribed a short-acting b2-agonist by his general practitioner (GP). He is concordant with treatment and has a good inhaler technique.

His spirometry result is shown below.



Which is the next appropriate therapy to offer according to current NICE guidelines?

Your answer was incorrect

A Long-acting β2-agonist (LABA)

B Inhaled corticosteroids

C Leukotriene receptor antagonist (LTRA)

D No further treatment

E Long-acting muscarinic antagonist

Explanation

B Inhaled corticosteroids

This clinical scenario and flow–volume loop are suggestive of asthma. The spirometry suggests an obstructive picture, with a concave curve pattern. Inhaled corticosteroids at a low dose are the first-line maintenance therapy for adults with uncontrolled asthma.

A Long-acting β2-agonist (LABA)

A LABA should only be offered after inhaled corticosteroids and an LTRA have already been offered. The LTRA should then be reviewed after six to eight weeks after the addition of a LABA to see if it should be continued.

C Leukotriene receptor antagonist (LTRA)

An LTRA is only recommended as a stage in adult asthma management after low-dose inhaled corticosteroids have been initiated.

D No further treatment

The patient is experiencing nocturnal breathlessness, which is an indication for increasing maintenance therapy for asthma.

E Long-acting muscarinic antagonist

A long-acting muscarinic antagonist is used as the next stage in the management of COPD. However, the spirometry result is more in keeping with asthma than COPD, making this answer incorrect.

71788

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

National Institute for Health and Care Excellence. 2017. Asthma: diagnosis, monitoring and chronic asthma management.
nice.org.uk/guidance/ng80/chapter/Recommendations
(https://www.nice.org.uk/guidance/ng80/chapter/Recommendations)

Asthma

Media

Expanded explanations

Flashcards

A 42-year-old man, who is known to be HIV positive, presents with a cough productive of rusty-coloured sputum. He has been feeling unwell for around a week or so. On examination he is pyrexial (38.2°C) with a BP of 135/70 mmHg. His pulse is 95/min and regular. Auscultation of the chest reveals an area of bronchial breathing over the left lower lung field.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	124 g/l	135–175 g/l
White cell count (WCC)	13.1 × 10^9/l (Neutrophilia)	4.0–11.0 × 10^9/l
Platelets (PLT)	195 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	100 µmol/l	50–120 µmol/l
Chest X-ray (CXR)	Left. lower lobe consolidation	

His chest x-ray is below



Which of the following types of pneumonia is the most likely diagnosis?

Your answer was incorrect

A *Pneumocystis*

B *Staphylococcus aureus*

C *Mycoplasma*

D *Streptococcus pneumoniae*

E *Klebsiella*

Explanation

D *Streptococcus pneumoniae*

This man has community-acquired pneumonia which is most likely to be *S. pneumoniae*. The estimated attack rate is higher in HIV-infected patients (5.9/1000 vs 0.31/1000 in non-HIV infected patients). Resistance patterns are similar in HIV and non-HIV infected patients, and invasive disease is rare and tends to occur in patients with a low CD4 count. The chest x-ray showing lobar consolidation (note the absence of the outline of the left hemidiaphragm indicating left lower lobe infection) and neutrophilia fits best with a *S. pneumoniae* episode rather than another cause for the respiratory tract infection.

Image source: http://www.svuhradiology.ie/case-study/left-lower-lobe-consolidation/

C *Mycoplasma*

Mycoplasma pneumonia often follows a slightly more subacute course than streptococcal pneumonia, with a dry cough, chest pain and wheeze, that can be present for a few days before patients engage with medical services.

A *Pneumocystis*

Pneumocystis pneumonia is associated with diffuse patchy consolidation, and the hallmark of the condition is desaturation on exercise (usually measured with a hand-held pulse oximeter).

B *Staphylococcus aureus*

Staphylococcus aureus pneumonia, even in the context of underlying HIV infection, normally follows an episode of influenza.

E *Klebsiella*

Klebsiella pneumonia is associated with lobar infection with cavitation and is seen most commonly in patients with a history of alcoholism.

20635

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 28-year-old man who is currently prescribed highly active anti-retroviral therapy (HAART) comes to the clinic for review. He is extremely worried as he has been working in a homeless shelter with a man who has recently been identified as having active pulmonary tuberculosis. He is currently feeling well.

On examination, his BP is 122/72 mmHg, pulse is 80/min and regular and he is apyrexial. His chest is clear. His BMI is 25 and has been stable over the past six months at least. His CD4 count is 420, and an interferon gamma screen for TB is positive.

Which of the following is the most appropriate next step?

Your answer was incorrect

A Quadruple anti-tuberculous therapy with rifampicin, pyrazinamide, ethambutol and isoniazid for six months

B Dual anti-tuberculous therapy with rifampicin and pyrazinamide for two months

C Dual anti-tuberculous therapy with rifampicin and isoniazid for three months

D Monotherapy with isoniazid for six to nine months

E Monotherapy with isoniazid for one year

Explanation

D Monotherapy with isoniazid for six to nine months

Monotherapy with isoniazid for six months can be associated with poor compliance but is the regime of choice as per the WHO guidelines on treatment of latent TB in patients with HIV on HAART. This man will hopefully engage with treatment as he seems to have a stable lifestyle. This will give his the best chance of both treating the latent TB and avoiding dangerous drug interactions if a rifampicin containing regime was used.

B Dual anti-tuberculous therapy with rifampicin and pyrazinamide for two months

In an attempt to improve adherence, a 2-month rifampicin and pyrazinamide regime was considered but this is not widely used because of reported cases of hepatotoxicity.

A Quadruple anti-tuberculous therapy with rifampicin, pyrazinamide, ethambutol and isoniazid for six months

Quadruple anti-tuberculous therapy is the intervention for patients suffering from active TB.

C Dual anti-tuberculous therapy with rifampicin and isoniazid for three months

This patient fits the criteria for a diagnosis of latent TB and, in this situation, against a background of HIV, anti-tuberculous chemotherapy is considered. A commonly used option is rifampicin and isoniazid for a 3-month period, but this has high incidence of harmful drug interactions with HAART and this patient likely does not have a chaotic lifestyle as he works, so would hopefully engage with monotherapy treatment.

E Monotherapy with isoniazid for one year

One year of monotherapy with isoniazid is not a commonly recognised regime. The 2015 WHO guidelines on treatment of latent TB discuss monotherapy for up to 36 months in high incidence areas, in lower incidence areas the usual course is six to nine months.

71624

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BHIVA guidelines for the treatment of TB/HIV coinfection 2011
bhiva.org/TB-HIV-coinfection-guidelines.aspx
(http://www.bhiva.org/TB-HIV-coinfection-guidelines.aspx)

A 51-year-old man, while an inpatient for a routine hernia operation, was noted to have a pneumothorax on his preoperative chest X-ray. He has a history of chronic bronchitis and a 50 pack-year smoking history. His chest X-ray shows a unilateral pneumothorax of 2.2 cm. He is breathless but does not have chest pain.

Which of the following would be the ideal management for this patient?

Your answer was incorrect

A Admit the patient for observation only

B Aspirate the pneumothorax with a 16 G cannula

C Discharge the patient to Outpatient Ambulatory Clinic in two weeks

D Insert a 14 Fr gauge chest drain

E Give the patient 10 litres of oxygen by mask

Explanation

D Insert a 14 Fr gauge chest drain

This patient has a secondary pneumothorax, as he is over the age of 50 and has a significant smoking history. The British Thoracic Society Guidelines suggest that a chest drain 8–14 Fr should be inserted if the pneumothorax is > 2 cm in size or the patient is breathless. The intrapleural distance should be measured at the level of the hilum.

C Discharge the patient to Outpatient Ambulatory Clinic in two weeks

This patient has a secondary pneumothorax and he is symptomatic. Guidelines suggest that in this situation a chest drain should be inserted. Therefore outpatient management would be inappropriate.

A Admit the patient for observation only

Admission without intervention can be considered for patients with a secondary pneumothorax where the pneumothorax is < 1 cm in size and the patient is asymptomatic (not breathless).

B Aspirate the pneumothorax with a 16 G cannula

An attempt at aspirating a secondary pneumothorax should only be made where the pneumothorax is between 1 and 2 cm, and the patient is not breathless.

E Give the patient 10 litres of oxygen by mask

High-flow oxygen may be used as a treatment option for secondary pneumothoraces < 1 cm in size, provided that the patient is not at risk of oxygen sensitivity.

72961

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society (BTS): BTS Pleural Disease Guideline 2010 Management of spontaneous pneumothorax
thorax.bmj.com/content/65/Suppl_2/ii18
(https://thorax.bmj.com/content/65/Suppl_2/ii18)

Pneumothorax

Media

Expanded explanations

Flashcards

A 72-year-old man with a history of COPD and symptoms of benign prostatic hypertrophy presents to the clinic complaining of bone pain. He is a 40 pack-year smoker and takes a range of medications including ramipril, aspirin, atorvastatin, finasteride and a salmeterol/fluticasone combination inhaler. He has lost a few kilos in weight over the past few months without trying, and has seen his GP due to an episode of pneumonia. On examination his BP is 155/81 mmHg, with pulse 75/min (atrial fibrillation). There are coarse crackles on auscultation of the chest consistent with COPD. He looks clubbed.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	110 g/l	135–175 g/l
White cell count (WCC)	5.2 × 10^9/l	4–11 × 10^9/l
Platelets	198 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine	130 μmol/l	50–120 μmol/l
Erythrocyte sedimentation rate (ESR)	60 mm/hr	0–10 mm in the 1st hour
Corrected (Ca^{2+})	2.3 mmol/l	2.20–2.60 mmol/l

You arrange a bone scan, which reveals a symmetrical increase in uptake along the tubular bones and around the periarticular surfaces. The mandible and both scapulae also show signs of increased isotope uptake.

Which of the following is the most likely diagnosis?

Your answer was correct

A Multiple bony metastases from a lung primary

B Paget’s disease

C Hypertrophic pulmonary osteoarthropathy

D Osteomalacia

E Primary hyperparathyroidism

Explanation

C Hypertrophic pulmonary osteoarthropathy

Hypertrophic pulmonary osteoarthropathy is a condition seen in patients with a bronchial carcinoma, and may predate the discovery of the underlying lesion. It is characterised by finger clubbing and long bone pain. The most sensitive investigation for making the diagnosis is the isotope bone scan: the findings are typically uptake in long bones, around periarticular surfaces and sometimes also the mandible and scapulae. Symptoms of the condition may resolve when the primary tumour is removed. A chest X-ray has the potential to reveal the primary lung tumour.

A Multiple bony metastases from a lung primary

Symmetrical increase in uptake along tubular bones around periarticular surfaces is seen here. If multiple bony metastases were the cause of this patient's symptoms you would expect haphazard multiple areas of increased uptake over the bone scan.

B Paget's disease

Paget's disease often involves one joint only, and most frequently this is only one side of the pelvis. A rise in alkaline phosphatase is seen, in the presence of a normal calcium.

D Osteomalacia

Osteomalacia is associated with a borderline/low serum calcium, proximal bone pain and muscle weakness.

E Primary hyperparathyroidism

Primary hyperparathyroidism is associated with hypercalcaemia, rather than the normal calcium level seen here.

32453

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 72-year-old gentleman is referred to the Respiratory Outpatients Clinic with a 2-month history of increasing shortness of breath and cough. Over the last few weeks, he has coughed up a few streaks of blood. He has also noticed difficulty getting up from a chair, and that his clothes have been looser recently without trying to lose weight. He is taking regular inhaler therapy but is unsure of their names.

On examination, he is clubbed. Auscultation of his chest reveals reduced expansion and breath sounds on the left. Heart sounds are normal. He has a 2 cm hepatomegaly on palpation of his abdomen. He has reduced power (4/5) on hip flexion bilaterally and generally reduced tendon reflexes. Plantars both show a flexor response; proprioception and sensation are normal.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	109 g/l	135 - 175 g/l
White cell count (WCC)	12.2×10^9/l	4.0 - 11.0×10^9/l
Platelets (PLT)	491×10^9/l	150 - 400×10^9/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	58 mm/h	1 - 20 mm/h
Sodium (Na^+)	126 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.6 mmol/l	3.5 - 5.0 mmol/l
Urea	5.9 mmol/l	2.5 - 6.5 mmol/l
Creatinine	125 µmol/l	50 - 120 µmol/l
Bilirubin	49 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	120 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	441 u/l	30 - 150 u/l
Albumin	32 g/l	35 - 55 g/l
Corrected calcium (Ca^{2+})	2.74 mmol/l	2.2 - 2.7 mmol/l
Phosphate (PO_4^{3-})	0.81 mmol/l	1.1 - 1.45 mmol/l

His chest X-ray is shown below:



Which investigations would you organise next to aid diagnosis and treatment?

Your answer was incorrect

A Bone scan and bronchoscopy

B CT brain and liver ultrasound

C Bronchoscopy and PTH level

D MRI chest and bone scan

E CT chest/upper abdomen and bronchoscopy

Explanation

E CT chest/upper abdomen and bronchoscopy

The chest X-ray shows left lower lobe collapse. There is loss of volume of the left hemi-thorax and the classical ‘sail sign’ behind the heart.

In addition to this suggestive CXR, this man also has clinical features of lung malignancy weight loss, haemoptysis, dyspnoea and hypercalcaemia.

His investigations and clinical examination also demonstrate evidence of paraneoplastic syndromes which are associated with certain types of lung cancer:

- hyponatraemia secondary to the syndrome of inappropriate antidiuretic hormone secretion
- the proximal weakness and hyporeflexia which, along with autonomic features, suggests Lambert-Eaton myasthenic syndrome.

These are seen more frequently with small cell lung cancer. Ectopic ACTH production is also associated but rare.

Squamous cell carcinoma is associated with:

- gynaecomastia
- hyperthyroidism
- hypertropic pulmonary osteoarthropathy (HPOA) finger-clubbing, periostitis and arthritis
- hypercalcaemia usually secondary to bone metastases, but rarely may be secondary to secretion of PTH-related peptide.

Common to both:

- hypercalcaemia: the corrected calcium here is 2.74 mmol/ , due to bony metastases. If albumin is <40 g/l, corrected calcium = $[Ca^{2+}] + 0.02 \times 40\text{-}[alb]$ mmol/l
- clubbing
- smoking: he is likely to be a smoker or ex-smoker and have COPD as he is on inhalers.

A histological diagnosis is important, as is accurate staging. Therefore the most appropriate next investigation would be a bronchoscopy and biopsy, and a staging CT scan. Following this, a bone scan would be useful to confirm bone metastases. Since this patient has deranged liver function tests and hypercalcaemia, it is likely he has metastatic disease. Treatment of small cell lung cancer is primarily palliative with chemotherapy, as metastases tend to occur early. Surgical resection may be considered in combination with chemotherapy if it presented as a solitary pulmonary nodule, but this is rare. Palliative care involvement is important for symptomatic control or if the patient is unsuitable for other treatment options.

C Bronchoscopy and PTH level

While bronchoscopy could provide histological diagnosis, PTH would not be a diagnostic test for malignancy. NICE guidance recommends that ‘patients with known or suspected lung cancer should be offered a contrast-enhanced chest CT scan to further the diagnosis and stage the disease’.

A Bone scan and bronchoscopy

While the hypercalcaemia may suggest metastatic disease warranting a bone scan in due course, NICE guidance recommends that ‘Patients with known or suspected lung cancer should be offered a contrast-enhanced chest CT scan to further the diagnosis and stage the disease’; the CT scan should be performed first to stage disease, in association with bronchoscopy for histological diagnosis.

B CT brain and liver ultrasound

This would represent an inadequate combination of investigations in this case, and would provide neither histological nor radiological confirmation of the diagnosis. The patient’s neurological symptoms are more likely caused by paraneoplastic effects than a cerebral metastasis, and while liver ultrasound may be appropriate to examine for metastases, this alone would not provide adequate staging of the primary tumour.

D MRI chest and bone scan

CT scan, rather than MRI, is the recommended imaging modality for staging of lung cancer. Bronchoscopy is recommended for histological diagnosis.

70132

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Lung cancer: diagnosis and management
nice.org.uk/guidance/cg121/
(http://www.nice.org.uk/guidance/cg121/)

A 55-year-old asthmatic patient presents to the Emergency Department with increasing cough and wheeze. She describes a cough productive of brown sputum. There is no other past medical history of note. She is taking a steroid inhaler regularly and increasing doses of beta-agonist, with no alleviation of her symptoms. She has worked in a bakery for many years. She is an ex-smoker and stopped smoking 20 years ago; she has a pet dog and one cat.

On examination she has a temperature of 38°C and is breathless at rest. Her BP is 122/82 mmHg, with pulse 88/min and regular. There is reduced chest expansion anteriorly on the left side, with corresponding reduced breath sounds. She has a mild expiratory wheeze.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	141 g/l	135–175 g/l
White cell count (WCC)	12.7 × 10^9/l	4.0–11.0 × 10^9/l
Neutrophils	7.37 × 10^9/l	1.5–7.0 × 10^9/l
Lymphocytes	2.55 × 10^9/l	1.5–4.0 × 10^9/l
Eosinophils	2.1 × 10^9/l	0.1–0.4 × 10^9/l
Monocytes	0.6 × 10^9/l	0.0–0.8 × 10^9/l
Basophils	0.05 × 10^9/l	< 0.1 × 10^9/l
Mean corpuscular volume (MCV)	87.3 fl	80–100 fl
Platelets (PLT)	314 × 10^9/l	150–400 × 10^9/l
Erythrocyte sedimentation rate (ESR)	28 mm/h	1–20 mm/h
C-reactive protein (CRP)	16 mg/l	< 10 mg/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5–5.0 mmol/l
Urea	4.8 mmol/l	2.5–6.5 mmol/l
Creatinine	95 µmol/l	50–120 µmol/l
Bilirubin	12 µmol/l	1–22 µmol/l
Alanine aminotransferase (ALT)	49 u/l	7–55 u/l
Alkaline phosphatase (ALP)	61 u/l	30–150 u/l
Albumin	36 g/l	35–55 g/l

X-ray scan:



CT scan on presentation:



CT scan taken one month later:



What investigation would help make a diagnosis?

Your answer was correct

A p-ANCA

B c-ANCA

C Positive skin test/radioallergosorbent test for *Aspergillus fumigatus*

D Metacholine challenge test

E Lung function tests

Explanation

C Positive skin test/radioallergosorbent test for *Aspergillus fumigatus*

The chest X-ray shows left upper lobe collapse and demonstrates the ‘veil sign’. This patient has allergic bronchopulmonary aspergillosis (ABPA). Pulmonary infiltrates, blood eosinophilia and asthma are usually secondary to Churg–Strauss syndrome, or allergic bronchopulmonary aspergillosis.

ABPA is suspected in any patient with asthma who has an abnormal chest X-ray and high peripheral blood eosinophilia. The chest X-ray may show diffuse pulmonary infiltrates, and pulmonary, lobar or segmental collapse occurs as a transient feature. The most common cause is sensitivity to *A. fumigatus* spores.

The diagnostic criteria include:

- Asthma (in most cases)
- Peripheral blood and sputum eosinophilia
- Abnormal chest X-ray (infiltrates, segmental or lobar collapse)
- Positive skin tests/RAST to an extract of *A. fumigatus*
- *A. fumigatus* IgG serum-precipitating antibodies
- Raised total IgE >1000 ng/ml
- Fungal hyphae of *A. fumigatus* on microscopy of sputum.

A p-ANCA

This is an appropriate test for vasculitis and is associated with Churg–Strauss syndrome. The absence of other features of vasculitis, such as mononeuritis, counts against this as the underlying diagnosis.

B c-ANCA

This is an appropriate test for vasculitis. C-ANCA positivity is associated with Granulomatosis with polyangiitis, where features such as sinusitis, haemoptysis and renal vasculitis co-exist.

D Metacholine challenge test

The methacholine challenge test is used to diagnose asthma, usually where there is some doubt as to the status with respect to hyper-reactivity.

E Lung function tests

Lung functions tests would not be sufficiently discriminatory to diagnose allergic bronchopulmonary aspergillosis. They would merely demonstrate a mixed obstructive/restrictive picture.

70095

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A 54-year-old man presents to the Respiratory Clinic with his wife, who is extremely worried that he snores heavily every night and often seems to stop breathing. He also falls asleep each evening while watching the TV and has had to stop the car on a number of occasions to take a nap. He drinks two pints of beer each day and smokes five cigarettes per day.

On examination, his BP is 152/85 mmHg and his heart rate is 80 bpm and regular; his BMI is elevated at 32 kg/m^2, and his collar size is 18. His Epworth score is 15. He plans to lose weight.

Which initial intervention is the most appropriate?

Your answer was incorrect

A Continuous positive airway pressure

B Gastric banding

C Mandibular advancement device

D Modafinil

E Uvulopalatopharyngoplasty

Explanation

A Continuous positive airway pressure

For patients with mild symptoms that affect their quality of life and usual daytime activities, the National Institute for Health and Care Excellence (NICE) guidelines recommend fixed-level continuous positive airway pressure (CPAP). Therapy with CPAP is effective in reducing symptoms and is associated with an improvement in cardiovascular morbidity in patients with hypertension and obstructive sleep apnoea. Forty per cent of patients, however, have an elevated Epworth score after CPAP is initiated, and these individuals should, of course, be encouraged to take other measures such as losing weight.

B Gastric banding

Gastric banding would not be appropriate here as it would not treat all presentations and should be considered for patients who are unable to achieve significant weight loss.

C Mandibular advancement device

Patients with mild symptoms that affect their quality of life and usual daytime activities and who are unable to tolerate or decline to try CPAP may try a mandibular advancement device (MAD). A MAD may relieve obstruction and give this patient the opportunity to lose weight and improve his symptoms significantly. He should, of course, also be referred for a sleep study and advised that until his symptoms significantly improve, he is not to drive. In the event that the device is ineffective and/or he fails to lose significant weight, he can be referred for an alternative management option.

D Modafinil

Modafinil would not be appropriate here as it is used to enhance wakefulness in patients with central sleep disorders.

E Uvulopalatopharyngoplasty

Uvulopalatopharyngoplasty is considered in patients who are resistant to other interventions or who have significant obstructive sleep apnoea despite weight loss.

72129

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Obstructive sleep apnoea syndrome
cks.nice.org.uk/obstructive-sleep-apnoea-syndrome#!scenario
(https://cks.nice.org.uk/obstructive-sleep-apnoea-syndrome#!scenario)

A 64-year-old man comes to the Respiratory Clinic for review. He has been referred by his General Practitioner because of increasing shortness of breath due to chronic obstructive pulmonary disease (COPD). He continues to smoke ten cigarettes per day.

A few years ago, he had a trial of oral steroids which did lead to a positive impact on his peak flow, and he reports being diagnosed with asthma as a child. His most recent forced expiratory volume in 1 second (FEV_1) is 40% of predicted, and he can walk only 50 m to the local bus stop.

Current medications include a salbutamol inhaler as needed (prn) and carbocisteine.

Examination reveals a blood pressure of 135/82 mmHg and a pulse of 78 bpm and regular. He has signs of right heart failure.

Which of the following is most appropriate to reduce the risk of future exacerbations?

Your answer was incorrect

A Long-acting beta agonist (LABA) and long-acting muscarinic antagonist (LAMA)

B Short-acting muscarinic antagonist (SAMA)

C Long-term oxygen therapy (LTOT)

D LABA and high-dose inhaled corticosteroid (ICS)

E Theophylline

Explanation

D LABA and high-dose inhaled corticosteroid (ICS)

In clinical trials, the combination of LABA and high-dose ICS has shown an ability to reduce the frequency of exacerbations, although this is at the expense of an increase in the risk of pneumonia. As evidence has developed from large intervention trials comparing LABA + LAMA to LABA + ICS, it has become apparent that LABA + ICS is the most effective in patients who have asthmatic features/features of steroid responsiveness in their disease, as is seen here.

B Short-acting muscarinic antagonist (SAMA)

Ipratropium may improve symptoms, but it does not have a positive impact on outcomes in COPD, so that it would not represent the intervention of choice.

A Long-acting beta agonist (LABA) and long-acting muscarinic antagonist (LAMA)

LABA and LAMA in combination is the intervention of choice for patients with symptomatic COPD where there are no features of asthma or steroid responsiveness.

C Long-term oxygen therapy (LTOT)

LTOT improves mortality by reducing progression of right heart failure and pulmonary hypertension rather than by reducing exacerbations, but its use is less than desirable in patients who continue to smoke because of fire risk.

E Theophylline

Theophylline may improve symptoms, but it does not improve mortality in COPD and increases the risk for atrial arrhythmias in patients who take it. Inhaled beta agonists are the preferred option.

39521

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

National Institute for Health and Care Excellence. 2019. Chronic obstructive pulmonary disease in over 16s: non-pharmac...
nice.org.uk/guidance/ng115/resources/visual-summary-treatment-algorithm-pdf-6604261741
(https://www.nice.org.uk/guidance/ng115/resources/visual-summary-treatment-algorithm-pdf-6604261741)

You review a 48-year-old man in the Respiratory Clinic. He is diagnosed with tuberculosis and you want to start quadruple anti-tuberculous therapy. He is very concerned about the potential side effects of therapy. Previous history of note includes abnormal liver function tests and a history of alcohol abuse; most recently he has been diagnosed with type 2 diabetes. Recent ALT was 65 U/l, creatinine 102 μmol/l and a liver biopsy has shown early hepatic fibrosis.

Which of the following is important to bear in mind when managing his anti-tuberculous therapy?

Your answer was incorrect

A Ethambutol should not be given

B Isoniazid should not be given

C Pyrazinamide should not be given

D Pyridoxine should not be given

E Rifampicin should not be given

Explanation

C Pyrazinamide should not be given

Drug-induced hepatitis associated with anti-tuberculous therapy can be fatal; for this reason pyrazinamide should not be used in patients with known chronic liver disease, even if it is thought to be stable. Baseline and regular monitoring of liver function should be undertaken in patients on anti-tuberculous therapy, and where there is a history of chronic liver disease it is advised that this is weekly for the first two weeks and then two-weekly thereafter.

B Isoniazid should not be given

In stable liver disease it is acceptable to give up to two drugs which may have hepatotoxicity, usually isoniazid and rifampicin only.

A Ethambutol should not be given

Ethambutol can accumulate in patients with chronic kidney disease, leading to optic neuropathy, and therefore should be given at a lower dose. The creatinine of 102 μmol/l seen here means that dose adjustment is not required.

D Pyridoxine should not be given

Pyridoxine should be given in addition to isoniazid in those with renal impairment, to prevent peripheral neuropathy.

E Rifampicin should not be given

Rifampicin should only be avoided where there is decompensated liver disease and in those patients taking sulphonylureas where it may reduce the effectiveness of these agents.

70925

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 67-year-old non-smoker was referred to the Respiratory Consultant with a 3-month history of progressive shortness of breath. She was also complaining of a right-sided chest pain, which she described as a constant severe ache. She had lost about 5 kg in the last month and felt lethargic. She had initially put the symptoms down to the stress of recently being widowed. Her husband, who used to work as a plumber, died of a 'lung problem'. She has never worked outside the home, and has two daughters who are fit and well. On examination she looked unwell and was breathless at rest. Examination of her chest revealed reduced expansion, percussion note, vocal fremitus and breath sounds throughout the right hemithorax.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	102 g/l	135 - 175 g/l
White cell count (WCC)	10.8 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	471 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	81 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	69 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	44 mg/l	< 10 mg/l
Sodium (Na^+)	136 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.7 mmol/l	3.5 - 5.0 mmol/l
Urea	6.6 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	77 µmol/l	50 - 120 µmol/l
Bilirubin	34 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	21 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	97 u/l	30 - 150 u/l
Albumin	37 g/l	35 - 55 g/l

Her chest X-ray is shown below:



What is the most likely diagnosis?

Your answer was incorrect

A Asbestosis

B Squamous-cell carcinoma

C Pleural adenocarcinoma

D Mesothelioma

E Metastatic adenocarcinoma

Explanation

D Mesothelioma

The clue is in the question – her husband who has died worked as a plumber. She is a housewife (she has never worked outside the home), so it is assumed she may have been exposed to her husband’s overalls and was therefore exposed to asbestos. She has systemic symptoms as well as a pleural effusion, making the most likely diagnosis malignant mesothelioma.

Video-assisted thoracoscopy surgery is the investigation of choice as it has the highest diagnostic yield.

The chest X-ray shows pleural thickening and an effusion on the right. There is a holly-leaf pleural plaque on the left (directly above the heart to the left of the aortic knuckle), consistent with previous asbestos exposure. Incidentally, there is pericardial calcification.

C Pleural adenocarcinoma

This would be the main differential diagnosis but the probable history of asbestos exposure makes mesothelioma more likely.

A Asbestosis

Whilst the history is suggestive of an asbestos related pathology the history of right chest pain with radiographic evidence of significant pleural thickening is suggestive of mesothelioma with chest wall invasion.

B Squamous-cell carcinoma

The lady is a non-smoker with a history to suggestive of asbestos exposure and the chest radiograph shows significantpleural thickening. All these factors make mesothelioma more likely than small cell lung cancer.

E Metastatic adenocarcinoma

Again, the history is suggestive of asbestos exposure and with the significant unilateral pleural thickening mesothelioma is the most likely diagnosis.

70096

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 62-year-old woman presents to the Emergency Department with a 3-day history of a productive cough, left-sided chest pain, fever and shortness of breath.

On examination, she is confused and febrile. Her BP is 89/60 mmHg, heart rate 102 bpm and respiratory rate 31 breaths per minute. Her oxygen saturation is 87%. She has left-sided upper lobe bronchial sounds on auscultation.

Investigations:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	123 g/l	115–155 g/l
White cell count (WCC)	15.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	178 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	133 mmol/l	135–145 mmol/l
Potassium (K^+)	3.2 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	112 μmol/l	50–120 μmol/l
Mean Corpuscular Volume (MCV)	89 fl	76–98 fl
Urea	8.1 mmol/l	2.5–6.5 mmol/l
C-Reactive Protein (CRP)	59 mg/l	< 10 mg/l

Her CXR is shown below:



What is the next most important step in this patient's immediate management?

Your answer was correct

A	Hypertonic saline solution
B	IV fluids
C	Non-rebreather oxygen mask for delivery of fraction of inspired oxygen of 60–80%
D	Potassium replacement
E	Start oral antibiotics

Explanation

C Non-rebreather oxygen mask for delivery of fraction of inspired oxygen of 60–80%

This patient has severe pneumonia, which has a mortality of approximately 10% in patients admitted to the hospital. Features of severity, according to the CURB-65 score, include: confusion, urea concentration greater than 7.0 mmol/l, respiratory rate over 30 breaths per minute, BP less than 90 mmHg systolic and age over 65 years. A score of greater than 2 indicates severe pneumonia. She would, of course, need IV antibiotics, fluids, oxygen and analgesia as rapidly as possible. In this situation, the most marked abnormality is the low O2 saturation, and this should therefore be addressed first. The CXR shows left upper lobe consolidation, which is in keeping with this diagnosis.

Image source: https://commons.wikimedia.org/wiki/File:Lobaerpneumonie.jpg

A Hypertonic saline solution

This is inappropriate as there is only mild hyponatraemia, which does not need to be corrected urgently, and standard crystalloid fluids could be used for resuscitation. Hypertonic saline can be used in patients with chronic hyponatraemia who are symptomatic and with sodium levels of less than 120 mmol/l.

B IV fluids

While IV fluids should be given very urgently, it is of greater importance to ensure that the patient has good oxygenation.

D Potassium replacement

Potassium supplements should be given either as part of fluid management or with PO supplementation, but are of no immediate urgency for this patient.

E Start oral antibiotics

This would be inappropriate as the patient is septic secondary to a chest infection and needs IV antibiotics. Sepsis is defined as meeting two or more of systemic inflammatory response syndrome (SIRS) criteria with a suspected source of infection. SIRS criteria include temperature outside of 36–38 °C, heart rate greater than 90 bpm, a respiratory rate greater than 20 breaths per minute or partial pressure of carbon dioxide less than 32 mmHg and WCC outside of 4–12/mm3. This patient has at least three of these criteria.

73461

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 56-year-old man presented to his GP with a 4-week history of feeling generally unwell. He described nasal congestion, with pain under his left eye, flu-like symptoms and lethargy. His GP made a diagnosis of sinusitis and gave him a course of antibiotics. He did not improve and developed breathlessness, cough and chest pain. He had previously been fit and his only past medical history was of an inguinal hernia repair 10 years previously. He worked in a shop. He smoked 10 cigarettes a day and drank about 10 units of alcohol a week. He was taking no regular medication.

On examination, he looked unwell and pale. Observations: respiratory rate 34/min, BP 140/85 mmHg, pulse 120 bpm, temperature 36.6 °C. He had crusting of his nasal septum. His JVP was not elevated, and heart sounds were normal. Auscultation of his chest revealed fine crackles bilaterally. His abdominal and neurological examinations were unremarkable.

Urinalysis:

Protein	++
Blood	++
Nitrites	-
Leucocytes	-
Bilirubin	-

Investigations reveal;

Investigation	Result	Normal Values
Haemoglobin (Hb)	81 g/l	135 - 175 g/l
White cell count (WCC)	6.3 × 10^9/l	4.0 - 11.0 × 10^9/l
Mean corpuscular volume (MCV)	81 fl	80 - 100 fl
Platelets (PLT)	510 × 10^9/l	150 - 400 × 10^9/l
Erythrocyte sedimentation rate (ESR)	114 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	45 mg/l	< 10 mg/l
Sodium (Na^+)	142 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.9 mmol/l	3.5 - 5.0 mmol/l
Urea	21 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	341 μmol/l	50 - 120 μmol/l
Bilirubin	18 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	31 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	191 u/l	30 - 150 u/l

Albumin	34 g/l	35 - 55 g/l

Spirometry:

FEV1	72% predicted
FVC	77% predicted
KCO	122% predicted

His chest X-ray is shown below.



What is the most likely diagnosis?

Your answer was incorrect

A Granulomatosis with polyangiitis (GPA)

B Microscopic polyangiitis

C Goodpasture's syndrome/anti-GBM disease

D Pulmonary oedema

E *Legionella pneumophila*

Explanation

A Granulomatosis with polyangiitis (GPA)

This is a question about pulmonary–renal syndromes (for a list see below).
The raised KCO and the CXR should make one think of pulmonary haemorrhage. This should narrow the diagnosis down to a vasculitis, Goodpasture syndrome and SLE.

This patient has Granulomatosis with polyangiitis (GPA). The clinical diagnosis is made by the triad of upper respiratory tract (nasal polyps and sinusitis), lower respiratory tract and renal involvement. It is a small-vessel vasculitis associated with granulomas.

The diagnosis of GPA is suggested by a positive c-ANCA with the presence of an antibody to proteinase-3. The chest radiograph usually shows single or multiple nodules, fixed infiltrates or cavities. Pleural effusions and pulmonary infiltrates may also been seen. Diagnosis is usually confirmed with a biopsy. If untreated the median survival is just five months. Treatment is with intravenous cyclophosphamide and prednisolone, which reduces mortality to approximately 10%. Plasmapheresis would be first-line treatment if the disease was primarily renal.

Causes of pulmonary–renal syndromes:

- Systemic diseases:
  Granulomatosis with polyangiitis
  Microscopic polyangiitis
  Goodpasture syndrome
  Systemic lupus erythematosus
  Polyarteritis nodosa
  Henoch–Schönlein purpura
  Churg–Strauss syndrome (renal involvement less common)
- Primary pulmonary disease:
  Legionella pneumonia and interstitial nephritis
  Bacterial pneumonia with renal compromise secondary to sepsis
- Others:
  Pulmonary oedema with acute kidney disease
  Uraemic pneumonitis
  Right-sided bacterial endocarditis – may cause pulmonary embolic lesions and glomerulonephritis
  Iatrogenic glomerulonephritis with ciprofloxacin, e.g. in patients given ciprofloxacin for cystic fibrosis.

Image source: https://upload.wikimedia.org/wikipedia/commons/7/77/Wegeners_Granulomatosis_CXR.jpg (https://upload.wikimedia.org/wikipedia/commons/7/77/Wegeners_Granulomatosis_CXR.jpg)

C Goodpasture’s syndrome/anti-GBM disease

Goodpasture syndrome can present with haemoptysis, although it is not associated with midline sinusitis. It’s important to differentiate from microscopic polyangiitis with granuloma formation, because only anti-GBM disease responds to plasma exchange.

B Microscopic polyangiitis

Microscopic polyangiitis is characterised by a very similar small-vessel vasculitis, but without granuloma formation and involvement of the upper respiratory tract.

D Pulmonary oedema

Pulmonary oedema is the least likely diagnosis, as the JVP is not elevated and the KCO would be reduced.

E *Legionella pneumophila*

Legionella pneumonia is unlikely as he is afebrile and there is no evidence of foreign travel or exposure. In addition, this does not have a predilection for the upper respiratory tract.

70088

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 40-year-old single male carpenter presents to casualty with a three-week history of increasing shortness of breath. He is a lifelong non-smoker. Temperature on admission is 38.2°C. On auscultation of his chest he has bilateral inspiratory crepitations. There are no signs of peripheral oedema, and he is thin with a BMI of 19.5. You ask him to walk the length of the ward holding a portable pulse oximeter, and note that his oxygen saturation begins to fall rapidly.

Arterial blood gases (on air) reveal the following:

Investigation	Result	Normal Value
pH	7.36	7.35–7.45
pO_2	8.0 kPa	10.5–13.5 kPa
pCO_2	5.1 kPa	4.6–6.0 kPa
Bicarbonate (HCO_3^-)	20.1 mmol/l	24–30 mmol/l

His chest x-ray is shown below.



Which of the following represents the most likely underlying diagnosis in this case?

Your answer was correct

A Community-acquired pneumonia

B Fibrosing alveolitis

C *Pneumocystis jirovecii* pneumonia (PCP)

D Pulmonary oedema

E Pulmonary embolism

Explanation

C *Pneumocystis jirovecii* pneumonia (PCP)

PCP is most commonly seen related to immunodeficiency in patients who are HIV-positive. The fact that he desaturates on a walk test is typical of this condition, as are the clinical signs, which are much more prominent than the subtle ground-glass appearance x-ray changes. Incidence is highest when the CD4 lymphocyte count falls below 100/mm^3. Diagnosis is via sputum to look for cysts of PCP or via bronchioalvelolar lavage fluid if sputum is negative. Treatment is with co-trimoxazole given either orally or IV; alternatives include pentamidine. After one episode of infection, patients should be maintained on prophylactic co-trimoxazole to prevent further acute episodes.

Image source: https://upload.wikimedia.org/wikipedia/commons/9/9f/X-ray_of_ground_glass_opacities_of_pneumocystis_pneumonia.jpg

A Community-acquired pneumonia

The subacute course of this patient’s disease, coupled with the patchy bilateral areas of consolidation, is inconsistent with common causes of community-acquired pneumonia such as *S. pneumoniae*.

B Fibrosing alveolitis

The time course of this patient’s symptoms would be considered too short for the onset of fibrosing alveolitis. The desaturation on exercise is also characteristic of PCP.

D Pulmonary oedema

Patchy consolidation and pyrexia are more consistent with infection versus cardiac failure. The lack of significant cardiac risk factors also counts against pulmonary oedema as the cause of symptoms seen here.

E Pulmonary embolism

Acute-onset pleuritic chest pain is absent from the history given here, and the patchy consolidation and pyrexia don’t fit with an acute PE.

6523

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A patient with known bronchiectasis grows fully sensitive *Pseudomonas aeruginosa* in their sputum for the first time. This is associated with a deterioration in his breathlessness.

Which of the following antibiotics would be most suitable as first-line treatment?

Your answer was correct

A IV ticarcillin plus IV gentamicin

B IV ceftazidime

C IV gentamicin

D IV colistin

E Nebulised Colistin

Explanation

B IV ceftazidime

Standard treatment is with a third-generation cephalosporin such as ceftazidime, meropenem or aztreonam, used in combination with gentamicin if required. *Pseudomonas aeruginosa* is resistant to amoxicillin.

A IV ticarcillin plus IV gentamicin

Ticarcillin is primarily used against urinary *Pseudomonas* infections rather than respiratory infections.

C IV gentamicin

IV gentamicin can be an adjunct to meropenem in *Pseudmonas* infection but unsuitable on its own. In urinary infections, gentamicin can be used as monotherapy but it is generally otherwise used in combination.

D IV colistin

Colistin is an effective antibiotic against most Gram-negative bacilli. It can be used as last-line treatment against *Pseudomonas*, *Klebsiella* and *Acinetobacter* infections in the presence of multi-drug resistance. It is generally not used due to its renal toxicity. This makes it inappropriate at this stage for this patient.

E Nebulised Colistin

Nebulised treatments, such as aminoglycosides and colistin, can be used for prevention of a flare-up in a patient colonised with *Pseudomonas aeruginosa,* and their use as prophylactic therapy has been shown to improve lung function in patients with cystic fibrosis, but this not adequate as a treatment for acute infection.

2321

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 35-year-old woman presents to her general practitioner complaining of breathlessness. The only history of note is that her mother died of a respiratory illness of unknown aetiology during her early 40s. She is a non-smoker who works as a secretary in a solicitor's office. On examination her blood pressure is 122/72 mmHg, with pulse 85/min and regular. There are scattered crackles more marked at the lung bases on auscultation of the chest. Routine bloods are normal, pulmonary function testing reveals a restrictive defect.



Which of the following diagnoses fits best with this clinical picture?

Your answer was incorrect

A Sarcoidosis

B Obliterative bronchiolitis

C Adenocarcinoma of the lung

D Alveolar microlithiasis

E Silicosis

Explanation

D Alveolar microlithiasis

This rare condition, which is sometimes familial with autosomal recessive inheritance, is associated with the deposition of diffuse calcified microliths in the alveolar space. In familial cases there is often a 2:1 female preponderance. The chest X-ray here shows fine micronodular calcification seen, predominantly in the lung bases and around the hilar. Computed tomography (CT) scan may confirm this micronodular calcification, and there is a progressive restrictive lung defect. Unfortunately, progressive calcium deposition leads to respiratory failure. No medical intervention has proved successful, although lung transplantation has been used in some cases. So much calcium is deposited that lungs examined at post-mortem have been reported to be so stiff as to require a saw to enable sectioning for examination.

Image source: https://radiopaedia.org/cases/alveolar-microlithiasis-2?lang=gb (https://radiopaedia.org/cases/alveolar-microlithiasis-2?lang=gb)

B Obliterative bronchiolitis

Obliterative bronchiolitis is associated bronchiolar inflammation and scarring that leads to fixed airway obstruction. It isn't associated with the micronodular calcification seen here.

A Sarcoidosis

Sarcoidosis is not associated with areas of micronodular calcification, but is associated with hilar lymphadenopathy.

C Adenocarcinoma of the lung

Adenocarcinoma of the lung is a peripheral lung tumour that arises in distal airways and isn't associated with the diffuse calcification seen here.

E Silicosis

Silicosis is associated with eggshell calcification rather than the fine micronodular calcification seen here, and occurs in patients exposed to mineral dust.

32410

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 41-year-old woman is admitted with a rash and shortness of breath.

On examination, she is tachycardic at 118 bpm and her oxygen saturations are 92% on air, while a painful purpuric rash is noted across both legs. She is visibly dyspnoeic and is complaining of coughing up blood.

She has no past medical history of note and takes no regular medication. She has three children and is not planning to have any more.

Investigations:

Haemoglobin (Hb)	121 g/l	(115–135 g/l)
White cell count (WCC)	16.1×10^9/l	($4–11 \times 10^9$/l)
Creatinine (Cr)	251 μmol/l	(50–120 μmol/l)
MPO antibodies	Negative	
PR3 antibodies	Positive	
First sputum sample	Negative	

Her chest X-ray is shown below.



The patient is started on prednisolone.

What is the most appropriate addition to the management of this patient?

Your answer was incorrect

A Azathioprine orally

B Pulsed intravenous (IV) cyclophosphamide

C Methotrexate orally

D No addition required

E IV rituximab

Explanation

B Pulsed intravenous (IV) cyclophosphamide

With a purpuric sounding rash, joint pain, evidence of possible pulmonary haemorrhage and positive PR3 antibodies, the diagnosis here is likely to be granulomatosis with polyangiitis. This is severe and here presents with end-organ damage to the lungs and kidneys. Therefore, cyclophosphamide is required to achieve remission. This is given IV and pulsed because of its potential for renal toxicity.

D No addition required

In granulomatosis with polyangiitis, oral steroids alone are insufficient, and guidelines suggest they must be accompanied by pulsed IV cyclophosphamide.

A Azathioprine orally

This patient has evidence of ANCA-positive vasculitis – specifically, granulomatosis with polyangiitis. These patients need to be started on pulsed cyclophosphamide to help achieve remission. After remission, cyclophosphamide may be replaced by azathioprine or methotrexate.

C Methotrexate orally

In patients with a diagnosis of granulomatosis with polyangiitis, without symptoms or evidence of organ damage, immunosuppression may not be required and methotrexate may be offered instead. However, there is clear evidence of end-organ damage in this case. If methotrexate was being considered as a treatment, female patients must be counselled regarding family planning, due to its teratogenicity.

E IV rituximab

In granulomatosis with polyangiitis, rituximab may be offered in place of cyclophosphamide if it is not tolerated, if the patient’s family is incomplete, if there is a history of uroepithelial malignancy or if the patient has already reached their maximum cumulative cyclophosphamide dose.

71777

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Guidance: Rituximab
nice.org.uk/guidance/TA308/chapter/1-guidance
(https://www.nice.org.uk/guidance/TA308/chapter/1-guidance)

A 56-year-old man comes to the clinic for review. He complains of excessive daytime sleepiness such that he falls asleep a number of times during the day, and is unable to hold down his job as a taxi driver. He has a history of hypertension for which he takes Ramipril, but is otherwise well. His wife tells you he snores for most of the night and often stops breathing. On examination his blood pressure (BP) is 158/88 mmHg, his pulse is 78 beats per minute (bpm) and his body mass index (BMI) is 38 kg/m^2. He has a collar size of 18.5.

Investigations reveal the following;

Hb	135 g/l
WCC	6.9 x10^9/l
PLT	188 x10^9/l
Na$^+$	137 mmol/l
K$^+$	4.3 mmol/l
Creatinine	108 micromol/l
Glucose	8.1 mmol/l (fasting)
Sleep study	>10 hypoxic episodes associated with apnoea during the overnight period

Which of the following is the best long-term intervention for this patient?

Your answer was incorrect

A Caffeine

B CPAP

C Modafinil

D Uvuloplasty

E Weight loss

Explanation

E Weight loss

The answer is Weight loss -

This patient has obstructive sleep apnoea (OSA), contributing to excessive day time sleepiness, and potentially worsening BP and glucose control because of significant sympathetic activation associated with the night time wakening. Given that fasting glucose is suggestive of a new diagnosis of type 2 diabetes mellitus, aggressive attempts at weight loss are the most appropriate intervention here.

B CPAP

CPAP is incorrect because it is a short-term fix for the apnoeic episodes, but does not impact significantly with respect to weight reduction.

A Caffeine

Caffeine, is incorrect because it does not impact on the underlying cause of his excessive sleepiness, and runs the risk of daytime sleep episodes becoming more unpredictable.

C Modafinil

Modafinil is incorrect, it is a treatment for narcolepsy / cataplexy, and does not contribute to weight loss.

D Uvuloplasty

Uvuloplasty is incorrect, it is primarily a treatment for snoring, not for sleep apnoea.

40410

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Obstructive Sleep Apnoea

Media

Expanded explanations

Flashcards

A 65-year-old man presents with a 6-week history of anorexia, malaise and breathlessness which he relates to rapidly worsening asthma. He takes NSAIDs for osteoarthritis of his knees, paroxetine and also inhalers for mild asthma. He was on lithium carbonate 2–5 years ago.

Examination reveals mild peripheral oedema, bilateral pleural effusions and a skin rash. He is hypertensive at 210/100 mmHg.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	70 g/l	135 - 175 g/l
Eosinophils	2.2 × 10^9/l	0.04 - 0.4 × 10^9/l
Serum IgG	8.0 g/l	6.0 - 13.0 g/l
Corrected calcium (Ca^{2+})	1.9 mmol/l	2.2 - 2.7 mmol/l
Serum IgA	1.2 g/l	0.8 - 3.0 g/l
Phosphate (PO_4^{3-})	2.5 mmol/l	1.1 - 1.45 mmol/l
Serum IgM	1.5 g/l	0.4 - 2.5 g/l
Urea	79.0 mmol/l	2.5 - 6.5 mmol/l
Serum IgE	200 u/l	< 120 u/l
Creatinine (Cr)	800 µmol/l	50 - 120 µmol/l
Potassium (K^+)	7.1 mmol/l	3.5 - 5.0 mmol/l
Urinalysis	Blood ++, protein ++	
Abdominal ultrasound scan (USS)	No obstruction, kidneys 11 and 12 cm	

The likely diagnosis is:

Your answer was incorrect

A Amyloidosis

B Chronic analgesic nephropathy

C Eosinophilic granulomatosis with polyangiitis (EGPA)

D Lithium nephrotoxicity

E Membranous glomerulonephritis

Explanation

C Eosinophilic granulomatosis with polyangiitis (EGPA)

This man has presented with biochemical evidence of kidney disease. The key to the diagnosis here is the significantly raised eosinophil count and raised serum IgE both of which are characteristic of EPGA. EPGA is characterised by asthma and blood eosinophilia together with an eosinophilic vasculitis. The initial phase of the disorder is one of asthma and allergic rhinitis, often followed by peripheral blood eosinophilia with eosinophilic tissue disease. The vasculitic phase that follows is life-threatening; however, it can often be treated effectively with immunosuppression. Renal disease consists of focal glomerulonephritis in the vasculitic phase with non-selective proteinuria, haematuria and red cell casts on microscopy.

B Chronic analgesic nephropathy

Chronic analgesic nephropathy is incorrect. Analgesic nephropathy will not present with blood and protein in the urine and there is no skin rash. There may be papillary necrosis with visible blood in the urine and abdominal pain or insidious renal impairment with no blood or protein in the urine.

A Amyloidosis

Amyloidosis is incorrect. Amyloidosis typically presents with a nephrotic picture. Presentation involves heavy proteinuria in the form of nephrotic syndrome and chronic kidney disease. There is fatigue, weight loss, bruising and easy bleeding (including the gastrointestinal tract). AL amyloidosis may affect other organs such as heart to produce cardiomegaly and liver with hepatomegaly.

D Lithium nephrotoxicity

Lithium nephrotoxicity is incorrect. Lithium nephrotoxicity is associated with nephrogenic diabetes insipidus and may be associated with tubule-intersitial fibrosis and tubular dilatation. The presentation would be gradual worsening of renal function over a period of years. There is no blood in the urine or skin rash.

E Membranous glomerulonephritis

Membranous glomerulonephritis is incorrect. Membranous gomerulonephritis is the most common cause of nephrotic syndrome in the adult population in the developed world. There would not be any blood in the urine or any skin rash. Progressive renal impairment with proteinuria is a common presentation. The diagnosis is by renal biopsy which would show glomerular basement membrane (GBM) thickening and ‘spikes’ of GBM extending around subepithelial deposits often best seen on silver stains. There is loss of podocyte foot processes with a variable degree of acute tubular necrosis and tubule-interstitial fibrosis.

70798

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 34-year-old HIV-positive man presented to hospital complaining of increasing shortness of breath and cough. He had noticed his vision was slightly worse over the last few weeks. He had prolonged admission 4 months previously with a chest complaint. He has been taking prophylactic co-trimoxazole. He is a current smoker with a 10- pack year history. On examination he looked unwell. Pulse was 110/min, respiratory rate 28/min, BP 95/65 mmHg and saturations 91% on air. Auscultation of his chest revealed fine crackles bilaterally. The CXR shows reticular shadowing throughout both lung fields.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	94 g/l	135 - 175 g/l
White cell count (WCC)	3.2×10^9/l	4.0 - 11.0×10^9/l
Platelets (PLT)	511×10^9/l	150 - 400×10^9/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
$CD4^+$ count	10/mm^3	430 - 1690 mm^3
Sodium (Na^+)	139 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Urea	6.1 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	111 µmol/l	50 - 120 µmol/l
Bilirubin	52 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	113 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	121 u/l	30 - 150 u/l
Albumin	34 u/l	35 - 55 u/l
Transfer factor for carbon monoxide (TLCO)	80% predicted	

What is the likely diagnosis?

Your answer was incorrect

A *Pneumocystis jirovecii* pneumonia

B CMV pneumonitis

C Nocardia infection

D Tuberculosis

E Streptococcal pneumonia

Explanation

B CMV pneumonitis

This patient has a cytomegalovirus (CMV) infection. CMV is one of the most frequent opportunistic infections in patients with advanced HIV disease. A CD4 count <50 cells/mm^3 carries high risk of the disease.

It can cause hepatitis, colitis, retinitis, pneumonitis, radiculopathy and encephalitis. The reduced visual acuity and abnormal liver function tests also are more in keeping with CMV.

In the immunosuppressed, serology may be of little value. Diagnosis is made by polymerase chain reaction (PCR) of serum. Histological staining of transbronchial biopsies may demonstrate the pathognomic 'owl's eye' cells. Other techniques include rapid culture methods such as DEAFF (detection of early antigen fluorescent foci) by monoclonal antibody.

Treatment is with intravenous ganciclovir.

E Streptococcal pneumonia

There is multi-organ involvement here, making streptococcal pneumonia less likely.

A *Pneumocystis jirovecii* pneumonia

Clinically and radiologically, CMV pneumonia mimics *P. jirovecii* pneumonia (PJP). However, this patient was on prophylactic co-trimoxazole, making PJP less likely.

C Nocardia infection

Nocardiosis can occur in immunocompromised patients, but CXR usually shows cavitation plus/minus pleural effusion. Skin abscesses may be present. The diffuse CXR shadowing combined with eye symptoms and deranged LFTs make CMV disease more likely.

D Tuberculosis

TB is more common in immunocompromised patients, and miliary TB may affect multiple organs, but the combination of diffuse CXR shadowing, deranged LFTs and eye symptoms in particular make CMV disease more likely.

70145

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 55-year-old woman who is asthmatic is admitted to the Emergency Department with wheeze and shortness of breath, although she is not yet showing signs of fatigue. She is given continued nebulised salbutamol, intravenous (IV) steroids and high-flow oxygen. She remains short of breath after these initial therapies. The Nursing Sister in charge asks whether giving IV magnesium is appropriate.

According to the British Thoracic Society (BTS) guidelines, which of the following would be a reason to give IV magnesium at this stage?

Your answer was incorrect

A A peak expiratory flow rate (PEFR) of 49% of the patient's best PEFR after initial therapy

B A repeat blood gas showing a low $P_a(CO_2)$ of 3 kPa

C Magnesium at the lower end of the normal range

D Oxygen saturation of 93% on air after initial therapy

E Tachycardia with a heart rate of 105 bpm

Explanation

A A peak expiratory flow rate (PEFR) of 49% of the patient's best PEFR after initial therapy

According to the British Thoracic Society (BTS) guidelines, 50% magnesium sulfate at 1.2–2 g (4–8 mmol) over 20 minutes can be given intravenous (IV) to a patient with severe asthma not responding to inhaled bronchodilator or who has severe life-threatening asthma. A peak expiratory flow rate (PEFR) of 49% of best PEFR grades this patient as having a severe attack and the fact that she continues to have wheeze and shortness of breath after initial therapy would justify giving intravenous (IV) magnesium therapy.

B A repeat blood gas showing a low $P_a(CO_2)$ of 3 kPa

Due to hyperventilation in asthma exacerbations, patients 'blow off' their carbon dioxide and, therefore, the level on the blood gas is often low. A rising carbon dioxide level is concerning due to the possibility that the patient is becoming fatigued and not ventilating efficiently. In this context, a low $P_a(CO_2)$ is, therefore, reassuring.

C Magnesium at the lower end of the normal range

Magnesium concentration is irrelevant with regard to the treatment of asthma. Supra-therapeutic levels are given to achieve a pharmacological response with reduced acetylcholine release and improved smooth muscle relaxation.

D Oxygen saturation of 93% on air after initial therapy

If there are life-threatening features of an acute asthma exacerbation, IV magnesium should be considered. An oxygen saturation of < 92% would count as one of these features and therefore, this patient does not have a life-threatening exacerbation on the basis of her oxygen saturation.

E Tachycardia with a heart rate of 105 bpm

A heart rate of > 110 bpm is an indicator of severe asthma and is more likely to drive the use of magnesium.

72953

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BTS Guidelines: Asthma
brit-thoracic.org.uk/quality-improvement/guidelines/asthma/
(https://www.brit-thoracic.org.uk/quality-improvement/guidelines/asthma/)

Asthma

Media

Expanded explanations

Flashcards

A 71-year-old patient undergoes radical surgery for an oesophageal adenocarcinoma with oesophagectomy, gastrectomy and splenectomy. Initially he makes a good recovery but you are asked to review him a few days later because he is increasingly short of breath. On examination BP is 142/72 mmHg and pulse 79/min and regular. There is reduced expansion of the left side of the chest and dullness to percussion over the left lung fields.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	109 g/l	135–175 g/l
White cell count (WCC)	7.9 × 10^9/l	4–11 × 10^9/l
Platelets	172 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5–5.0 mmol/l
Creatinine	105 μmol/l	50–120 μmol/l
Chest X-ray	Large left sided pleural effusion	
Diagnostic tap	milky coloured effusion, protein content 48g/l	

Which of the following investigations on the pleural fluid is most likely to contribute to confirming the underlying diagnosis?

Your answer was incorrect

A Triglycerides

B LDH

C Culture and sensitivity

D pH

E Glucose

Explanation

A Triglycerides

A triglyceride level greater than 1.2 mmol/l indicates a 99% chance that the fluid in the pleural space is chyle. This is likely to be a complication of surgery, with intra-operative damage to the duct or one of its tributaries. This can cause up to 4 l of fluid per day to leak into the pleural space, leading to the classic symptoms of a large pleural effusion. Initial management is conservative as the thoracic duct leak reduces spontaneously over a few days. If the effusion persists, imaging is needed, with referral for surgical ligation as indicated.

D pH

Testing the acidity of pleural fluid is used as a guide to treatment in para-pneumonic and malignant pleural effusions: a pH of <7.2 in the former indicates the need to drain, and <7.3 in the latter is associated with worse outcome. Neither is relevant in this clinical scenario.

B LDH

LDH is used to distinguish an exudate from a transudate, and is therefore utilised to elucidate the cause of pleural effusion when heart failure versus inflammation or infection needs to be delineated. It is often done as part of the routine tests on pleural fluid, but will not be the most useful in confirming the underlying diagnosis.

C Culture and sensitivity

This would identify the organism in an infective pleural effusion but would not aid the diagnosis here.

E Glucose

Glucose levels are of value in para-pneumonic effusions and identify causes such as tuberculosis or malignancy.

23330

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pleural Effusion

Media

Expanded explanations

Flashcards

A 69-year-old smoker presents for review. He has a 60-pack year history and has recently begun losing weight. In addition he has a chronic cough and is thirsty all the time. Past history of note includes two left carotid territory TIAs in the past year. Blood testing reveals mild hypercalcaemia, a left upper lobe lung mass is seen on chest X-ray and trans-bronchial biopsy confirms this to be a squamous cell carcinoma of the bronchus.

Investigations reveal the following:

Investigation	**Result**
Computed tomography (CT) scan	Left upper lobe mass, with maximal dimension of 5.2 cm. No evidence of lymph node, pleural or rib involvement
Spirometry	FEV_1 pre surgery less than 40% predicted (1.2 Litre)
Carotid ultrasound (US)	Left carotid 90% stenosis
Corrected calcium (Ca^{2+})	2.7 mmol/l

Which of the following represents the most appropriate management in this case?

Your answer was incorrect

A Refer for left pneumonectomy

B Refer for left upper lobectomy

C Refer for radiotherapy

D Refer for chemotherapy

E Refer for combined chemo- and radiotherapy

Explanation

E Refer for combined chemo- and radiotherapy

Unfortunately, this patient makes a very poor operative candidate due to his spirometry and carotid stenosis, which means he is at risk of not having sufficient residual lung function postoperatively, and of suffering a perioperative stroke. As such, radical radio- and chemotherapy is the intervention of choice.

C Refer for radiotherapy

Radiotherapy alone risks denying this patient access to chemotherapy for squamous cell carcinoma, which may significantly impact on his 2-year survival.

A Refer for left pneumonectomy

Pneumonectomy is likely not survivable for this patient, given that his FEV_1 pre-surgery is less than 40% predicted.

B Refer for left upper lobectomy

Although left upper lobectomy may preserve sufficient lung function in this patient, his carotid atherosclerosis puts him at high risk of a perioperative stroke.

D Refer for chemotherapy

Chemotherapy alone risks denying this patient access to the benefits of aggressive radiotherapy – for example, continuous hyperfractionated radiotherapy (CHART).

18548

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

The airway pressure graph of a patient receiving ventilator support is shown below.



What form of ventilatory support is this patient receiving?

Your answer was incorrect

A	Continuous positive airways pressure (CPAP)
B	Positive end-expiratory pressure (PEEP)
C	None
D	Intermittent mandatory ventilation (IMV)
E	Inverse ratio ventilation

Explanation

A Continuous positive airways pressure (CPAP)

CPAP generates continuous increased pressure, where the shape of the curve of inspiration and expiration stays the same but the whole curve is shifted upwards.

B Positive end-expiratory pressure (PEEP)

In PEEP, positive pressure is applied at the end of expiration and hence this portion of the curve remains positive for a more prolonged period, with a rapid fall-off at the end of the breath, versus normal respiration.

C None

In normal respiration the airway pressure returns to baseline at the beginning and end of each breath.

D Intermittent mandatory ventilation (IMV)

Intermittent mandatory ventilation intersperses ventilator-driven breathing with normal breathing, when the patient’s own respiratory effort is inadequate.

E Inverse ratio ventilation

Inverse ratio ventilation extends the inspiratory phase of breathing, leading to a greater AUC for the inspiratory phase of ventilation, versus the symmetrical curves seen here.

32451

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Ventilatory Support

Expanded explanations

A 25-year-old man presents with left-sided pleuritic chest pain. It had come on suddenly 3 days previously. Initially he thought it was secondary to his workout in the gym. However, since it did not improve and he noticed that he was slightly more breathless than usual, he went to the Emergency Department. He has no significant past medical history. He is a smoker of 20/day. On examination he looked well. He was a tall, thin man and was not short of breath at rest. Examination was unremarkable except for a clicking sound which was synchronous with the heart sounds. CXR was unremarkable.

Which of the following investigations may help with the diagnosis?

Your answer was incorrect

A ECG

B Echocardiogram

C Expiratory CXR

D Lateral decubitus CXR

E Repeat CXR

Explanation

D Lateral decubitus CXR

A click synchronous with the heart sounds is a recognised sign of a small left apical pneumothorax. He is a tall, thin man and a smoker, which are both risk factors for developing a pneumothorax. There is a strong association between pneumothoraces and smoking. The most recent British Thoracic Society (BTS) guidelines suggest that if the postero-anterior (PA) chest X-ray is normal and a small pneumothorax is suspected, a lateral decubitus chest X-ray provides added information in up to 14% of cases (see weblink below). For patients in the lateral decubitus position, air rises to the non-dependent lateral location. Only 5 mL of pleural air may be needed to detect pneumothorax in this position.

The treatment of a primary pneumothorax in a non-smoker, <50 years old, with no evidence of underlying lung disease, with a rim of air of <2 cm and no breathlessness is to discharge and follow up as an outpatient. Although this patient is a smoker there is nothing to suggest underlying lung disease, and it is likely that his apical pneumothorax is too small to safely aspirate. Patients should be given analgesia if required and clear instructions to return if their symptoms get any worse.

C Expiratory CXR

BTS guidelines recommend the use of lateral decubitus CXR. Whilst there is a thought process that expiration results in the lungs reducing in volume, and theoretically improving visability of pneumothorax, they are not recommended in BTS guidelines.

A ECG

BTS guidelines recommend the use of lateral decubitus CXR. ECG may be abnormal in pneumothorax, which may include abnormal left axis deviation and right axis deviation. There may be QRS abnormalities as well.

B Echocardiogram

BTS guidelines recommend the use of lateral decubitus CXR. Ultrasound may be useful in diagnosing traumatic pneumothorax, and the cardiac function may be affected by expanding pneumothorax.

E Repeat CXR

BTS guidelines recommend the use of lateral decubitus CXR. Repeating a CXR will not help as the CXR may look the same.

70119

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total:	301
Responses - % Correct:	27%

External Links

BTS Pleural Disease Guideline 2010

brit-thoracic.org.uk/document-library/clinical-information/pleural-disease/pleural-disease-guidelines-2010/appendix-3-spontaneous-pneum...

(https://www.brit-thoracic.org.uk/document-library/clinical-information/pleural-disease/pleural-disease-guidelines-2010/appendix-3-spontaneous-pneumothorax-poster-pleural-disease-guideline/)

Pneumothorax

Media

Expanded explanations

Flashcards

A 21-year-old man presents with a 3-month history of progressive wheeze and dyspnoea. Over the last few weeks, he has been coughing up thick, rubbery ‘plugs’ of mucus. There is no haemoptysis. He has a background of mild asthma, which was well controlled until a few months ago, and eczema. He has had recurrent chest infections recently, requiring several courses of antibiotics from his general practitioner, and he attributes this to the damp autumn weather. He also has a feeling of general malaise. He had been more acutely short of breath for the last few days.

On examination he was dyspnoeic at rest, but not cyanosed. His temperature was 37.8 °C, pulse 100/min, blood pressure 132/86 mmHg, $S_a(O_2)$ 94% on air, respiratory rate 22/min. His heart sounds were normal. On auscultation of his chest, he had widespread expiratory wheeze, with some coarse crepitations in the mid-zones and reduced air entry in the right lower zone. Abdominal and neurological examinations were unremarkable.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	150 g/l	135–175 g/l
White cell count (WCC)	13.1×10^9/l	$4.0–11.0 \times 10^9$/l
Neutrophils	8.5×10^9/l	$2.5–7.58 \times 10^9$/l
Lymphocytes	2.2×10^9/	$1.5–3.5 \times 10^9$/l
Monocytes	0.2×10^9/	$0.0–0.8 \times 10^9$/l
Eosinophils	2.1×10^9/	$0.04–0.4 \times 10^9$/l
Basophils	0.1×10^9/	$< 0.1 \times 10^9$/l
Platelets (PLT)	390×10^9/	$150–400 \times 10^9$/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Bilirubin	14 µmol/l	2–17 µmol/l
Potassium (K^+)	4.2 mmol/l	3.5–5.0 mmol/l
Alkaline phosphatase (ALP)	90 IU/l	30–130 IU/l
Urea	5.2 mmol/l	2.5–6.5 mmol/l
Alanine aminotransferase (ALT)	32 IU/l	5–30 IU/l
Creatinine (Cr)	100 µmol/l	50–120 µmol/l
Gamma glutamyl-transferase (γGT)	30 IU/l	5–30 IU/l
Chest X-ray (CXR)	Patchy infiltrates throughout both lung fields, particularly affecting the upper lobes	

What is the most likely diagnosis?

Your answer was incorrect

A Cystic fibrosis

B Invasive pulmonary aspergillosis

C Cryptogenic eosinophilic pneumonia

D Churg–Strauss syndrome

E Allergic bronchopulmonary aspergillosis (ABPA)

Explanation

E Allergic bronchopulmonary aspergillosis (ABPA)

ABPA is due to endobronchial colonisation by the fungus *Aspergillus fumigatus*. It tends to present in early adult life against a background of worsening asthma. There is initially bronchospasm, which can progress to proximal bronchiectasis and occasionally upper zone lung fibrosis, similar to TB.

Exacerbations are more common in the autumn and winter months. Patients often get recurrent episodes of eosinophilic pneumonia, and fleeting infiltrates can often be seen on the chest X-ray. Mucus plugging can sometimes cause lobar collapse.

The diagnosis is made by skin-prick testing, which causes an immediate hypersensitivity reaction. Serum precipitins are present in 90% of cases and there is a high peripheral eosinophil count.

Treatment is with oral prednisolone, which usually results in a rapid clinical and radiological improvement. Sometimes maintenance steroids are needed to prevent relapse.

A number of studies have suggested a possible role for itraconazole therapy, e.g. Wark et al., *Journal of Clinical Immunology* 2003, and this may be a useful adjunctive therapy in some patients.

C Cryptogenic eosinophilic pneumonia

This is characterised by accumulation of eosinophils in the interstitium and alveolar spaces without provocation. Strongly associated with asthma and atopy, there is gradual onset in the 4th or 5th decades of life. There is commonly dyspnoea, productive cough and fever with wheeze and crepitations. For this diagnosis, the eosinophil count would be much higher.

A Cystic fibrosis

Cystic fibrosis (CF) is a common autosomal recessive disorder of the CF trans-membrane conductance regular. This is a chloride channel, and when it does not function properly the composition of the mucus electrolytes changes, making it thicker. This causes chronic pulmonary infections and bronchiectasis in the lungs, and pancreatic insufficiency, gallstones, cirrhosis and male infertility. History is with recurrent infections or gastrointestinal signs such as steatorrhoea. Management is with physiotherapy for postural drainage, mucolytics and pancreatic enzyme replacements. First presentation at the age of 21 would be considered very unusual.

B Invasive pulmonary aspergillosis

Invasive aspergillosis occurs in immunocompromised patients or following the use of broad-spectrum antibiotics. Diagnosis can be very difficult. Sputum cultures of alveolar lavage may demonstrate *Aspergillus* on microscopy, but often biopsy is needed. Early CT scan of the chest or serial measurements of galactomannan may be helpful. Treatment is with voriconazole or miconazole. There is 30% mortality. The absence of risk factors makes this an unlikely diagnosis.

D Churg–Strauss syndrome

The correct name for this disorder is now eosinophilic granulomatosis with polyangitis (EGPA), a disorder of rhinosinusitis, asthma and eosinophilia with vasculitis. Patients normally develop problems in their 20s to 30s with atopy, rhinitis and asthma, followed later by eosinophilic infiltration of the lungs and vasculitis. The moderate elevation only in eosinophil count, coupled with the absence of other features of vasculitis, counts against a diagnosis of EGPA.

9156

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A 64-year-old man was referred to the clinic with insomnia. He complained of falling asleep during the day and having an early morning headache. He had a past history of a cholecystectomy and gout. He took no other regular medication. Examination was unremarkable. His weight and height were 120 kg and 175 cm, respectively.

Investigations:

Investigation	Result	Normal Values
Weight	120 kg	
Height	175 cm	
Haemoglobin (Hb)	161 g/l	135 - 175 g/l
White cell count (WCC)	6.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	94 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	101 fl	80 - 100 fl
Thyroid-stimulating hormone (TSH)	1.2 mu/l	0.4 - 5.0 mu/l

What underlying cause should be suspected in this patient?

Your answer was incorrect

A Acromegaly

B Alcohol excess

C Hypothyroidism

D Cushing’s disease

E Enlarged tonsils

Explanation

B Alcohol excess

This patient has obstructive sleep apnoea (OSA), the symptoms of which include insomnia, daytime somnolence, morning headache and obesity. Other symptoms include poor concentration during the day and partners may describe snoring followed by apnoeic episodes. For screening of OSA, the Epworth score can be helpful which is a measure of daytime somnolence.

Diagnosis is made using sleep studies, which would show apnoeic/hypopnoeic episodes associated with desaturations, an increase in heart rate and arousal from sleep. Treatment will depend on severity of symptoms and desaturations. In most cases, weight loss is advisable.

In mild OSA syndrome, a mandibular advancement device may control symptoms in mild to severe OSA syndrome nocturnal non-invasive ventilation (NIV) using continuous positive airway pressure (CPAP). Early morning headaches can be caused by a transient hypercapnia. This normal corrects to normal when the patient is awake. If a patient has hypercapnia in waking hours (and not just on an early morning blood gas) then overlap syndrome with obesity hypoventilation syndrome should be considered, although severe OSA may cause persistent hypercapnia in some case. Patients with persistent hypercapnia should be considered for bilevel positive airway pressure (eg BiPAP).

It is important to exclude underlying causes. This patient has a history of gout which, together with the raised MCV and low platelets, suggests alcohol consumption, which can precipitate the problem. Hypothyroidism, acromegaly and sedating drugs also need to be excluded. Retrognathia can cause OSA and large tonsils may obstruct the airway.

C Hypothyroidism

The patient’s TSH is not raised, making this an unlikely diagnosis.

A Acromegaly

Acromegaly is associated with OSA but there are no features in the history to suggest this over the more likely diagnosis of alcohol excess.

D Cushing’s disease

Cushing’s disease may be associated with OSA, but there are no features in the history or blood tests to suggest this as the more likely diagnosis versus alcohol excess.

E Enlarged tonsils

Enlarged tonsils may be associated with OSA, but the history and blood tests are more strongly suggestive of alcohol excess.

70141

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Obstructive Sleep Apnoea

Media

Expanded explanations

Flashcards

A 71-year-old man presents to his GP along with his wife. He has severe problems with sleeping at night, with episodes where he snores, stops breathing, is restless and often disturbs his wife from sleep. During the day his family complain that he often falls asleep watching the television, after his dinner or even whilst talking to his wife. He is a smoker of 20 cigarettes per day, has hypertension and suffered a TIA 3 years earlier. Medication includes ramipril 10 mg, simvastatin 40 mg and aspirin 75 mg. On examination his BMI is 32, his BP is 150/90 mmHg and he has signs of right heart failure.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	140 g/l	135–175 g/l
White cell count (WCC)	5.8 × 10^9/l	4.0–11.0 x 10^9/l
Platelets (PLT)	220 × 10^9/l	150–400 x 10^9/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	140 μmol/l	50–120 μmol/l

Which of the following is the most likely diagnosis?

Your answer was correct

A Central sleep apnoea

B Nocturnal epilepsy

C Obstructive sleep apnoea

D Multiple TIAs

E Narcolepsy

Explanation

C Obstructive sleep apnoea

This patient is obese, with hypertension and right heart failure. He has significant symptoms of nocturnal airway obstruction and daytime sleepiness, and this constellation of symptoms and signs fits with a diagnosis of obstructive sleep apnoea. Management of the condition involves aggressive weight reduction and the use of nocturnal CPAP to relieve nocturnal airway obstruction. Clearly this gentleman should also receive very strong advice to stop smoking and should be advised not to drive.

A Central sleep apnoea

Central sleep apnoea is usually seen in patients with significant neurological impairment, and the fact this patient is a smoker and obese is a much stronger pointer to obstructive sleep apnoea.

B Nocturnal epilepsy

Nocturnal epilepsy is more likely to be associated with tonic–clonic movements, automatisms or sleep-walking, rather than the clinical picture seen here.

D Multiple TIAs

There are no focal features to suggest TIAs; the recurrent snoring and cessation of breathing is much more suggestive of obstructive sleep apnoea.

E Narcolepsy

Narcolepsy is associated with excessive daytime sleepiness, cataplexy, hypnagogic hallucinations and sleep paralysis. The snoring and periods of apnoea are not consistent with a diagnosis of narcolepsy.

18628

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 50-year-old woman undergoes routine cholecystectomy. Her preoperative full blood count was normal and her operative progress was unremarkable other than a small postoperative bleed that settled with simple pressure to the wound. You are asked to see her, as she has become confused. She has no previous medical history of note and works as a computer programmer.

On examination, she is disorientated in time and place. Temperature 37.9 °C, pulse 110 beats per minute (bpm), blood pressure 110/70 mmHg, respiratory rate 24 breaths per minute. Heart sounds are normal and jugular venous pressure is not raised. Chest reveals some bronchial breathing on the right lower and middle zones. Her O_2 saturation is 91% on air. Abdominal examination reveals mild tenderness around the operative scar but is otherwise normal.

Investigations reveal the following:

Investigation	Result	Normal Value
Haemoglobin (Hb)	95 g/l	115–155 g/l
White cell count (WCC)	12.3×10^9/l	4.0–11.0 x 10^9/l
Platelets (PLT)	790×10^9/l	150–400 x 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	197 µmol/l	50–120 µmol/l
C-reactive Protein (CRP)	272 mg/l	< 10 mg/l

A blood film shows a dramatic increase in platelets with many enlarged forms. There is an increase in left-shifted granulocytes including metamyelocytes, myelocytes and promyelocytes.

In addition to basic resuscitation, what action would you take?

Your answer was incorrect

A Aspirin 300 mg with ceftazidime 1 g twice daily intravenously

B Aspirin 75 mg plus hydroxyurea 1 g twice daily

C Apheresis to reduce the platelet count urgently

D Anticoagulation with heparin to therapeutic levels

E Broad spectrum antibiotic intravenously

Explanation

E Broad spectrum antibiotic intravenously

This patient has developed postoperative pneumonia, which is making her confused secondary to hypoxia. The platelet count is raised secondary to infection – the normal preoperative count makes a primary haematological diagnosis unlikely in the extreme; the left shift is also secondary to infection. First-line therapy in this situation depends upon the presence or absence of risk factors for multi-drug resistant pathogens, knowledge of the local predominant pathogens and the patient's prior microbiology results. Additional consideration for treating anaerobic bacteria must be made if aspiration pneumonia is a possibility.

C Apheresis to reduce the platelet count urgently

Given that the platelet count increase is reactive as a result of ongoing infection, there is no indication for apheresis to take place.

A Aspirin 300 mg with ceftazidime 1 g twice daily intravenously

Aspirin is not required in this situation given that the increased platelet count is reactive; ceftazidime is an appropriate antibiotic choice where *Pseudomonas* is suspected.

B Aspirin 75 mg plus hydroxyurea 1 g twice daily

Hydroxyurea with aspirin is standard treatment for essential thrombocytosis, a clonal stem cell disorder associated with increased risk of thrombosis and haemorrhage.

D Anticoagulation with heparin to therapeutic levels

Anticoagulation with heparin to therapeutic levels isn’t indicated, and may in the postoperative period increase the risk of haemorrhage. Prophylactic dosing with low-molecular weight heparin may, however, be used here.

6385

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 63-year-old man comes to the Respiratory Clinic for review. He has chronic obstructive pulmonary disease (COPD), with a 40-pack-year smoking history, having stopped smoking only two years earlier. Despite taking salbutamol inhalers four to five times per day, he is still significantly short of breath, has a chronic dry cough, can walk only 30–40 m and only climbs the stairs once a day.

Examination reveals a blood pressure of 142/72 mmHg and a pulse of 80 bpm and regular. His chest is hyperexpanded; air entry is poor, and there is fine wheeze on auscultation. Lung function testing showed a forced expiratory volume in one second (FEV$_1$) of 30% of predicted, with no evidence of reversibility.

According to the National Institute of Health and Care Excellence (NICE), which of the following is the most appropriate next step?

Your answer was incorrect

A Add a combination of a long-acting muscarinic antagonist (LAMA) + long-acting beta-2 agonist (LABA) inhaler

B Add a combination of a LABA + inhaled corticosteroid (ICS) inhaler

C Add a combination of LAMA + LABA + ICS

D Add Salmeterol

E Add tiotropium

Explanation

A Add a combination of a long-acting muscarinic antagonist (LAMA) + long-acting beta-2 agonist (LABA) inhaler

Recently updated NICE guidelines suggest using a LAMA + LABA inhaler as second-line treatment if there is no evidence of reversibility on lung function testing that would suggest steroid responsiveness.

B Add a combination of a LABA + inhaled corticosteroid (ICS) inhaler

The addition of a LABA + ICS inhaler would be appropriate if the patient had asthmatic features or signs suggesting steroid responsiveness. Importantly, in this case, the patient has no evidence of reversibility on lung function testing, which suggests that this patient has no asthmatic features. Other tests that could be used to check steroid responsiveness are a high blood eosinophil count, substantial variation in FEV$_1$ over time (at least 400 ml) or substantial diurnal variation in peak expiratory flow (at least 20%).

C Add a combination of LAMA + LABA + ICS

This option would be an appropriate third-line therapy to consider if the patient fails to improve with the combination of LAMA + LABA therapy or if they have one to two severe exacerbations within a 12-month period.

D Add Salmeterol

This patient has significant airway obstruction, which is evidenced by an FEV_1 of 30% and his significant symptoms of breathlessness. He has already been established on a short-acting beta-2 agonist, which has failed to control his symptoms. Guidelines suggest that this should no longer be used second line as a monotherapy and should instead be used in combination.

E Add tiotropium

NICE guidelines now suggest that tiotropium should be used second line in combination with a LABA or an ICS, depending on the presence of steroid responsiveness, and not as a monotherapy as previously advised. 72027

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total:	301
Responses - % Correct:	27%

External Links

National Institute of Health and Care Excellence. 2018. Chronic obstructive pulmonary disease in over 16s: diagnosis and...
nice.org.uk/guidance/ng115
(https://www.nice.org.uk/guidance/ng115)

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 67-year-old man presents to the Emergency Department with a 3-day history of a productive cough. He has long-standing chronic obstructive pulmonary disease (COPD), has had six admissions over the past 18 months because of exacerbations, has home oxygen and takes nebulisers (fluticasone/salmeterol and tiotropium). He is only able to walk 2–3 yards to the toilet.

On examination, he has a respiratory rate of 15 breaths per minute. He mumbles incoherently after a vigorous sternal rub. His BP is 124/82 mmHg, and his heart rate is 85 bpm and regular. There is polyphonic wheeze on auscultation of the chest and ankle swelling consistent with right heart failure. He is receiving 60% oxygen (O_2) via a Venturi mask.

Arterial blood gas (ABG) analysis on arrival:

Investigations	Results	Normal Values
pH	7.22	7.35–7.45
Bicarbonate (HCO_3^-)	34 mmol/l	24–30 mmol/l
Partial pressure of carbon dioxide$p$(CO_2)	12 kPa	4.6–6.0 kPa
Partial pressure of oxygen (pO_2)	10.8 kPa	10.5–13.5 kPa

His CXR is shown below:



Which of the following is the next most appropriate intervention?

Your answer was incorrect

A 28% O_2

B Continuous positive airway pressure

C Doxapram

D Intubation and ventilation

E Non-invasive positive pressure ventilation

Explanation

A 28% O_2

This patient is adequately oxygenated but is clearly drowsy and acidotic because of marked carbon dioxide (CO_2) retention. The CXR shows subtle findings in keeping with chronic obstructive pulmonary disease (COPD), minor hyper expansion and diaphragmatic flattening. The most appropriate next step is to reduce the inspired

O_2 and recheck the ABG results. It is likely the CO_2 level will reduce significantly, accompanied by a small reduction in the O_2 level and a consequent improvement in consciousness.

Image source: https://commons.wikimedia.org/wiki/File:Chest_X-ray_-_posteroanterior_view.jpg

C Doxapram

Doxapram is a centrally-acting respiratory stimulant. It was extensively used in the management of COPD before the advent of NIPPV in patients for whom invasive ventilation was undesirable.

B Continuous positive airway pressure

Continuous positive airway pressure is more commonly used as a treatment for patients with sleep apnea. In this situation, if consciousness does not improve with decreased inspired O_2. Nasal intermittent positive pressure ventilation (NIPPV) is preferred.

D Intubation and ventilation

Invasive ventilation is undesirable in patients with end-stage COPD if it can be avoided because of the difficulties in weaning patients. Here a number of stages before this are necessary, including turning down the inspired oxygen and a trial of NIPPV.

E Non-invasive positive pressure ventilation

NIPPV is the obvious next step if hypercapnia does not rapidly improve once inspired O_2 is reduced, although ideally, the pH should not be allowed to fall much further before NIPPV is considered.

39520

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 55-year-old woman presents with a 6-month history of recurrent chest infections. She had had a few episodes of wheeze and haemoptysis in the last few weeks. She also noticed intermittent hot flushes, which her GP had put down to the menopause. These could occur at any time but on occasions were precipitated by alcohol. She had no significant past medical history and she had never smoked.

On examination, she was anxious; examination of her chest revealed reduced expansion on the right side. Cardiovascular examination revealed a pansystolic murmur. Blood results were unremarkable.

Her chest X-ray is shown below.



What is the treatment of choice?

Your answer was incorrect

A Surgery

B Steroid inhaler

C Intravenous antibiotics

D CHOP chemotherapy

E Alpha-adrenergic blockade, e.g. Phenoxybenzamine

Explanation

A Surgery

The treatment of choice for bronchial carcinoid is surgical resection, as lesions are vascular and obstructive, leading to potential significant haemoptysis and/or pneumonia.

The disease commonly presents between the ages of 50 and 70 and is unrelated to smoking. Bronchial carcinoid tumours are the most indolent form of a spectrum of neuroendocrine tumours of the lung that include small cell lung cancer as the most malignant. They are thought to arise from Kulchitsky cells in the bronchial mucosa. They usually secrete serotonin and arise from the large bronchi. Patients may present with a mass, which is usually centrally located (as in this case - CXR shows partially obstructive an endobronchial mass involving the bifurcation of the left main bronchus), or with recurrent chest infections, haemoptysis, chest pain and wheeze. Flushing (which may be precipitated by alcohol, food ingestion, stress or emotion) and diarrhoea may occur. However, patients are commonly asymptomatic and the tumour is found incidentally.

Carcinoid syndrome occurs when there are secondary tumours in the liver that release serotonin into the systemic circulation. However, symptoms of carcinoid syndrome may rarely occur in bronchial carcinoid in the absence of metastases, as the bronchial tree drains directly into the systemic circulation.

Diagnosis is by detection of 5-hydroxyindoleacetic acid (the metabolite of serotonin) in the urine, and bronchial carcinoid is usually seen and diagnosed by bronchoscopy and biopsy.

Pellagra may occur due to tumour uptake of tryptophan, the precursor of nicotinic acid.

Image source: https://radiopaedia.org/cases/endobronchial-carcinoid-tumor?lang=gb (https://radiopaedia.org/cases/endobronchial-carcinoid-tumor?lang=gb)

C Intravenous antibiotics

This is an appropriate treatment for pulmonary infection, which is not suggested here by the chronicity and periodicity of symptoms.

B Steroid inhaler

This is an appropriate treatment for asthma. In this case a steroid inhaler won’t impact on bronchial hyper-reactivity and won’t prevent/reduce the risk of lobar infection due to bronchial obstruction.

D CHOP chemotherapy

This is an appropriate treatment for lymphoma, which is less likely given the symptoms of intermittent flushing.

E Alpha-adrenergic blockade, e.g. Phenoxybenzamine

This is an appropriate treatment for phaeochromocytoma, where patients should be subject to complete alpha blockade prior to surgery.

70104

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 54 year old male with a history of alcohol excess presents to A+E with cough, dyspnoea and drenching night sweats. Following investigation, the diagnosis of TB is confirmed, and he is commenced on directly observed quadruple anti-TB combination treatment. Prior to beginning treatment, he attends the clinic for discussion of the therapy and potential side effects to be aware of.

Which of the following listed options is a possible side effect of anti-tuberculous therapy?

Your answer was incorrect

A Hepatic obstruction

B Haematuria

C Vitamin A deficiency

D Cortical visual loss

E Peripheral neuropathy

Explanation

E Peripheral neuropathy

Isoniazid can cause a peripheral neuropathy, particularly if used in patients with malnourishment, HIV or alcoholism, or other patients at risk of peripheral neuropathy.

Among of the main side effects to be aware of with anti-TB therapy are: peripheral neuropathy with isoniazid; orange-red staining of secretions with rifampicin; and optic neuritis with ethambutol. Rifampicin, isoniazid and pyrazinamide are all associated with GI upset and hepatitis.

B Haematuria

Rifampicin is known to cause orange-red discoloration of secretions such as urine, sweat and tears, but is not associated with haematuria.

A Hepatic obstruction

Anti-TB therapy is more commonly associated with drug-related hepatitis than obstruction.

C Vitamin A deficiency

Rifampicin and isoniazid can cause a relative vitamin D deficiency, probably already present in the patient but unmasked by the anti-TB therapy, but anti-TB therapy is not associated with vitamin A deficiency.

D Cortical visual loss

Rifampicin causes an orange discolouration of the urine. Ethambutol tends to cause colour blindness, restriction of visual fields and loss of visual acuity because of optic neuropathy, but not cortical blindness.

7172

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 16-year-old boy presents to the Outpatients Department having coughed up a small amount of bright-red blood. He is mildly short of breath and feels ‘wheezy’. His mother says he has always been a ‘chesty’ child. He was teased at school as he has always been the smallest in his class. His brother and sister were fit and well. He had a dog at home.

On examination he looked well and was not breathless at rest. He had clubbing. He had nasal polyps and auscultation of his chest revealed coarse crackles throughout both lung fields. His abdomen was distended but soft.

Urinalysis reveals the following:

Investigation	Result
Protein	+
Glucose	++
Nitrites	negative
Leucocytes	negative

Investigations reveal the following:

Investigation	Result	Normal values
Haemoglobin (Hb)	110 g/l	135–175 g/l
White cell count (WCC)	12.1 × 10^9/l	4.0–11.0 × 10^9/l
Neutrophils	91%	
Lymphocytes	8%	
Monocytes	0.4%	
Mean corpuscular volume (MCV)	84 fl	80–100 fl
Platelets (PLT)	450 × 10^9/l	150–400 × 10^9/l
Erythrocyte sedimentation rate (ESR)	35 mm/hour	1–20 mm/hour
C-reactive protein (CRP)	40 mg/l	< 10 mg/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5–5.0 mmol/l
Urea	6.4 mmol/l	2.5–6.5 mmol/l
Creatinine (Cr)	62 μmol/l	50–120 μmol/l
Bilirubin	31 μmol/l	1–22 μmol/l

Aspartate aminotransferase (AST)	46 U/l	10–40 U/l
Alkaline phosphatase (ALP)	221 U/l	30–150 U/l
Albumin	38 g/l	35–55 g/l
Protein	74 g/l	60–83 g/l
Magnesium (Mg^{2+})	0.64 mmol/l	0.6–1.1 mmol/l
Calcium (Ca^{2+})	2.1 mmol/l	2.2–2.7 mmol/l
Phosphate (PO_4^{3-})	0.75 mmol/l	1.1–1.45 mmol/l

Spirometry reveals the following:

Forced expiratory volume (FEV_1)	60% predicted
Forced vital capacity (FVC)	80% predicted
Peak expiratory flow rate (PEF)	350 l/min
Carbon monoxide transfer coefficient (KCO)	70% predicted

His Chest X-ray (CXR) is displayed below;



Which pathogen is most likely to colonise this patient?

Your answer was correct

A Streptococcus pneumoniae

B Haemophilus influenzae

C Aspergillus fumigatus

D Pseudomonas aeruginosa

E Mycobacterium tuberculosis

Explanation

D Pseudomonas aeruginosa

The answer is Pseudomonas aeruginosa -

This boy has cystic fibrosis. He has had recurrent chest infections secondary to bronchiectasis (confirmed by examination findings and the chest X-ray, also note long line inserted via left antecubital fossa), he is small for his age, has nasal polyps (which occur in one third of patients with CF) and clubbing. Pulmonary function tests show airflow obstruction He has mildly deranged liver function tests – secondary to obstruction of biliary ductules in the liver. This can eventually lead to cirrhosis. He has glucose in his urine – pancreatic islet cells are destroyed as the pancreas becomes fibrotic resulting in CF related diabetes in 30% of patients by adulthood.

The predominant bacteria causing infection is initially Staphylococcus aureus which declines in early adulthood (see external link). By the time patients are in their early teenage years, Pseudomonas aeruginosa is the second most common organism to colonise the patient before becoming the most common in early adulthood. As Staphylococcus aureus is not an option here, Pseudomonas aeruginosa is the next logical choice.

Male cystic fibrosis patients are infertile due to failure of the development of the vas deferens and epididymis.

Treatment is with pancreatic supplements, prophylactic inhaled or nebulised antibiotics, strict glucose control, mucolytics/DNAase and regular physiotherapy with postural drainage. Prophylactic Azithromycin is used for its immunomodulatory effects. Non-invasive ventilation may be used in patients with chronic type 2 respiratory failure as a bridge to lung transplantation Lung transplantation is considered as a treatment option once there is severe respiratory compromise.

A Streptococcus pneumoniae

Streptococcus pneumoniae is incorrect, it is not commonly seen in CF.

B Haemophilus influenzae

Haemophilus influenzae is incorrect, it is more common in early childhood in CF. Until age 9 or 10 it is the second most common pathogen, behind Staphylococcus aureus, and thereafter declines in incidence. Pseudomonas aeruginosa becomes the second most common pathogen at this point before becoming the dominant strain in early adulthood.

C Aspergillus fumigatus

Aspergillus fumigatus is incorrect, it is commonly found in sputum cultures of individuals with CF but is less likely to be found consistently than Pseudomonas.

E Mycobacterium tuberculosis

Isolation of Mycobacterium tuberculosis is incorrect, this is very uncommon in CF.

70087

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total: 301

Responses - % Correct: 27%

External Links

Prevalence of Respiratory Microorganisms by Age Cohort, 2019
cysticfibrosis.org.uk/sites/default/files/2020-12/2019%20Registry%20Annual%20Data%20report_Sep%202020.pdf
(https://www.cysticfibrosis.org.uk/sites/default/files/2020-12/2019%20Registry%20Annual%20Data%20report_Sep%202020.pdf)

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 49-year-old nurse comes to the surgery. She is a smoker of 20 cigarettes per day and is complaining of increased shortness of breath, a chronic cough and wheezing. On examination she smells strongly of cigarettes. You can hear coarse wheezing and occasional crackles on auscultation.

Which of the following is most useful in determining appropriate long-term management of her condition?

Your answer was incorrect

A Chest X-ray

B Alpha-1-antitrypsin deficiency testing

C CT thorax

D Bronchoscopy

E Spirometry with reversibility

Explanation

E Spirometry with reversibility

This woman has evidence of chronic obstructive pulmonary disease (COPD), but up to 30% of patients fit the definition of reversibility (>15% improvement in FEV_1 post-bronchodilator). In patients who are reversible, there is good evidence for treating with inhaled corticosteroids, as in asthma. Spirometry also correlates well with long-term mortality, and can be helpful in persuading patients to give up smoking.

C CT thorax

CT thorax is useful for elucidating the underlying cause of restrictive lung disease. In the event that lung function testing reveals a restrictive lung defect, CT thorax may be a more important part of the work-up.

A Chest X-ray

Although a chest X-ray may show hyper-expansion consistent, for example, with emphysema and alternative diagnoses such as lung cancer, it won’t establish response to therapeutic intervention.

B Alpha-1-antitrypsin deficiency testing

Given this patient's age and smoking history, it's likely her shortness of breath is related to COPD, rather than having any link to alpha-1-antitrypsin deficiency.

D Bronchoscopy

Neutrophils and eosinophils may both be seen in bronchial fluid when there is underlying airway inflammation related to asthma, but this doesn't link strongly to the response to corticosteroids.

20605

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 64-year-old woman presents to the Emergency Department with severe shortness of breath, fever and cough productive of rusty-coloured sputum. She has no past medical history but reports an allergy to penicillin that she experienced in childhood. On examination, she is alert and orientated. On auscultation of her chest, crackles are heard at the right base. Her blood pressure is 110/70 mmHg. She is tachycardic, with a heart rate of 101 bpm. Her respiratory rate is 34 breaths per minute, and she is pyrexial at 38.2 °C.

Investigations:

Chest X-ray reveals right basal consolidation.

Investigation	**Result**	**Normal value**
Haemoglobin	121 g/l	115–155 g/l
White cell count (WCC)	13.5 × 10^9/l	4–11 × 10^9/l
Platelets	204 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	140 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Urea	8.0 mmol/l	2.5–6.5 mmol/l
Creatinine	110 micromol/l	50–120 µmol/l
C-reactive protein (CRP)	180 mg/l	< 5 mg/l
$P_a(O_2)$	7.6 kPa	10.5–13.5 kPa
$P_a(CO_2)$	6.1 kPa	4.6–6.0 kPa
Electrocardiogram	Sinus tachycardia – 101 bpm	

What is the most appropriate treatment regime for this patient?

Your answer was incorrect

A Intravenous (IV) levofloxacin and oral clarithromycin

B IV cefuroxime, oral clarithromycin and oxygen therapy via face mask

C Oral amoxicillin, oral clarithromycin and oxygen therapy via nasal cannulae

D Oral doxycycline and oxygen therapy via nasal cannulae

E Oral erythromycin

Explanation

D Oral doxycycline and oxygen therapy via nasal cannulae

This patient has community-acquired pneumonia (CAP). The British Thoracic Society (BTS) guidelines recommend the use of the CURB-65 score to stratify patients into different management groups.

The CURB-65 score is a simple additive score, ranging from 0 to 5, based on the presence of confusion, urea, respiratory rate, blood pressure and age, which can be used to stratify patients with CAP into different management groups.

Each of the following features scores 1 point:

- confusion (< 8/10 score on Abbreviated Mental Test (AMT))
- urea > 7 mmol/l
- respiratory rate > 30 breaths per minute
- blood pressure – systolic < 90 mmHg and/or diastolic < 60 mmHg
- age > 65.

Patients with scores of 0 and 1 are at low risk of mortality (< 2%) and might be suitable for management as hospital outpatients or by their General Practitioner. Patients with a score of 2 are at intermediate risk of mortality (9%) and should be considered for hospital-supervised treatment. Patients with a score of > 2 are at high risk of mortality (> 19%). They should be treated in the hospital. Initial care in an Intensive Care or High Dependency Unit may be appropriate for those with the highest scores.

This patient has a score of 2 and therefore requires treatment for moderate severity. Recommended treatment for patients with moderate-severity CAP is oral amoxicillin with a macrolide. If patients are intolerant to penicillins or macrolides, oral doxycycline is the main alternative agent.

In addition to antibiotic therapy, the patient should receive oxygen therapy to maintain a $p_a(O_2)$ of > 8 kPa and therefore, this patient should also receive oxygen treatment. However, given that $p_a(O_2)$ is only just below 8 kPa, delivery of low-flow oxygen therapy via nasal cannulae should achieve the required target.

C Oral amoxicillin, oral clarithromycin and oxygen therapy via nasal cannulae

Oral amoxicillin and clarithromycin with oxygen therapy would be the treatment of choice for this patient if she did not have a known penicillin allergy.

A Intravenous (IV) levofloxacin and oral clarithromycin

This would be the antibiotic treatment of choice in a patient with a penicillin allergy who is unable to tolerate oral medications. However, there is nothing in the scenario to suggest this patient would not tolerate oral treatment. Furthermore, she requires treatment with oxygen therapy so this would not be appropriate.

B IV cefuroxime, oral clarithromycin and oxygen therapy via face mask

In patients with moderate-severity community-acquired pneumonia (CAP) who cannot tolerate oral treatment and are allergic to penicillin, intravenous (IV) levofloxacin or a second-or third-generation cephalosporin (eg cefuroxime or cefotaxime) can be used in combination with clarithromycin. There is nothing in the scenario to suggest this patient cannot tolerate oral treatment and furthermore oxygen via a face mask is unlikely to be needed to improve her $p_a(O_2)$.

E Oral erythromycin

Oral erythromycin can be used as an alternative to clarithromycin in pregnant patients. However, it should be used in combination with a penicillin, tetracycline or fluoroquinolone antibiotic to cover for atypical pathogens.

21468

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation s...
ncbi.nlm.nih.gov/pubmed/12728155
(https://www.ncbi.nlm.nih.gov/pubmed/12728155)

BTS management of CAP
brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-mana...
(https://www.brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/)

A 21-year-old woman who is visiting the United Kingdom from East Africa comes to the Emergency Department for review. She has been suffering from chronic cough, fevers and weight loss over the last six months and is worried there may be a serious underlying problem. There is no past medical history of note, apart from an episode of shingles on the chest wall some 12 months earlier.

On examination, she is thin, with a body mass index (BMI) of 20 kg/m^2, and there is generalised lymphadenopathy. There are coarse breath sounds in the mid and lower zones bilaterally on auscultation.

Investigations reveal the following:

Investigations	**Results**	**Normal Values**
Haemoglobin (Hb)	109 g/l	115–155 g/l
White cell count (WCC)	4.8 × 10^9/l	4–11 × 10^9/l
Lymphocytes	1.0 × 10^9/l	1.5–3.5 × 10^9/l
Platelets (PLT)	171 × 10^9/l	150–400 × 10^9/l
Sodium (Na$^+$)	137 mmol/l	135–145 mmol/l
Potassium (K$^+$)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	95 μmol/l	50–120 μmol/l

Mantoux test: Negative

Her chest X-ray is shown below:



Which of the following is the most likely diagnosis?

Your answer was correct

A	Pertussis scarring
B	Sarcoidosis
C	Scleroderma
D	Tuberculosis
E	Varicella pneumonia

Explanation

D Tuberculosis

Human immunodeficiency virus (HIV) infection is a recognised cause of negative Mantoux, in the presence of active TB, the most likely clinical picture here. The chest X-ray picture fits with miliary TB, and the symptoms of fevers, cough and weight loss also support the diagnosis. Increased exposure to HIV is seen in patients from the

African subcontinent. Initiation of anti-tuberculous therapy, coupled with highly active antiretroviral therapy (HAART) once the diagnosis of HIV is confirmed, are the most appropriate next steps.

Image source: https://commons.wikimedia.org/wiki/File:Chest_radiograph_of_miliary_tuberculosis_1.jpg (https://commons.wikimedia.org/wiki/File:Chest_radiograph_of_miliary_tuberculosis_1.jpg)

A Pertussis scarring

Scarring over the long term from pertussis leads to symptoms and signs of bronchiectasis. With respect to the chest X-ray findings, these can include parallel line opacities, known as tram tracks, caused by dilated, thickened bronchi. Ring opacities or cystic spaces may be seen which are large (2 cm in diameter), and these result from cystic bronchiectasis. The signet ring sign, which is said to be characteristic of bronchiectasis, corresponds to a dilated bronchus, immediately adjacent to a smaller pulmonary artery.

B Sarcoidosis

Sarcoidosis is characteristically associated with bilateral hilar lymphadenopathy. It does not fit with the multiple peripheral nodules seen here on the chest X-ray.

C Scleroderma

Scleroderma (systemic sclerosis) does not fit with the picture seen here on chest X-ray. The chest X-ray may be normal in early pulmonary fibrosis related to systemic sclerosis. Over the longer term, generalised patchy opacification consistent with pulmonary fibrosis is seen, rather than the nodular changes seen here.

E Varicella pneumonia

Previous varicella pneumonia does lead to nodular calcification seen on chest X-ray, but it would not explain the symptoms of active infection reported here.

39657

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 25-year-old man is brought to the Emergency Department following a road traffic accident in which he was thrown off his motorbike. He is the subject of a trauma call. He is complaining of pain in his chest and upper abdomen which radiates to his back and left shoulder. He was previously fit and well.

On examination, he looks unwell. His BP is 90/55 mmHg, pulse 130 bpm. He is bruised and tender over his anterior chest wall and on palpation crackling can be felt. Auscultation of his chest reveals quiet breath sounds at the left base.

His chest X-ray is shown below:

A chest drain is inserted in the Emergency Department.

Investigation	Result	Normal Values
Pleural fluid protein	45 g/l	25 - 35 g/l
Pleural amylase	240 u/l	30 - 110 u/l
Serum amylase	190 u/l	23 - 85 u/l



What is the most likely diagnosis?

Your answer was incorrect

A Oesophageal rupture

B Pancreatitis

C Rib fractures and haemothorax

D Chylothorax

E Aortic dissection

Explanation

A Oesophageal rupture

Oesophageal rupture is a serious condition following trauma, violent vomiting or endoscopic procedures, or can be associated with malignancy. Spillage of gastric contents into the pleurae causes shock and pain. The chest X-ray usually shows a left pleural effusion or hydro-pneumothorax. There may also be air in the soft tissues (as in this patient), termed surgical emphysema. Pleural fluid is an exudate, and amylase levels are high. The ratio of pleural fluid amylase to serum amylase is >1.0. Iso-enzyme analysis can be useful, as oesophageal rupture will show the amylase to be salivary in origin.
Diagnosis is made radiologically using water-soluble contrast. Surgical treatment is required and mortality is high.

C Rib fractures and haemothorax

There are no rib fractures visible on the CXR and no mention of blood in the pleural tap.

B Pancreatitis

Serum amylase is only mildly raised, making an acute pancreatitis unlikely. Serum amylase can be normal in chronic pancreatitis, but from the history the patient was previously fit and well. The history of trauma in this case makes oesophageal rupture a more likely diagnosis.

D Chylothorax

A chylothorax may be caused by a tear in the thoracic duct, which could occur with trauma. Pleural fluid has a milky appearance and pleural triglyceride levels are high, neither of which is mentioned here. The amylase values in serum and pleural fluid make oesophageal rupture more likely.

E Aortic dissection

The mediastinum is not broadened, and pleural effusion with high amylase levels is more in keeping with oesophageal rupture.

70143

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pleural Effusion

Media

Expanded explanations

Flashcards

A patient comes to the Emergency Department with difficulty in breathing which started the previous night. He has difficult giving history, as his lower jaw is painful when he tries to talk. On examination, he is dyspnoeic, febrile and anxious. His lower jaw is inflamed, swollen, oedematous and tender.

What is the immediate appropriate step in the management of this patient?

Your answer was incorrect

A Examine the inside of his mouth with a spatula

B Request for an urgent chest X-ray

C Take blood cultures

D Take bloods for U/E, full blood count (FBC) and C-reactive protein (CRP)

E Call for an anaesthetist

Explanation

E Call for an anaesthetist

This patient has most likely got Ludwig’s angina, which is a rapidly spreading, life-threatening cellulitis of the sublingual and submandibular spaces that usually starts in an infected lower molar. Airway obstruction may result as the infection spreads to the supraglottic tissues.

C Take blood cultures

Following definitive management of the patient’s airway, treatment with intravenous antibiotics active against streptococci and oral anaerobes should be started.

A Examine the inside of his mouth with a spatula

This patient’s breathless may be secondary to partial airway obstruction. An anaesthetist should be called to visualise the airway and institute definitive management as required.

B Request for an urgent chest X-ray

This patient is at significant risk of airway obstruction; therefore management of their airway is the most appropriate next immediate step.

D Take bloods for U/E, full blood count (FBC) and C-reactive protein (CRP)

Routine bloods will be helpful in tracking the patient’s progress over coming days, but taking them is not an immediate priority.

7505

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

An 18-year-old man presents to the Outpatients Department having coughed up a small amount of bright red blood. He is mildly short of breath and feels ‘wheezy’. His mother says he has always been a ‘chesty’ child. He was teased at school as he has always been the smallest in his class. His brother and sister were fit and well. He had a dog at home.

On examination he looked well and was not breathless at rest. He had clubbing. He had nasal polyps and auscultation of his chest revealed coarse crackles throughout both lung fields. His abdomen was distended but soft.

Urinalysis:

Protein	+
Glucose	++
Nitrites	-
Leucocytes	-

Haematology:

Investigation	Result	Normal Values
Haemoglobin (Hb)	110 g/l	135 - 175 g/l
White cell count (WCC)	12.1 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	450 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	137 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	62 µmol/l	50 - 120 µmol/l
Neutrophils	91%	
Lymphocytes	8%	
Monocytes	0.4%	
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	35 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	40 g/l	< 10 g/l
Urea	6.4 mmol/l	2.5 - 6.5 mmol/l
Bilirubin	31 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	46 u/l	10 - 40 u/l

Alkaline phosphatase (ALP)	221 u/l	30 - 150 u/l
Albumin	38 g/l	35 - 55 g/l
Protein	74 g/l	
Magnesium (Mg^{2+})	0.64 mmol/l	0.6 - 1.1 mmol/l
Corrected calcium (Ca^{2+})	2.1 mmol/l	2.2 - 2.7 mmol/l
Phosphate (PO_4^{3-})	0.75 mmol/l	1.1 - 1.45 mmol/l

Spirometry:

FEV1	60% predicted
FVC	80% predicted
PEF	350 l/min
KCO	70% predicted

His chest X-ray is shown below.



What is the test most likely to establish a diagnosis?

Your answer was incorrect

A Metacholine challenge test

B Anti-GBM antibodies

C Electron microscopy of cilia

D Sweat test

E Aspergillus precipitins

Explanation

D Sweat test

This boy has cystic fibrosis. He has had recurrent chest infections secondary to bronchiectasis (confirmed by examination findings and the chest X-ray, also note long line inserted via left antecubital fossa), he is small for his age, has nasal polyps (which occur in one third of patients with CF) and clubbing. Pulmonary function tests show airflow obstruction He has mildly deranged liver function tests – secondary to obstruction of biliary ductules in the liver. This can eventually lead to cirrhosis. He has glucose in his urine – pancreatic islet cells are destroyed as the pancreas becomes fibrotic resulting in CF related diabetes in 30% of patients by adulthood.

The diagnosis of cystic fibrosis is made by a sweat test and genetic analysis. A sweat chloride concentration over 60 mmol/l is indicative of cystic fibrosis.

Treatment is with pancreatic supplements, prophylactic inhaled or nebulised antibiotics, strict glucose control, mucolytics/DNAase and regular physiotherapy with postural drainage. Prophylactic Azithromycin is used for its immunomodulatory effects. Non invasive ventilation may be used in patients with chronic type 2 respiratory failure as a bridge to lung transplantation Lung transplantation is considered as a treatment option once there is severe respiratory compromise.

B Anti-GBM antibodies

This is a test for Goodpasture syndrome, which is associated with a deterioration in renal function accompanied by haemoptysis over a relatively short period of time.

A Metacholine challenge test

This is a test for asthma. It is rarely performed, given that a diagnosis of asthma can usually be made according to other clinical criteria.

C Electron microscopy of cilia

This test is used to diagnose primary ciliary dyskinesia or Kartagener syndrome.

E Aspergillus precipitins

This is a test used in the diagnosis of ABPA, which is associated with symptoms of asthma and patchy shadowing on the CXR. A poor response is usually seen to conventional inhaled asthma medications.

70086

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 54-year-old man comes to the Emergency Department complaining of shortness of breath. He has been unwell for a few days and most recently has begun to spike fevers. He has a history of a previous myocardial infarction (MI) some 4 years earlier and mild left ventricular failure as a consequence. Medication includes ramipril 10 mg daily, aspirin 75 mg and bendroflumethiazide 2.5 mg. On examination his BP is 115/70 mmHg, pulse 95/min and his temperature is 37.9°C. He has bilateral crackles on auscultation, perhaps more marked on the left than the right, and a respiratory rate of 36/min.

Investigations:

Investigation	Result	Normal value
pH	7.6	7.35–7.45
pCO_2	3.5 kPa	4.6–6.0 kPa
pO_2	7.2 kPa	10.5–13.5 kPa
Bicarbonate	33 mmol/l	24–30 mmol/l

Which of the following best describes his acid–base disturbance?

Your answer was correct

A Compensated respiratory alkalosis

B Compensated respiratory acidosis

C Mixed respiratory and metabolic alkalosis

D Compensated metabolic alkalosis

E Metabolic and respiratory acidosis

Explanation

C Mixed respiratory and metabolic alkalosis

This situation with an elevated bicarbonate and low CO_2 and O_2 can be seen in patients who are chronic diuretic users and who suffer an acute respiratory event such as a pulmonary embolus, where they hyperventilate to maintain adequate oxygenation.

A Compensated respiratory alkalosis

Compensation implies that bicarbonate would be reduced to maintain pH in the normal range; however, a bicarbonate of 33 mmol/l means that is clearly not the case here.

B Compensated respiratory acidosis

In a compensated respiratory acidosis, bicarbonate and CO_2 are both elevated.

D Compensated metabolic alkalosis

Respiratory compensation for metabolic alkalosis is limited because of hypoxic drive; it may, however, be associated with a small rise in Pa CO2.

E Metabolic and respiratory acidosis

A mixed metabolic and respiratory acidosis presents with low pH and low bicarbonate, with elevated CO2, in contrast to the picture seen here.

20982

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Respiratory Physiology

Expanded
explanations

A 52-year-old farmer presented with a 5-month history of progressively worsening exertional dyspnoea and a persistent dry cough. He also described 6 kg weight loss and general malaise over this time period. He had no significant past medical history or family history. He was an ex-smoker who stopped 10 years ago and drank 10–12 units of alcohol per week.

On examination, he was dyspnoeic on minimal exertion, but not at rest. He was not cyanosed and there was no finger clubbing. He had low-grade pyrexia of 37.4 °C. His heart sounds were normal, jugular venous pressure was not raised and there was no peripheral oedema. He had bilateral fine apical crepitations on chest auscultation. Abdominal and neurological examinations were unremarkable.

Investigations reveal the following:

Investigation	**Result**	**Normal value**
Haemoglobin (Hb)	112 g/l	135–175 g/l
White cell count (WCC)	12.4×10^9/l	$4.0–11.0 \times 10^9$/l
Platelets (PLT)	420×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	140 mmol/l	135–145 mmol/l
Potassium (K^+)	3.6 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	110 μmol/l	50–120 μmol/l
Corrected calcium (Ca^{2+})	2.42 mmol/l	2.20–2.60 mmol/l
Bilirubin	14 μmol/l	2–17 μmol/l
Urea	6.1 mmol/l	2.5–6.5 mmol/l
Alanine aminotransferase (ALT)	32 IU/l	5–30 IU/l
Alkaline phosphatase (ALP)	92 IU/l	30–130 IU/l
Gamma glutamyl-transferase (GGT)	44 IU/l	5–30 IU/l
Mean corpuscular volume (MCV)	92 fl	76–98 fl
Erythrocyte sedimentation rate (ESR)	60 mm/hour	1–20 mm/hour
Pulmonary function tests (PFTs)	Restrictive defect, reduced gas transfer	

A high-resolution computed tomography (CT) scan of the thorax is shown below:



What is the most likely diagnosis?

Your answer was incorrect

A	Sarcoidosis
B	Allergic bronchopulmonary aspergillosis
C	Idiopathic pulmonary fibrosis
D	Extrinsic allergic alveolitis

E Pulmonary tuberculosis

Explanation

D Extrinsic allergic alveolitis

This man has chronic extrinsic allergic alveolitis (EAA). This is caused by inhaled allergens provoking a hypersensitivity reaction in the lungs and resulting in diffuse inflammation involving the small airways and alveoli. In the chronic form, lymphocytic and giant cell infiltration can eventually lead to pulmonary fibrosis. Typical allergens include microbial spores, especially *Micropolyspora faeni* in mouldy hay, which causes farmer's lung, and avian proteins causing bird fancier's lung. The diagnosis can be made by a combination of imaging, respiratory function tests, serum precipitins and bronchoalveolar lavage. The computed tomography (CT) scan shows multiple small, ill-defined nodules throughout both lungs, with areas of ground-glass opacity in the basal segments. Treatment consists of avoidance of the causative allergen and long-term corticosteroid therapy. In some patients progression to end-stage fibrosis and respiratory failure cannot be prevented.

C Idiopathic pulmonary fibrosis

Idiopathic pulmonary fibrosis in this case is a diagnosis of exclusion. The occupation as a farmer means that disease related to exposure to inhaled allergens should be excluded first.

A Sarcoidosis

Sarcoidosis is associated with progressive pulmonary fibrosis if there is marked disease activity which is left untreated, but other features associated with active sarcoid, such as erythema nodosum and elevated serum calcium, appear absent here.

B Allergic bronchopulmonary aspergillosis

ABPA is associated with exaggerated asthma symptoms that do not respond to inhaled therapies, rather than the restrictive lung picture seen here.

E Pulmonary tuberculosis

There is no suggestion of TB exposure, and the CT image is suggestive of fibrosis rather than showing any sign of focal infection.

9150

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 68-year-old man presents to the hospital with a 12-hour history of acute, severe epigastric pain radiating to the back, as well as profuse vomiting. He has a long history of alcohol excess and admitted to a drinking binge 24 hours before the onset of pain. His past medical history includes coronary artery bypass graft (CABG) surgery two years ago and hypertension.

On examination, he has pyrexia of 38 °C. His heart rate is 120 bpm and regular and his blood pressure is 92/60 mmHg. Oxygen saturation is 91% on air. Heart sounds are normal, with few basal crepitations on chest auscultation. He is very tender in the upper abdomen, with rebound and guarding.

Investigations:

Investigation	Result	Normal value
Haemoglobin (Hb)	128 g/l	135–175 g/l
White cell count (WCC)	21.0 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	140 × 10^9/l	150–400 × 10^9/l
Mean corpuscular volume (MCV)	106 fl	76–98 fl
Corrected calcium (Ca^{2+})	1.96 mmol/l	2.20–2.60 mmol/l
Bilirubin	19 µmol/l	2–17 µmol/l
Alanine aminotransferase (ALT)	50 IU/l	5–30 IU/l
Urea	6.2 mmol/l	2.5–6.5 mmol/l
Alkaline phosphatase (ALP)	120 IU/l	30–130 IU/l
Gamma glutamyl-transferase (γGT)	110 IU/l	5–30 IU/l
Glucose	11.0 mmol/l	3.5–5.5 mmol/l
Amylase	1320 U/l	< 200 U/l
C-reactive protein (CRP)	350 mg/l	< 0 mg/l
Electrocardiogram (ECG)	Sinus tachycardia, no acute changes	
Chest X-ray (CXR)	Patchy shadowing lower zones, no free air under diaphragm	

Initial blood gases on air reveal the following:

Investigation	Result
pH	7.160

$p_a(O_2)$	7.8 kPa
$p_a(CO_2)$	3.2 kPa
HCO^{3-}	21
Lactate	7

He is treated with oxygen, intravenous fluids, antibiotics and opiate analgesia. However, 24 hours later he has not improved and is getting progressively more breathless and hypoxic.

His repeat blood gas with a FiO_2 of 35% reveals the following:

Investigation	**Result**
pH	7.25
$p_a(O_2)$	7.8 kPa
$p_a(CO_2)$	7.2 kPa
HCO^{3-}	26
Lactate	4

What is the most likely cause of this deterioration?

Your answer was incorrect

A Acute respiratory distress syndrome (ARDS)

B Aspiration pneumonia

C Cardiogenic pulmonary oedema

D Pulmonary haemorrhage

E Pulmonary embolism

Explanation

A Acute respiratory distress syndrome (ARDS)

This patient has developed acute respiratory distress syndrome (ARDS) as a complication of severe acute pancreatitis. It is defined as acute respiratory failure with non-cardiogenic pulmonary oedema secondary to severe pulmonary or systemic illness and may be associated with multiorgan failure. The most common cause is sepsis. There is increased permeability of the pulmonary microcirculation due to inflammation causing leakage of fluid into the lungs. Later features include pulmonary vasoconstriction and hypertension secondary to hypoxia, and lung exudates which lead to reduced compliance. Initial blood gases can reveal respiratory alkalosis, and the development of acute hypercapnic respiratory acidosis can represent severe ARDS with impending respiratory arrest. ARDS should be managed on an Intensive Care Unit. Supportive measures include positive-pressure ventilation, circulatory support with inotropes and treatment of sepsis. Inhaled nitrogen oxide and haemofiltration can also be used. Mortality is currently around 20–30%.

B Aspiration pneumonia

Aspiration pneumonia may be caused by vomiting and result in hypoxaemia and respiratory acidosis. However, the presence of pancreatitis as a precipitant and the finding of bilateral chest X-ray changes are more in keeping with ARDS as the cause.

C Cardiogenic pulmonary oedema

Cardiogenic pulmonary oedema is a possible differential due to administration of fluids and a background of coronary artery bypass graft indicating ischaemic heart disease and likely acute illness due to reduced ventricular compliance. However, the presence of acute pancreatitis is more likely to be associated with ARDS, and the lower lobe changes are more consistent with ARDS than with pulmonary oedema.

D Pulmonary haemorrhage

Pulmonary haemorrhage is not associated with pancreatitis, although it is associated with patchy bilateral lung infiltrates on the chest X-ray.

E Pulmonary embolism

Pulmonary embolism is associated with shortness of breath and chest pain. Although it can be precipitated by hypovolaemia, it is not in keeping with the patchy lower zone changes seen on the chest X-ray.

9152

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

You are called to a cardiac arrest on the coronary care unit. A 74-year-old woman with a 40-pack year smoking history and COPD on high-dose salmeterol/fluticasone combination inhaler has arrested two days after admission with an inferior myocardial infarction.

Arterial blood gasses on admission reveal the following:

Investigation	**Result**	**Normal value**
pH	7.39	7.35–7.45
pO_2	8.5 mmol/l	10.5–13.5 kPa
pCO_2	5.8 mmol/l	4.6–6.0 kPa
HCO_3^-	31 mmol/l	24–30 mmol/l

Arterial blood gasses on 35% O_2 post-cardiac arrest reveal the following:

Investigation	**Result**	**Normal values**
pH	7.29	7.35–7.45
pO_2	7.5 kPa	10.5–13.5 kPa
pCO_2	6.9 kPa	4.6–6.0 kPa

She responds appropriately to commands and is cooperative and conscious around 15 min after her cardiac arrest. A portable chest X-ray is shown below.



Which of the following is the most appropriate course of action with respect to her ventilatory support?

Your answer was correct

A	Continue 35% O_2 therapy
B	Reduce her inspired oxygen to 28%
C	Increase her inspired oxygen to 60%
D	Consider her for non-invasive positive pressure ventilation (NIPPV)
E	Move to intubation and ventilation

Explanation

D Consider her for non-invasive positive pressure ventilation (NIPPV)

On admission her arterial blood gasses revealed a compensated respiratory acidosis with some degree of CO_2 retention. Post cardiac arrest she has decompensated with relative hypoxia and CO_2 retention. At this point the concern is that the acidosis may promote further respiratory weakness, driving CO_2 retention and worsening the pH further. NIPPV should reverse the respiratory acidosis and may have a positive impact on any pulmonary oedema associated with the MI. The chest x-ray shows hyperinflation, in keeping with COPD, and an absence of any sign of resuscitation injury.

Image source: https://commons.wikimedia.org/wiki/File:COPD.JPG (https://commons.wikimedia.org/wiki/File:COPD.JPG)

A Continue 35% O_2 therapy

Continuing her 35% O_2 may further exacerbate CO_2 retention and lead to respiratory arrest, because of increasing drowsiness and lethargy.

B Reduce her inspired oxygen to 28%

This risks worsening any hypoxia, which could exacerbate myocardial ischaemia leading to chest pain and/or rhythm disturbance.

C Increase her inspired oxygen to 60%

This, like continuing 35% O_2, risks worsening respiratory acidosis without significantly increasing PaO_2 because of consequent respiratory muscle weakness and steadily decreasing consciousness.

E Move to intubation and ventilation

This should be carefully considered based on pre-morbid status, and ideally would be a step only after NIPPV had not succeeded in maintaining oxygenation and reversing respiratory acidosis.

18537

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Ventilatory Support

Expanded explanations

A 32-year-old African-Caribbean nurse is referred to the Chest Clinic with several months of progressive dyspnoea and a dry cough. She is now short of breath on minimal exertion. She was previously fit and well, has no relevant family history and is on no medications. She is a non-smoker.

On closer questioning, she also described fatigue and weight loss and a painful eye with blurred vision for the last few weeks.

On examination, she was comfortable at rest with mild pyrexia of 37.4 °C. Her heart sounds were normal and she had fine crackles at both mid-zones in her chest. Her abdomen was non-tender with no palpable masses. She had no focal neurological signs. She had an inflamed, watery right eye with reduced visual acuity.

On inspection of her skin, she had several raised erythematous lesions on her shins that were tender to palpation.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	98 g/l	115–155 g/l
White cell count (WCC)	7.6×10^9/l	$4–11 \times 10^9$/l
Platelets	360×10^9/l	$150–400 \times 10^9$/l
γGT	32 IU/l	5–30 IU/l
Mean corpuscular volume (MCV)	92 fl	76–98 fl
Erythrocyte sedimentation rate (ESR)	80 mm/hr	0–10 mm in the 1st hour
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K+)	4.2 mmol/l	3.5–5.0 mmol/l
Corrected Ca^{2+}	2.78 mmol/l	2.20–2.60 mmol/l
Albumin	36 g/l	35–55 g/l
Bilirubin	14 µmol/l	2–17 µmol/l
Urea	4.2 mmol/l	2.5–6.5 mmol/l
Creatinine	82 µmol/l	50–120 µmol/l
Alanine aminotransferase (ALT)	38 IU/l	5–30 IU/l
Alkaline phosphatase	60 IU/l	30–130 IU/l

Pulmonary function tests:

Forced expiratory volume (FEV_1)	78% of predicted
Forced vital capacity (FVC)	70% of predicted
Transfer co-efficient (Kco)	78% of predicted

His chest x-ray is shown below.



Which of the following is the most appropriate initial therapy for this patient?

Your answer was correct

A Anti-tuberculous chemotherapy

B Ciclosporin

C Cyclophosphamide

D Methotrexate

E Prednisolone

Explanation

E Prednisolone

The chest x-ray shows bilateral hilar lymphadenopathy (BHL) with clear lung fields. Taken together with the anterior uveitis and erythema nodosum the underlying diagnosis is sarcoidosis. Steroid are indicated as a first line treatment, as the patient has hypercalcaemia. Without hypercalcaemia, the management would simply be supportive. Local corticosteroids may be appropriate for her eye disease.

Image source:
https://commons.wikimedia.org/wiki/File:Pulmonary_sarcoidosis_in_a_50_year_old_male_patient,_X-ray_thorax_image,_2006.jpg
(https://commons.wikimedia.org/wiki/File:Pulmonary_sarcoidosis_in_a_50_year_old_male_patient,_X-ray_thorax_image,_2006.jpg)

A Anti-tuberculous chemotherapy

This woman has pulmonary sarcoidosis with anterior uveitis and erythema nodosum, with evidence of bilateral hilar lymphadenopathy on her chest x-ray. Therefore, this option is incorrect as this is the treatment for TB.

B Ciclosporin

Ciclosporin could be considered further down the line as a steroid-sparing agent if the patient was unable to wean off prednisolone.

C Cyclophosphamide

Cyclophosphamide could be considered further down the line if the patient is particularly resistant to high-dose steroids.

D Methotrexate

Methotrexate could be considered at a later stage as a steroid-sparing agent if the patient fails to wean off prednisolone.

32406

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

You review a 35-year-old woman who is overweight, with a body mass index (BMI) of 31. She presents with increasing lethargy and shortness of breath over the past few weeks. She has a past history of deep vein thrombosis three years ago. Her BP is 145/82 mmHg, with pulse 92/min and regular. Saturations are 91% on air and fall on only minor exercise. D-dimers are above the upper limit of normal.

Electrocardiogram (ECG) is shown below:



Courtesy of J.S.S.G. de Jong, MD, AMC, The Netherlands

ECGPEDIA.ORG

Which of the following diagnoses fits best with this clinical picture?

Your answer was incorrect

A Right heart failure

B Sleep apnoea syndrome

C Single massive pulmonary embolus

D Multiple pulmonary emboli

E Cardiomyopathy

Explanation

D Multiple pulmonary emboli

The ECG shows right ventricular hypertrophy and right atrial enlargement. When combined with the previous history of deep venous thrombosis, and raised D-dimer, the the diagnosis of recurrent thromboembolic disease is most likely. Lifelong warfarinisation is the treatment of choice. If further pulmonary emboli occur, then the patient should be considered for insertion of an inferior vena cava (IVC) filter.

Image source: https://en.ecgpedia.org/index.php?title=File:E000004.jpg#filelinks

C Single massive pulmonary embolus

There is no evidence of circulatory collapse – this patient does not have a low systolic blood pressure drop to suggest a massive pulmonary embolus.

A Right heart failure

There is no evidence of right heart failure symptoms, such as elevated JVP, peripheral oedema or ascites.

B Sleep apnoea syndrome

Although this patient is overweight (a risk factor for sleep apnoea), she does not have the features suggestive of sleep apnoea such as sleepiness and snoring.

E Cardiomyopathy

Unless this patient has arrhythmogenic right cardiomyopathy (a rare genetic condition), right ventricular hypertrophy on ECG does not fit with this picture.

32449

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total: 301

Responses - % Correct: 27%

Pulmonary Embolism

Media

Expanded explanations

Flashcards

A 64-year-old man is referred to the Respiratory Clinic with slowly progressive shortness of breath and a chronic cough. He has been suffering from fevers over the past few months and has lost 3 kg in weight. He worked making stone sculptures for many years and has smoked 10 cigarettes per day for the past 25 years. Examination reveals a BP of 132/72 mmHg; his heart rate is 85 bpm and regular. There are scattered crackles on auscultation, more marked in the upper zones. Pulmonary function tests (PFTs) show mixed obstructive/restrictive defect.

A CXR is taken:



Which of the following is the most likely diagnosis?

Your answer was correct

A Aspergillosis

B Chronic obstructive pulmonary disease

C Sarcoidosis

D Silicosis

E Tuberculosis

Explanation

D Silicosis

This patient has a very long history of exposure to silica dust, and the CXR changes seen are suggestive of patchy fibrosis. Removal of exposure to silica dust and stopping smoking are key steps in the management of this patient. No therapies have been proven to alter the course of the disease, although corticosteroids may be of benefit in treating acute exacerbations of silicosis. Hypercalcaemia can also occur in granulomatous silicosis.

A Aspergillosis

Allergic bronchopulmonary aspergillosis (ABPA) may produce similar changes on a CXR, although given this patient's occupation, silicosis is considered more likely. ABPA is treated with a combination of oral corticosteroids and itraconazole.

B Chronic obstructive pulmonary disease

The CXR changes are consistent with patchy fibrosis, not with chronic obstructive pulmonary disease (COPD). In COPD, increased bronchovascular markings and sometimes bullae (in emphysema) are seen. Additionally, the PFTs expected in COPD would show an obstructive pattern.

C Sarcoidosis

In sarcoidosis, there is hilar lymphadenopathy and patchy areas of fibrosis when the disease is severe. The relative absence of hilar lymphadenopathy here makes sarcoidosis less likely. Only around 15% of patients with sarcoidosis have respiratory symptoms, and the use of steroids to treat the disease is reserved for symptomatic patients.

E Tuberculosis

Changes due to tuberculosis (TB), such as upper zone consolidation or opacities, are more likely to be confined to one side of the chest rather than the diffuse pattern seen here, and we are given no history of potential exposure to TB.

38297

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Silicosis

Media

Expanded explanations

Flashcards

A 17-year-old woman with a history of cystic fibrosis presents to the Emergency Room with rapidly worsening shortness of breath, and a cough productive of much more sputum than usual with marked wheeze. She has started to lose weight and is off her food. On examination her BMI is 21, her BP is 105/72 mmHg and pulse 89/min with a temperature of 37.9°C. Auscultation of the chest reveals a mix of crackles and wheeze.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	110 g/l	115–155 g/l
White cell count (WCC)	12.2 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	199 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	140 mmol/l	135–145 mmol/l
Potassium (K^+)	5.0 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	140 µmol/l	50–120 µmol/l
Sputum sample	Gram-negative rods	

Which of the following is the most likely identity of the bacteria found in the sputum sample?

Your answer was incorrect

A *Klebsiella*

B *Mycoplasma*

C *Pseudomonas*

D *Bacteroides*

E *Staphylococcus*

Explanation

C *Pseudomonas*

Pseudomonas colonisation is associated with deterioration of respiratory symptoms, loss of appetite and weight loss in patients with CF. It is seen as Gram negative rods on a sputum sample. Severity of pulmonary disease, including the degree of *Pseudomonas colonisation*, is closely related to outcome and prognosis. Inhaled tobramycin given intermittently over a prolonged period has been shown to delay the deterioration in lung function and improve prognosis for patients chronically infected with *Pseudomonas.*

B *Mycoplasma*

Mycoplasma is a mollicute which lacks a peptidoglycan cell wall; it is most recognised as a cause of pneumonia in epidemics, often in individuals who reside in close proximity – for example, soldiers in the army.

A *Klebsiella*

Klebsiella are Gram-negative rod-shaped bacteria, but rather than causing chronic infection they are more likely to be associated with consolidation and abscess formation in patients who are immunosuppressed or who have chronic alcoholism.

D *Bacteroides*

Bacteroides is generally recognised as part of the normal colonic flora, and colonisation therefore doesn't play a key role in deteriorating lung function associated with cystic fibrosis.

E *Staphylococcus*

Staphylococci are Gram-positive cocci that lead to acute pneumonia, usually after influenza infection.

20952

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 63-year-old man was brought to the Emergency Department by a ‘blue light’ ambulance after having been found at home by his daughter. It transpires that the daughter had gone out for 15 min and, upon her return, had discovered her father collapsed but conscious. She does not live with her father so was not aware of any recent health problems that her father had, but she knew that he was on medication for blood pressure and his heart. She recalls that her father was complaining of some retrosternal discomfort earlier after coughing and bringing up a little blood, and that he was slightly more short of breath than usual. She reports that he is known to have had a heart attack 6 months previously.

The paramedics have initiated volume replacement and protected the airway. In the casualty department, clinical examination shows the patient to be dyspnoeic and tachypnoeic, with a raised JVP. Appropriate bloods are obtained and fluid replacement continued. An ECG is obtained and shown below.



Unfortunately, shortly after his arrival in the Emergency Department, his condition deteriorates and he has a cardiac arrest. He is resuscitated and transferred to the ITU where he is ventilated but his condition continues to deteriorate.

What is the most likely diagnosis?

Your answer was correct

A Haemothorax

B Acute pulmonary embolus

C Left ventricular failure

D Acute respiratory distress syndrome (ARDS)

E Fat embolus

Explanation

B Acute pulmonary embolus

Pulmonary embolism (PE) is an often fatal complication of underlying venous thrombosis in the legs or pelvis. Microthrombi break off and pass through the heart before lodging in the pulmonary circulation. The classical triad of symptoms of PE – haemoptysis, shortness of breath and pleuritic chest pain – occur in less than a fifth of patients eventually diagnosed with PE. Massive PE causes hypotension due to acute right heart failure. The ECG here shows the uncommonly seen but classical S_1Q_3T3.

Image source: https://commons.wikimedia.org/wiki/File:Pulm_embolism.jpg (https://commons.wikimedia.org/wiki/File:Pulm_embolism.jpg)

A Haemothorax

A haemothorax typically occurs in the context of trauma, which is not seen in the history here, and is not as likely a diagnosis as PE in this case.

C Left ventricular failure

Left ventricular failure can occur after an acute myocardial infarction, but this patient's symptoms (specifically haemoptysis) and the ECG changes are not suggestive of this diagnosis.

D Acute respiratory distress syndrome (ARDS)

ARDS may be caused by lung injury or may occur secondary to a systemic illness such as sepsis or pancreatitis. ARDS is characterised by respiratory failure not of cardiogenic or hypovolaemic origin, occurring within 1 week of a precipitating trauma/condition, with the presence of bilateral opacities on CXR. Clinical features include dyspnoea, tachypnoea, tachycardia, cyanosis and low pO_2.

E Fat embolus

Fat embolism occurs following trauma to the long bones or pelvis, following parenteral lipid infusion, or recent corticosteroid administration.

6809

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 19-year-old man presents with a 10-day history of headaches and malaise, with the more recent development of a dry cough and breathlessness.

On examination, he was pyrexial, and he had a rash which looked like this:



Investigations:

Investigation	Result	Normal values
Haemoglobin (Hb)	94 g/l	135 - 175 g/l
White cell count (WCC)	10 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	9.2 × 10^9/l	1.5 - 7.0 × 10^9/l
Platelets (PLT)	200 × 10^9/l	150 - 400 × 10^9/l
Reticulocytes	5.1%	0.5 - 2.5 %
Sodium (Na^+)	129 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.0 mmol/l	3.5 - 5.0 mmol/l
Urea	4.4 mmol/l	2.5 - 6.5 mmol/l
Bilirubin	30 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	44 u/l	10 - 40 u/l
Alanine aminotransferase (ALT)	30 u/l	7 - 55 u/l
Alkaline phosphatase (ALP)	121 u/l	30 - 150 u/l
Albumin	35 g/l	35 - 55 g/l
Lactate dehydrogenase (LDH	495 u/l	140 - 280 u/l
Creatinine (Cr)	90 µmol/l	50 - 120 µmol/l

His chest X-ray is shown below:



Which is the most appropriate investigation in order to reach a diagnosis?

Your answer was incorrect

A	EBV serology
B	Urine *Legionella* antigen
C	Sputum culture
D	Mycobacterial culture
E	*Mycoplasma* serology

Explanation

E *Mycoplasma* serology

Mycoplasma usually affects young individuals and occurs in epidemics every 3–4 years. Chest symptoms are usually preceded by non-specific symptoms such as malaise and headaches.

The CXR usually shows only one lobe to be involved; however, about 20% show bilateral pneumonia. There is often a discrepancy between X-ray appearances and the clinical condition of the patient.

Diagnosis is by *Mycoplasma* serology – cold agglutinins occur in 50%. The image shows erythema multiforme, which further supports the diagnosis. The blood results suggest a haemolytic anaemia with a reticulocytosis, hyperbilirubinaemia and elevated LDH; autoimmune haemolytic anaemia (AHIA), when caused by cold agglutinins, is associated with *Mycoplasma* and can be diagnosed with a direct Coomb's test.

Extra-pulmonary complications of *Mycoplasma* include:

- Cardiovascular – myocarditis and pericarditis

- Dermatological – erythema multiforme (as shown here), erythema nodosum, Stevens–Johnson syndrome and non-specific rashes
- Gastrointestinal – hepatitis, pancreatitis, nausea, vomiting, anorexia and transient abdominal pain
- Neurological – meningoencephalitis, meningitis, ascending paralysis, transient myelitis, cranial nerve palsies, peripheral neuropathy, bullous myringitis
- Haematological – cold autoimmune haemolytic anaemia, thrombocytopenia, disseminated intravascular coagulation
- Renal – glomerulonephritis
- Others – arthralgia, arthritis, bullous myringitis.

Treatment: macrolides.

Causes of erythema multiforme:

- Infections – herpes simplex virus (most common cause), orf, HBV, HIV, EBV, mumps (paramyxovirus), *Mycoplasma*, psittacosis, *Rickettsiae*, *Streptococcus*, typhoid, diphtheria
- Drug reactions – barbiturates, penicillin, sulphonamides, phenytoin
- Connective tissue disease – SLE
- Vasculitis – polyarteritis nodosa, granulomatosis with polyangiitits

Others – underlying malignancy, sarcoidosis, rheumatoid arthritis, ulcerative colitis.

C Sputum culture

Mycoplasma is a fastidious organism, taking 3 weeks or more to grow in culture. Specialised culture medium is required, and the test lacks sensitivity compared to other investigations. For these reasons, sputum culture is not the first-line investigation for *Mycoplasma*.

A EBV serology

EBV serology would be appropriate if the diagnosis was suspected to be EBV infection causing infectious mononucleosis.

B Urine *Legionella* antigen

Urine *Legionella* antigen would be appropriate if the diagnosis was suspected to be *Legionella pneumophila* infection.

D Mycobacterial culture

Mycobacterial culture would be appropriate if TB infection was suspected.

70076

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 62-year-old man presents to the Emergency Department with a 5-day history of shortness of breath and a cough productive of green sputum. There are no other symptoms of note and he has no past medical history. He smokes 15–20 cigarettes per day.

On examination, his Glasgow Coma Scale score is 15/15. He has a respiratory rate of 32 breaths per minute, and oxygen saturations of 91% on air. His heart sounds are normal, with a heart rate of 120 bpm and a blood pressure of 101/62 mmHg. On chest auscultation, there are coarse crackles at the left lung base.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	126 g/l	135 - 175 g/l
White cell count (WCC)	14.5 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	10 × 10^9/l	1.5 - 7.0 × 10^9/l
Urea	6.5 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	130 μmol/l	50 - 120 μmol/l

Which factor in this patient’s presentation is the biggest prognostic indicator?

Your answer was incorrect

A Age

B Blood pressure

C Neutrophilia

D Smoking history

E Respiratory rate of 32 breaths/min

Explanation

E Respiratory rate of 32 breaths/min

This forms part of the CURB-65 criteria, an evidence-based scoring system used to estimate severity and guide management of patients with community-acquired pneumonia. The components of CURB-65 are: confusion (not present in this patient); urea concentration > 7 mmol/l (urea concentration here is < 7.0 mmol/l); respiratory rate ≥ 30 breaths/minute (this patient meets the criteria as his respiratory rate is 32 breaths per minute); systolic blood pressure < 90 mmHg (not met in this patient who has a systolic blood pressure of 110 mmHg); diastolic blood pressure < 60 mmHg; and age over 65 years (this patient is 62 years old). A CURB-65 score of zero corresponds to a 0.7% risk of mortality, whereas a score of 5 corresponds to a 57% risk. With a score of 1, this patient's risk is 3.2%. He would be suitable for outpatient management. If he scores 2 or more, he should be managed as an inpatient.

C Neutrophilia

Neutrophilia is not one of the features of CURB-65; the only CURB-65 criterion met by this patient is his respiratory rate which is 30 breaths per minute.

A Age

Age of ≥ 65 years counts towards the CURB-65 criteria. This patient is 62 years old, which does score a point on this scale.

B Blood pressure

This question relates to the CURB-65 criteria – these clinical criteria provide evidence-based guidance on the severity of community-acquired pneumonia and can be used to determine the appropriate management of a patient (admission vs outpatient care, and antibiotic therapy). Blood pressure does form part of the CURB-65 criteria; however, patients score a point if their systolic blood pressure is < 90 mmHg or diastolic blood pressure is < 60 mmHg. This patient has a systolic blood pressure of 110 mmHg, and therefore, this is not the correct answer. The only criterion on which this patient scores against the CURB-65 is his respiratory rate which is ≥ 30 breaths/minute.

D Smoking history

Although smoking is a risk factor for more severe respiratory disease, it does not form part of the CURB-65 severity criteria for pneumonia.

71041

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 62-year-old man presents to the Emergency Department with sudden-onset shortness of breath and right-sided pleuritic chest pain. His past medical history includes chronic obstructive pulmonary disease, for which he takes fluticasone and salmeterol (combined) and salbutamol, hypertension and ischaemic heart disease.

On examination, his blood pressure is 155/90 mmHg and his heart rate is 92 bpm and regular. On auscultation, he has diminished breath sounds over the right-hand side. Oxygen saturation is reduced at 90%. A chest X-ray (CXR) reveals a right-sided pneumothorax with a 1.5-cm rim of air.

An arterial blood gas is taken:

Investigation	**Result**	**Normal result**
pH	7.32	7.35–7.45
$p(O_2)$	8.9 kPa	11–14.4 kPa
$p(CO_2)$	6.9 kPa	4.6–6.4 kPa
Sodium (Na^+)	136 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Lactate	1.6 mmol/l	< 1.9 mmol/l

Which of the following is the most appropriate intervention?

Your answer was incorrect

A Admit for high-flow oxygen

B Discharge and review in 2–4 weeks

C Insert size 8–14 Fr chest drain

D Needle aspiration

E Non-invasive ventilation

Explanation

C Insert size 8–14 Fr chest drain

Insertion of a chest drain would be the most appropriate intervention for this patient. Although the rim of air is < 2 cm, the British Thoracic Society states that in the presence of significant symptoms (ie shortness of breath), a chest drain is preferred to air aspiration.

B Discharge and review in 2–4 weeks

Discharge and review in 2–4 weeks is recommended for patients with a primary pneumothorax of < 2 cm in size or after successful air aspiration of a primary pneumothorax.

A Admit for high-flow oxygen

Admission for high-flow oxygen is the correct course of action after successful air aspiration for secondary pneumothorax.

D Needle aspiration

This is the correct course of action for a secondary pneumothorax with no symptoms. However, given the shortness of breath and hypoxia, a chest drain is the preferred course of action.

E Non-invasive ventilation

Although this patient is hypoxic, with a raised $p(CO_2)$, increased pressures with non-invasive ventilation can transform a simple pneumothorax into a tension pneumothorax and non-invasive ventilation is therefore contraindicated.

72933

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society (BTS): BTS Pleural Disease Guideline 2010 Management of spontaneous pneumothorax
thorax.bmj.com/content/65/Suppl_2/ii18
(https://thorax.bmj.com/content/65/Suppl_2/ii18)

A 45-year-old woman presents with increasing wheeze and paroxysms of coughing. Most recently she has had two episodes of pneumonia in succession and an episode of haemoptysis. She is a non-smoker. Her chest X-ray is shown below. Biopsy of the lesion reveals a multitude of abnormal cells but no necrosis, occasional nuclear pleomorphism and absent mitoses.



Which of the following diagnoses fits best with this clinical picture?

Your answer was incorrect

A Small cell carcinoma of the bronchus

B Adenocarcinoma of the bronchus

C Squamous cell carcinoma of the bronchus

D Organised lung abscess

E Bronchial carcinoid

Explanation

E Bronchial carcinoid

The chest x-ray shows a solitary lesion in the left hilar region. When combined with the biopsy demonstrating occasional nuclear pleomorphism and absent or late mitoses the diagnosis fits the picture of ‘typical’ bronchial carcinoid. Endobronchial lesions fit with this clinical picture of symptoms of obstruction and bouts of pneumonia. Patients with isolated lesions should be considered for resection. If the tumour shows atypical features or is close to the resection margin, then patients should be followed up with serial chest X-rays. Aggressive carcinoid may be managed with chemotherapy; symptoms of the carcinoid syndrome may be managed with octreotide, but by the time patients present with carcinoid syndrome, lesions are irresectable.

Image source: https://commons.wikimedia.org/wiki/File:Carcinoid_tumor_L._hilum_(3931156757).jpg

C Squamous cell carcinoma of the bronchus

Squamous cell carcinoma usually arises centrally, in larger bronchi, with metastases to hilar lymph nodes early in its development. It is strongly associated with cigarette smoking.

A Small cell carcinoma of the bronchus

Small cell carcinoma is associated with marked nuclear pleomorphism and a large number of mitoses. Early metastasis is very common, and it is associated with exposure to tobacco smoke.

B Adenocarcinoma of the bronchus

Adenocarcinoma presents with atypical cells showing frequent mitoses, often potentially forming ducts or glands.

D Organised lung abscess

Nuclear pleomorphism would not be expected in an organised lung abscess; white cell infiltration, particularly with neutrophils, would be a prominent feature.

32436

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 44-year-old man presented with a rash on his legs and feeling generally unwell with malaise, weight loss and fevers. He also had recently noticed some loss of sensation over the top of his right foot, with the foot-dragging when he walked. He had been diagnosed with asthma by his general practitioner many years ago and used a salbutamol inhaler as required. He also had a constant watery nasal discharge, but no epistaxis. He was a non-smoker.

On examination, he had low-grade pyrexia and scattered wheeze in his chest. Heart sounds were normal. His abdomen was soft and non-tender, with no masses. He had a purpuric rash on his lower limbs and several tender subcutaneous nodules on his forearms. He had a right foot drop but no other neurological abnormalities.

Investigations reveal the following:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	106 g/l	135–175 g/l
White cell count (WCC)	10.9 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	390 × 10^9/l	150–400 × 10^9/l
Neutrophils	6.2 × 10^9/l	1.5–7.0 × 10^9/l
Eosinophils	2.2 × 10^9/l	0.04–0.4 × 10^9/l
Monocytes	1.6 × 10^9/l	0.0–0.8 × 10^9/l
Basophils	0.1 × 10^9/l	< 0.1 × 10^9/l
Mean corpuscular volume (MCV)	94 fl	76–98 fl
Erythrocyte sedimentation rate (ESR)	82 mm/h	1–20 mm/hour
Sodium (Na^+)	120 mmol/l	135–145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5–5.0 mmol/l
Alkaline phosphatase (ALP)	80 IU/l	30–150 IU/l
Alanine aminotransferase (ALT)	30 IU/l	5–30 IU/l
Urea	5.2 mmol/l	2.5–6.5 mmol/l
Gamma-glutamyl transferase (GGT)	42 IU/l	5–30 IU/l
Bilirubin	14 μmol/l	2–17 μmol/l
Creatinine (Cr)	110 μmol/l	50–120 μmol/l
Antinuclear antibody (ANA)	Negative	
Anti-neutrophil cytoplasmic antibody (pANCA)	Positive	
Urinalysis	Normal	

His chest x-ray is shown below.



What is the most likely diagnosis?

Your answer was incorrect

A	Systemic lupus erythematosus
B	Polyarteritis nodosa
C	Eosinophilic granulomatosis with polyangitis
D	Anti-GBM disease
E	Granulomatosis with polyangitis

Explanation

C Eosinophilic granulomatosis with polyangitis

This man presents with the classic triad of asthma, eosinophilia and systemic vasculitis. There is usually a prodromal period for many years consisting of allergic rhinitis and late-onset asthma, which can be difficult to control.

The chest X-ray here shows pulmonary infiltrates bilaterally. Other findings occasionally found include nodules, transient hilar lymphadenopathy and pleural or pericardial effusions. Mononeuritis multiplex can occur. Anti-neutrophil cytoplasmic antibody (ANCA) is positive in approximately 40% of cases. Renal involvement is rare. Treatment is with high-dose corticosteroids, azathioprine or cyclophosphamide.

Image source: https://radiopaedia.org/cases/eosinophilic-granulomatosis-with-polyangiitis-3?lang=gb (https://radiopaedia.org/cases/eosinophilic-granulomatosis-with-polyangiitis-3?lang=gb)

B Polyarteritis nodosa

pANCA is classically not positive in patients with PAN. In addition, the symptoms of significant asthma poorly responsive to conventional therapies, and eosinophilia, are inconsistent with a diagnosis of PAN.

A Systemic lupus erythematosus

Although SLE is associated with an erythematous rash on sun-exposed areas, the eosinophilia, negative ANA and positive pANCA would be unusual for a diagnosis of lupus.

D Anti-GBM disease

This presents with rapidly progressive pulmonary and kidney disease, accompanied by haemoptysis and kidney disease due to glomerulonephritis. Eosinophilia isn’t a feature, nor is mononeuritis multiplex.

E Granulomatosis with polyangitis

This is associated with pulmonary and renal vasculitis with haemoptysis and a progressive rise in serum creatinine. There is usually sinusitis. cANCA rather than pANCA is more likely to be positive, and eosinophils are not elevated.

9147

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

You review a 67-year-old man with severe chronic obstructive pulmonary disease (COPD) who stopped smoking 3 years ago. He has a history of myocardial infarction 4 years ago and chronic peripheral vascular disease. His medications include ramipril, indapamide, salbutamol and ipratropium.

Which of the following clinical features would best support prescription of long-term oxygen therapy (LTOT)?

Your answer was incorrect

A Pa(O_2) of 8.2 kPa

B Pitting oedema of the lower limbs

C Pa(CO_2) of 5.1 kPa

D FEV_1 (forced expiratory volume in 1 s) of 1.7 l

E History of angina

Explanation

B Pitting oedema of the lower limbs

According the British Thoracic Society guidelines on long-term oxygen therapy (LTOT) in regard to patients with stable chronic obstructive pulmonary disease (COPD) (see weblink below), a resting Pa(O_2) ≤7.3 kPa should be assessed for LTOT, which offers survival benefit when used for at least 15 h per day and improves pulmonary haemodynamics. LTOT should be offered to those patients with stable COPD and a resting Pa(O_2) ≤8 kPa with evidence of peripheral oedema, polycythaemia (haematocrit ≥55%) or pulmonary hypertension. Swollen ankles indicate the possibility of right heart failure, which may be occurring as a result of pulmonary hypertension.

Two measures are required to make the decision on therapy, and samples should be taken at least 3 weeks apart whilst the patient is stable and free of an exacerbation. This leads to a reduction in secondary polycythaemia, reduced sympathetic outflow and consequent salt and water retention, a reduction in cardiac arrhythmias, and improved sleep quality due to reduced hypoxia-induced sleep arousals.

D FEV_1 (forced expiratory volume in 1 s) of 1.7 l

Spirometry findings do warrant criteria for oxygen therapy and do not indicate hypoxia.

A Pa(O_2) of 8.2 kPa

$Pa(O_2)$ >8kPa should not lead to a prescription of LTOT.

C Pa(CO_2) of 5.1 kPa

Although LTOT should be ordered for patients with resting hypercapnia (if they fulfil all other criteria for LTOT), this gentleman does not have resting hypercapnia.

E History of angina

Although it can be used in chronic heart failure, history of angina is not an appropriate use of LTOT. Anti-anginal medication or a coronary angiogram would be required if patient has recurrent episodes of angina.

32441

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Guidelines for Oxygen Use
brit-thoracic.org.uk/about-us/pressmedia/2017/new-bts-quality-standards-for-home-oxygen-use-in-adults/#
(https://www.brit-thoracic.org.uk/about-us/pressmedia/2017/new-bts-quality-standards-for-home-oxygen-use-in-adults/#)

A 35-year-old African female presents with a 6-months history of night sweats, shortness of breath on exertion and dry cough. She also gives a history of weight loss and arthralgia over the same period of time. She had been intermittently unwell for 1 year before the development of these symptoms. She recalls a painful rash on her shins 3 months ago, which resolved spontaneously. She gives no history of eye problems, polyuria or polydipsia. She denies chest pain or tightness. She denies any past medical problems and is not on any medications. On examination she appears well at rest. There is no joint swelling or redness. Cardiovascular, respiratory and abdominal examinations are normal. A high resolution CT of the thorax has revealed bilateral hilar lymphadenopathy with evidence of early fibrosis.

Which investigation is most likely to help confirm the diagnosis?

Your answer was incorrect

A Serum angiotensin-converting enzyme (ACE) concentration

B Pulmonary function tests

C Quantiferon gamma

D Bronchoalveolar lavage

E Transbronchial lung biopsy

Explanation

E Transbronchial lung biopsy

In this patient of African ethnicity with a description of probable erythema nodosum and a dry cough, the most likely diagnosis is sarcoidosis. Sarcoidosis is a multi-system granulomatous disorder of unknown aetiology and is characterised pathologically by the presence of non-caseating granulomas in involved organs. As with the diagnosis of tuberculosis, a tissue diagnosis is the only completely specific way that a confirmed diagnosis can be made, as all the other diagnostic tests are non-specific to sarcoidosis.

C Quantiferon gamma

This is an African patient with a dry cough and what sounds like erythema nodosum. Whilst tuberculosis would be in the differential diagnosis, given that she is having night sweats and is systemically unwell, a quantiferon gamma would not confirm or refute a diagnosis of active tuberculosis, as it cannot differentiate between active or latent disease.

A Serum angiotensin-converting enzyme (ACE) concentration

In this patient of African ethinicity with a description of probable erythema nodosum and a dry cough, the most likely diagnosis is sarcoidosis. Serum angiotensin-converting enzyme (ACE) activity is elevated in 60% of patients, presumably reflecting macrophage activity. It can also be elevated in patients with histoplasmosis, acute miliary tuberculosis (TB), Gaucher's hepatitis or lymphoma, and is non-specific. Liver disease slows the metabolic excretion of serum ACE and results in increased ACE activity.

B Pulmonary function tests

In this patient of African ethnicity with a description of probable erythema nodosum and a dry cough, the most likely diagnosis is sarcoidosis. However, pulmonary function tests are non-specific in making this diagnosis. Pulmonary function tests show restriction, decreased compliance and impaired diffusing capacity. However, serial pulmonary function tests are important for assessing disease progression and guiding treatment.

D Bronchoalveolar lavage

In this patient of African ethnicity with a description of probable erythema nodosum and a dry cough, the most likely diagnosis is sarcoidosis. Bronchoalveolar lavage (BAL) can be used as an adjunctive measure to support the diagnosis of sarcoidosis. Bronchoalveolar lavage shows lymphocytosis in most patients with active sarcoidosis, but is rarely indicated because patients with hypersensitivity pneumonitis show similar lymphocytosis.

32403

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

A 45-year-old patient presents to his GP with dyspnoea and cough over the past week. He has also noticed recurrent nose bleeds and crusting around his nose over the past month. He is a non-smoker, has no past medical history and is not on any medications.

Spirometry is performed and reveals the following:

Investigation	Result
Forced expiratory volume in 1 second (FEV_1)	80% predicted
Forced vital capacity (FVC)	65% predicted
Total lung capacity (TLC)	70% predicted
Lung CO transfer factor (T_{LCO})	120% predicted factor
CO transfer coefficient (K_{CO})	120% predicted coefficient

Urine dip: protein ++, blood +

Blood tests reveal the following:

Investigation	Result	Normal Value
Haemoglobin (Hb)	112 g/l	135–175 g/l
White cell count (WCC)	11.2 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	540 × 10^9/l	150–400 × 10^9/l
Mean corpuscular volume (MCV)	90 fl	76–98 fl
Estimated glomerular filtration rate (eGFR)	41	

Which is the most likely cause of the lung function tests?

Your answer was incorrect

A Pulmonary haemorrhage

B Pneumonectomy

C Left-to-right shunt

D Sarcoidosis

E Neuromuscular defect

Explanation

A Pulmonary haemorrhage

The spirometry shows a restrictive picture (FEV_1/FVC >80%). This suggests that there is a restrictive cause of the shortness of breath, such as haemorrhage, sarcoidosis, pneumonectomy or reduced work of the respiratory muscles due to a neuromuscular defect. Pulmonary haemorrhage would be a cause of increased K_{CO} and hence is the most likely diagnosis when taken with the history of nose bleeds, making a vasculitis such as Granulomatosis with polyangitis (GPA) the likely underlying diagnosis. GPA is also associated with collapse of the nasal bridge, leading to a ‘saddle nose’ deformity.

D Sarcoidosis

While this would result in a restrictive pattern of PFTs, we are not given any history of other symptoms such as erythema nodosum, and this does not explain the increased K_{CO} as adequately as pulmonary haemorrhage.

B Pneumonectomy

This would be expected to result in a reduced transfer factor due to loss of significant volume of alveolar membrane, with normal transfer coefficient assuming the residual lung is normal.

C Left-to-right shunt

While this would cause an increased transfer factor, it does not explain the nose bleeds as adequately as a diagnosis of underlying vasculitis. We are also not given any history of murmurs such as ASD, VSD or PDA on examination, nor features of underlying pulmonary hypertension.

E Neuromuscular defect

While this would result in a restrictive pattern of PFTs, we are not given any other features to suggest a neurological aetiology such as motor or sensory deficit. It also fails to adequately explain the patient’s epistaxis.

7161

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 31-year-old asthmatic woman was seen in the Respiratory Outpatient Clinic. She said she had been generally well although she had not been sleeping well at night due to coughing. She had also noticed that when playing tennis she was getting wheezy and requiring her salbutamol inhaler more frequently – at least 4 times a week. She was taking budesonide 200 micrograms two puffs twice a day. She had no significant past medical history. She smoked 10 cigarettes/day and had done so for 2 years. Auscultation of her chest revealed a mild expiratory wheeze. Examination was otherwise remarkable.

PEF	80% predicted
FEV_1	85% predicted
FVC	90% predicted

She is treated successfully and goes home.

Her symptoms improve and she goes back to playing regular tennis without a problem. Three months later she was feeling so much better she decided not to renew her prescription for her inhalers.

Two weeks following this she developed a chest infection and became acutely short of breath. On examination she is unable to speak full sentences. Respiratory rate is 30/min, pulse 115/min, BP 100/60 mmHg.

Auscultation of her chest reveals an expiratory wheeze throughout. Her initial investigations including chest x-ray are shown below.



PEFR	50 l/min
ABGs on air:	
pH	7.45
pCO_2	4.06 kPa
pO_2	12 kPa
Bicarbonate	21.5 mmol/l

She does not improve with treatment. Her chest examination now reveals quiet breath sounds. Repeat arterial blood gases on oxygen show:

pH	7.33
pCO_2	6.5 kPa
pO_2	12 kPa
Bicarbonate	25.1 kPa

What would you do next?

Your answer was correct

A Give intravenous antibiotics

B Give more regular nebulisers

C Refer to ICU for intubation

D Start non-invasive ventilation

E Trial of helium/oxygen

Explanation

C Refer to ICU for intubation

The patient deteriorates and develops signs of life-threatening asthma – a silent chest, a pCO_2 >6kPa and a fall in pH. Other signs of life-threatening asthma include: confusion, exhaustion or coma, bradycardia, hypotension, PEFR <30% predicted or best, and severe hypoxaemia (PaO_2 <8 kPa) despite oxygen therapy. This patient now needs intubation and transfer to the ICU. Non-invasive ventilation is not used in asthma at present and it is unlikely ever to replace intubation in life-threatening asthma.

The chest x-ray here is normal and it is likely the underlying cause of her exacerbation is a viral infection and poor compliance with medication.

Image source: https://commons.wikimedia.org/wiki/File:Chest_Xray_PA_3-8-2010.png

A Give intravenous antibiotics

The priority here is to ensure the patient is receiving adequate respiratory support. There is an absence of features indicating a bacterial infection, including a normal chest x-ray; this lady's exacerbation is likely to have been triggered by a virus in conjunction with non-adherence to her asthma medication.

B Give more regular nebulisers

This lady is now in acute type 2 respiratory failure with respiratory acidosis. This is near-fatal asthma and she requires immediate intensive care assessment for intubation and ventilation.

D Start non-invasive ventilation

Non-invasive ventilation is not an appropriate treatment for asthma. Invasive ventilation should be used in patients with acute respiratory failure secondary to as asthma exacerbation.

E Trial of helium/oxygen

Helium/oxygen is a treatment for vocal cord dysfunction – this lady is having a near-fatal asthma exacerbation

70125

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society. 2019. BTS/SIGN British guideline on the management of asthma.
brit-thoracic.org.uk/quality-improvement/guidelines/asthma/
(https://www.brit-thoracic.org.uk/quality-improvement/guidelines/asthma/)

Asthma

Media Expanded explanations Flashcards

An 18-year-old man is admitted to the Emergency Department drowsy and confused. He has recently complained of symptoms of an upper respiratory tract infection. He has a past medical history of Duchenne’s muscular dystrophy. Over the past few weeks, he noted that he seemed to be suffering intermittent morning headaches and intermittent confusion.

On examination, he is drowsy but able to obey simple commands. His respiratory rate is 13 breaths per minute. Air entry is poor on auscultation, with crackles at the left base.

Investigations:

Investigation	Result	Normal value
pH	7.3	7.35–7.45
pCO_2	7.0 kPa	4.6–6.0 kPa
pO_2	7.0 kPa	10.5–13.5 kPa

Which of the following would represent the most appropriate management in this case?

Your answer was incorrect

A 60% oxygen therapy

B 35% oxygen therapy

C 28% oxygen therapy

D Intubation and ventilation

E Non-invasive ventilation

Explanation

E Non-invasive ventilation

This man has evidence of chronic carbon dioxide retention, as evidenced by his morning headaches, intermittent confusion and drowsiness. A further indication of respiratory muscle, particularly diaphragmatic and weakness may be indicated by breathlessness during swimming. Chronic hypoventilation has likely led to an opportunistic respiratory tract infection associated with his cold. At present, he has mild respiratory acidosis, and the situation may be remediable with non-invasive ventilation and antibiotics. Since Duchenne's muscular dystrophy is progressive, intubation and ventilation should be carefully considered.

B 35% oxygen therapy

Like higher percentages of oxygen, 35% oxygen will not reverse the impact of hypoventilation with respect to respiratory acidosis. Therefore, it would not be appropriate here.

A 60% oxygen therapy

Therapy with 60% oxygen will have a very limited impact, if any, on improving this patient’s hypoxia due to his hypoventilation and worsening acidosis. Therefore, this is not an appropriate option for this patient.

C 28% oxygen therapy

Therapy with 28% oxygen will not improve oxygenation and will not reverse the impact of hypoventilation, making it unlikely to be appropriate for this patient.

D Intubation and ventilation

Intubation and ventilation should be carefully considered in the context of Duchenne’s muscular dystrophy. It is a progressive condition and the risk is that the patient may become un-weanable from the ventilator. As such, non-invasive ventilation is preferred here. Ideally, patients should consider an advanced directive before they become unwell so that their wishes with respect to invasive therapies are already established.

18554

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Ventilatory Support

Expanded explanations

An 18-year-old woman comes to the Emergency Room complaining of extreme lethargy. A couple of weeks earlier she had suffered from a nasty bout of gastroenteritis after a takeaway meal. She complains of abnormal sensation and numbness in her fingers, and weakness of her lower legs. She is generally hypotonic and has difficulty standing. On examination there is a postural drop in BP of 20 mmHg to 105/70 mmHg, and she has a resting tachycardia of 90 BPM. No deep tendon reflexes can be elicited in her lower limbs. Only medication of note includes the oral contraceptive pill.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	121 g/l	115–155 g/l
White cell count (WCC)	6.9 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	230 × 10^9/l	150–400 × 10^9/l
Sodium (Na$^+$)	141 mmol/l	135–145 mmol/l
Potassium (K$^+$)	4.8 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	140 μmol/l	50–120 μmol/l
Glucose	7.2 mmol/l	3.5–5.5 mmol/l
Thyroid stimulating hormone (TSH)	1.8 μU/l	0.17–3.2 μU/l

Given the likely diagnosis, which of the following is the most important way to monitor the progress of her condition?

Your answer was incorrect

A Lumbar puncture

B Anti-GQ1b antibodies

C FVC

D FEV$_1$

E Creatine kinase levels

Explanation

C FVC

This woman has a history consistent with Guillain–Barré syndrome (GBS), which is known to occur after an episode of bacterial gastroenteritis. Typical symptoms seen here include paraesthesia in the fingers, with progressive lower limb weakness. Autonomic signs such as tachycardia and postural hypotension also occur. FVC is the most helpful investigation in assessing prognosis. Respiratory failure, including supervening bacterial pneumonia, is the most important cause of morbidity/mortality associated with the condition.

B Anti-GQ1b antibodies

Anti-GQ1b antibodies are found in the Miller Fisher variant of GBS, which is associated with ophthalmoplegia; they are not predictive of outcomes in the condition.

A Lumbar puncture

Although an elevation in CSF protein may be seen, this may be normal in the early stages of the condition and therefore doesn't rule out the diagnosis.

D FEV_1

FEV_1 is a marker of airways obstruction, and is much more useful in the monitoring of asthma and COPD.

E Creatine kinase levels

CK is a marker of disease activity in dermatomyositis, and is not useful in monitoring of Guillain–Barré syndrome.

21025

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Respiratory Failure

Expanded explanations

A 42-year-old publican is admitted with increasing shortness of breath which is relieved by lying flat. He has been seen on one previous occasion with a haematemesis and admits to drinking 6–8 pints of strong lager per day. He also smokes 20 cigarettes per day. On examination he looks unwell; he has signs of chronic liver disease including clubbing, spider naevi, scratch marks, swollen ankles and ascites. He is tender in the right upper quadrant. Respiratory examination is suggestive of bilateral pleural effusions. His BP is 95/60 mmHg, and his pulse is 100/min.

Investigations reveal the following:

Investgation	Result	Normal value
Haemoglobin (Hb)	100 g/l	135–175 g/l
White cell count (WCC)	5.6 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	78 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	3.8 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	90 μmol/l	50–120 μmol/l
Alanine aminotransferase (ALT)	140 U/l	5–30 IU/l
Bilirubin	110 μmol/l	2–17 μmol/l
Albumin	22 g/l	35–55 g/l
pO_2	7.9 kPa	10.5–13.5 kPa
pCO_2	3.9 kPa	4.6–6.0 kPa

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Pulmonary embolus

B Cardiac failure

C Hepatopulmonary syndrome

D Chronic obstructive pulmonary disease (COPD)

E Pulmonary fibrosis

Explanation

C Hepatopulmonary syndrome

Given the history of significant liver disease, a link between this and his chest pathology, hepatopulmonary syndrome is more likely than a pulmonary embolus. Equally, there are no signs suggestive of either COPD or pulmonary fibrosis. Hepatopulmonary syndrome is due to arteriovenous shunting and is potentially progressive and life-threatening. It is associated with gross dilatation of pulmonary pre-capillary and capillary vessels; hypoxia may be exacerbated by sitting upright because of preferential perfusion of lung bases (with poor ventilation of these areas in the upright position). Lung bases are better ventilated when the patient lies flat. Hypoxia may be further worsened by the presence of significant pleural effusions. Nitric oxide is thought to play a role in the condition, largely as a result of studies in rodent models, although this is yet to be confirmed in human studies. The syndrome may be corrected by liver transplantation, and is adequate reason for expedited referral. Platypnoea is related to this abnormal pulmonary vasodilation, which leads to inappropriate shunting. Bubbles can be seen appearing late in the left atrium when agitated saline is injected, and the atrium is observed on echocardiography.

D Chronic obstructive pulmonary disease (COPD)

Bilateral pleural effusions would not fit with a diagnosis of COPD, and chronic COPD would usually be associated with an elevated bicarbonate because of metabolic compensation, not seen here.

A Pulmonary embolus

Although hypoxia and shortness of breath could be associated with pulmonary embolism, it’s less likely given decreased hepatic synthetic function and likely abnormal clotting, coupled with obvious decompensated cirrhosis, meaning hepatopulmonary syndrome is a much more obvious cause.

B Cardiac failure

Significant left ventricular failure is more likely to be associated with type 2 respiratory failure rather than the low CO_2 seen here.

E Pulmonary fibrosis

Pulmonary fibrosis is associated with crackles on auscultation, and not with the pleural effusions seen here. Shortness of breath would also not be improved by lying flat.

20999

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A young paint-sprayer who works in a car shop complains of shortness of breath and tiredness over last 2 months. He was prescribed inhalers by GP, but eventually needed a course of steroids for resolution of symptoms. It was found that after a 2-week holiday to Marbella his symptoms had resolved. He then saw GP on return as his symptoms had once again worsened. On examination his BP was 122/72 mmHg, pulse was 75/min and regular and there was scattered wheeze on auscultation of the chest. A peak flow diary revealed a significant reduction, which progressed during the working week and then improved over the weekend.

Which of the following would be the most appropriate intervention?

Your answer was incorrect

A Change of career

B Oral Monteleukast during the working week

C Regular inhaled beclomethasone

D Regular inhaled salmeterol/fluticasone

E Review of the work environment to ensure exposure to paint fumes is minimised

Explanation

E Review of the work environment to ensure exposure to paint fumes is minimised

The most likely cause of this man's symptoms is occupational asthma. A number of substances such as paint spray (isocyanates) and soldering kit (colophony fumes) used on cars are possible causes of the illness. In the first instance his working environment should be examined to ensure that exposure to fumes is minimised, both for his own health and for others in the same work environment. If changes do not improve his symptoms then he should be offered alternative work, as a commitment to regular asthma therapy is suboptimal in this situation.

C Regular inhaled beclomethasone

Regular inhaled beclomethasone is the initial intervention for patients with asthma not related to precipitating factors in the work environment and who do not respond to PRN salbutamol.

A Change of career

A change of career is only appropriate if despite minimising exposure to paint fumes, symptoms persist. A career change is preferable versus starting regular asthma therapy.

B Oral Monteleukast during the working week

Oral Monteleukast is a treatment option for patients who fail to respond to high-dose inhaled corticosteroids or to the combination of inhaled corticosteroids and a long-acting beta agonist. It should not be used in lieu of a review of the working environment.

D Regular inhaled salmeterol/fluticasone

Combination inhaled corticosteroid and long-acting beta agonist preparations are used for patients who fail to respond to inhaled corticosteroids alone.

32433

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Occupational Asthma

Expanded
explanations

A 45-year-old woman complains of shortness of breath, occasional fevers and mild weight loss, all getting worse over the past months. She does not complain of a cough but remembered that she had coughed once and produced a twig-shaped mucoid sputum mass. She has no haemoptysis. She has no past medical history and is on no medications. Her chest X-ray reveals bilateral, perihilar, dense airspace shadowing. She went on to have a HRCT of her thorax, which showed a ‘crazy paving’ pattern of extensive, dense, white infiltrates. Her spirometry was a restrictive pattern with reduced total lung capacity. She also had a bronchoscopy and lavage, which revealed periodic acid–Schiff (PAS)-positive proteinaceous fluid and elevated levels of surfactant proteins A and D.

What is the most likely diagnosis?

Your answer was incorrect

A Chronic pulmonary eosinophilia

B Sarcoidosis

C Extrinsic allergic alveolitis

D Pulmonary alveolar proteinosis

E *Pneumocystis jirovecii* infection

Explanation

D Pulmonary alveolar proteinosis

Pulmonary alveolar proteinosis (PAP) is a rare diffuse lung disease in which the alveolar sacs become filled with protein-rich fluid that characteristically stain for PAS, a protein derived from surfactant. It usually affects people aged 20–60 years who have no previous lung disease, men more frequently than women. It can be primary or secondary, the latter being related to infections with, for example, *Pneumocystis jirovecii* or atypical mycobacteria, or to immunosuppressants, organic dusts and haematological malignancy. It usually presents with shortness of breath and mild constitutional symptoms. Diagnosis is based on bronchial lavage or biopsy with PAS-positive stains and increased presence of surfactant proteins A and D. HRCT shows a classic picture of dense infiltrates, often referred to as crazy paving. Spirometry shows a restrictive pattern with reduced lung capacity and reduced carbon monoxide diffusion. Most patients do not require treatment unless their shortness of breath is disabling. Treatment is by washing the alveoli out with salt solution. This can be done by bronchoscopy or under general anaesthesia through the trachea. If both lungs need washing then they are done about 5 days apart. The number of washings depends on symptoms. The prognosis is good.

C Extrinsic allergic alveolitis

Hypersensitivity pneumonitis occurs in response to an antigen, such as occurs in, for example, bird fancier’s lung, etc. We are not given any history of exposure that would explain this diagnosis, and the CT findings are not in keeping, with a ‘ground glass’ pattern being more common.

A Chronic pulmonary eosinophilia

Histology would be expected to show eosinophilic and lymphocytic infiltration of the alveoli and interstitium, with fibrosis. HRCT typically shows non-segmental areas of consolidation, with a peripheral predilection.

B Sarcoidosis

This is classically associated histologically with the formation of non-caseating granulomas. We are not given any history of cutaneous features such as erythema nodosum. The CT findings also do not fit, with sarcoidosis commonly showing hilar lymphadenopathy and perilymphatic nodule distribution.

E *Pneumocystis jirovecii* infection

This is uncommon outwith the context of HIV infection or other cause of immunocompromise.

7163

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301

Responses - % Correct:	27%

A 62-year-old male smoker of 30 cigarettes per day presents with symptoms of cough and dyspnoea, accompanied by vague, non-specific chest pain. There is general malaise and joint pain, which has been relieved by using diclofenac as required (prn), and his cough has not been improved by a salbutamol inhaler prn.

His chest X-ray is shown below.



By James Heilman, MD [CC BY-SA 4.0 (https://creativecommons.org/licenses/by-sa/4.0)], from Wikimedia Commons

There is a non-specific increase in immunoglobulins and raised serum angiotensin-converting enzyme (ACE) level.

Which of the following diagnoses fits best with this clinical picture?

Your answer was incorrect

A Bronchial carcinoma

B Sarcoidosis

C Multiple myeloma

D Lymphoma

E Tuberculosis

Explanation

B Sarcoidosis

In this case, there is evidence of bilateral hilar lymphadenopathy on chest X-ray. More than 90% of patients with sarcoidosis have abnormal chest X-ray findings. Both immunoglobulins and serum ACE levels can be raised in active sarcoidosis. Renal function, liver function tests and serum calcium levels should also be assessed at initial appointment. High-resolution computed tomography (CT) scanning may demonstrate micronodules in a subpleural and bronchoalveolar distribution, fissural nodularity and bronchial distortion. Irregular linear opacities and ground-glass shadowing may also be seen. Air trapping due to endobronchial granulomas and 'honeycombing' may also be seen. Kveim tests are no longer performed due to perceived danger of infection. If the CT scan is diagnostic, then mediastinoscopy, bronchoscopy or biopsies can often be avoided.

C Multiple myeloma

Multiple myeloma is associated with the appearance of a paraprotein band on immunoglobulin testing, rather than the non-specific increases seen here.

A Bronchial carcinoma

Bronchial carcinoma is more usually associated with a unilateral mass seen on chest X-ray, with lymph node spread to the ipsilateral side.

D Lymphoma

Lymphoma is associated with hilar lymphadenopathy, although the dry cough and elevated serum ACE level are more consistent with a diagnosis of sarcoidosis.

E Tuberculosis

Tuberculosis is less likely, given the hilar lymphadenopathy is bilateral and no risk factors for tuberculosis are proffered in the clinical scenario.

32438

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

This high-resolution computed tomography (CT) scan is from a 42-year-old man with chronic dyspnoea and cough. PaO_2 is 8.9 kPa on air on arterial blood gas testing.



What is the most appropriate management strategy for this condition?

Your answer was correct

A Oral corticosteroids

B Antibiotics and regular postural drainage

C Long-term home oxygen therapy

D Intravenous cyclophosphamide

E Anti-mycobacterial therapy

Explanation

B Antibiotics and regular postural drainage

This computed tomography (CT) scan shows the typical appearance of bronchiectasis, with dilated, thickened bronchi. Causes of this condition include cystic fibrosis, post-infective TB, immune deficiency and Kartagener syndrome. Management includes daily postural drainage, antibiotic therapy for infective exacerbations (which should be guided by sputum culture) and bronchodilators. Surgical resection for localised areas of bronchiectasis is rarely indicated. Heart lung transplantation is an option in end-stage disease with cor pulmonale. Colonisation with *Pseudomonas aeruginosa* is a common problem. Complications of bronchiectasis include massive haemoptysis, empyema, pneumothorax and metastatic infections (e.g. brain abscess).

A Oral corticosteroids

In this patient, if the CT had demonstrated evidence of fibrosis with honeycombing, this would have indicated that he had interstitial lung disease which would be likely to be susceptible to steroid treatment.

C Long-term home oxygen therapy

If the CT scan had demonstrated features of idiopathic pulmonary fibrosis, with bilateral lower zone reticulo-nodular shadows and reduced lung volume, then a diagnosis of idiopathic pulmonary fibrosis would be likely and long-term home oxygen could be considered as a supportive measure.

D Intravenous cyclophosphamide

If the CT had demonstrated bilateral hilar lymphoadenopathy with pulmonary infiltrates and fibrosis, then a diagnosis of sarcoidosis could be considered, for which IV cyclophosphamide is a treatment option in severe disease.

E Anti-mycobacterial therapy

Anti-mycobacterial therapy is used to treat TB. TB would be suggested on a CT scan if it demonstrated consolidation and cavitating lesions, but would require further immunological or microbiological testing to confirm diagnosis.

9154

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 71-year-old man presents to the Emergency Department. He has suffered from a dry cough and diarrhoea over the past few days and is now drowsy and confused. According to his wife, they have recently returned from a winter holiday to Spain where they spent three weeks staying in a hotel. He has a past history of hypertension and inferior myocardial infarction, although he has had no ongoing symptoms of ischaemic heart disease over the past few years. He has no known allergies.

Examination reveals a blood pressure of 103/82 mmHg and a heart rate of 95 bpm and regular. His temperature is 38.4 °C. There is right-sided wheeze on auscultation and evidence of right-sided consolidation. His respiratory rate is elevated at 34 breaths per minute, and he looks unwell. His abdomen is soft, and there is minor left-sided tenderness on palpation.

Investigations:

Investigations	Results	Normal Value
Haemoglobin (Hb)	130 g/l	135–175 g/l
White cell count (WCC)	14.1 × 10^9/l	4–11 × 10^9/l
Platelets (PLT)	171 × 10^9/l	150–400 × 10^9/l
C-reactive protein (CRP)	178 mg/l	0–10 mg/l
Sodium (Na^+)	130 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Urea	8 mmol/l	2.5–6.5 mmol/l
Creatinine	112 × µmol/l	50–120 µmol/l
Glucose	6.1 mmol/l	3.5–5.5 mmol/l

Chest X-ray shown is taken:



What is the treatment of choice for this patient, given the likely diagnosis?

Your answer was incorrect

A	Azithromycin 500 mg intravenous once a day (OD) for ten days
B	Clarithromycin 500 mg oral (PO) OD for 14 days
C	Ciprofloxacin 400 mg IV three times a day (TDS) for ten days
D	Doxycycline 200 mg IV, followed by 100 mg twice a day for 21 days
E	Levofloxacin 750 mg IV OD for 14 days

Explanation

E Levofloxacin 750 mg IV OD for 14 days

The history of pneumonia acquired following a trip to Spain, coupled with hyponatraemia, diarrhoea and lobar pneumonia, raises the possibility of Legionella infection. Urinary antigen testing is a rapid potential route to confirm the diagnosis (commercial kits being widely available). Syndrome of inappropriate antidiuretic hormone is seen and potentially accounts for the low sodium seen here. Radiographic appearances can lag behind the clinical picture and are typically lobar (as here) or multilobar pneumonia, commonly with pleural effusion. Macrolides or quinolones are the antibiotics of choice, and a course of at least ten days is recommended. The

British Thoracic Society recommends treatment with fluoroquinolones for severe cases requiring hospitalisation. The use of levofloxacin has the best evidence and therefore, this is preserved for severe cases; the treatment course should be at least 14 days. In cases where there is a poor response to initial antibiotic therapy, a combination of fluoroquinolone and macrolide can be used. Mortality is approximately 10% and is higher in patients who are immunocompromised or older with significant co-morbidities.

Image source: https://commons.wikimedia.org/wiki/File:Pneumonia_of_Right_Lung_Middle_Lobe.jpg

C Ciprofloxacin 400 mg IV three times a day (TDS) for ten days

Ciprofloxacin is another fluoroquinolone antibiotic. There is less evidence for ciprofloxacin and therefore, its use is reserved for milder cases, which makes it an incorrect answer in this case, as the patient has severe pneumonia. The typical treatment course for ciprofloxacin is 7–10 days.

A Azithromycin 500 mg intravenous once a day (OD) for ten days

Azithromycin is a macrolide antibiotic. Macrolides can be used as a first-line treatment for Legionella infection. However, in severe cases (such as this patient with a CURB-65 score of 4), treatment with fluoroquinolone is recommended given for 7-14 days.

B Clarithromycin 500 mg oral (PO) OD for 14 days

Clarithromycin is a macrolide antibiotic that can be used as a first-line treatment for Legionella infection. Clarithromycin should be given for 14–21 days.

D Doxycycline 200 mg IV, followed by 100 mg twice a day for 21 days

Doxycycline is a tetracycline antibiotic. Tetracyclines may be used as an alternative treatment option where the patient has a known allergy to macrolides or fluoroquinolones. However, this patient does not have any known allergies. Treatment with doxycycline should be for 14–21 days.

72978

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society (BTS): Guidelines on pneumonia
brit-thoracic.org.uk/quality-improvement/guidelines/pneumonia-adults/
(https://www.brit-thoracic.org.uk/quality-improvement/guidelines/pneumonia-adults/)

Pneumonia

Media

Expanded explanations

Flashcards

A 32-year-old man, who has been human immunodeficiency virus (HIV)-positive for 5 years and started highly active antiretroviral therapy several months ago, presented with several days of malaise, rigors, a cough productive of blood-streaked green sputum and pleuritic chest pain. His latest CD4 count was 320 × 10^6/l several weeks ago.

On examination, he was unwell, with pyrexia of 39 °C, pulse of 110 bpm, blood pressure of 100/70 mmHg and oxygen saturation of 91% on air. On auscultation of his chest, he had bronchial breathing and coarse crepitations at the left lower zone.

His chest X-ray is shown below.



What is the most likely diagnosis?

Your answer was incorrect

A *Pneumocystis jiroveci* pneumonia (PCP)

B Streptococcal pneumonia

C Staphylococcal pneumonia

D *Mycobacterium avium intacellulare* infection

E Lymphoid interstitial pneumonitis

Explanation

B Streptococcal pneumonia

This man has community-acquired pneumonia, 60–75% of which are caused by *Streptococcus pneumoniae*. Disseminated bacterial sepsis is common in HIV patients and may be seen in the early stages of infection, before a significant decline in the CD4 count. The chest X-ray shows the consolidation of left lower lobe.

Image source- https://radiopaedia.org/cases/17760/studies/17517?lang=us&source_of=https%3A%2F%2Fradiopaedia.org%2Farticles%2Fleft-lower-lobe-consolidation%3Flang%3Dus (https://radiopaedia.org/cases/17760/studies/17517?lang=us&source_of=https%3A%2F%2Fradiopaedia.org%2Farticles%2Fleft-lower-lobe-consolidation%3Flang%3Dus)

C Staphylococcal pneumonia

Staphylococcal pneumonia tends to occur post-influenza infection or in intravenous drug users, and causes cavitation and multiple lung abscesses. There is no indication of previous influenza as a driver for infection here.

A *Pneumocystis jiroveci* pneumonia (PCP)

P. jiroveci (previously *carinii*) pneumonia and *Mycobacterium avium intracellulare* infection are acquired immune deficiency syndrome (AIDS)-defining conditions and require a CD4 count of < 200 × 10^9/l to develop. PCP also usually develops as a dry cough, with desaturation on exercise and patchy interstitial shadowing on chest X-ray.

D *Mycobacterium avium intacellulare* infection

Atypical mycobacterial infection, like PCP, occurs in patients with a significant fall in their CD4 count and is an AIDS-defining condition.

E Lymphoid interstitial pneumonitis

Lymphoid interstitial pneumonitis is due to infiltration of the alveolar tissue by lymphocytes and plasma cells and presents with dry cough, dyspnoea and reticulo-nodular shadowing on chest X-ray. It is uncommon in adults but well described in paediatric HIV infection.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total: 301

Responses - % Correct: 27%

Pneumonia

Media

Expanded explanations

Flashcards

An 82-year-old woman living in her own home presents with a 3-day history of confusion, as noted by her carer. She has had a purulent cough for several days. She has never smoked.

Her other health problems include controlled hypertension, osteoarthritis and venous leg ulcers, which limit her mobility.

She attends a day centre 3 days per week, where two other patients have recently been treated for pneumonia with according to the general practitioner (GP), although you have no further details regarding this.

On examination, she is mildly confused and drowsy, with mild pyrexia of 37.8 °C. Chest examination is abnormal. Routine bloods reveal a neutrophilia with CRP of 220.

Her chest X-ray below:



What is the most likely causative organism?

Your answer was incorrect

A *Streptococcus pneumoniae*

B *Legionella pneumophilia*

C *Mycoplasma*

D *Staphylococcus aureus*

E *Haemophilus influenzae*

Explanation

A *Streptococcus pneumoniae*

S. pneumoniae is the commonest cause of community-acquired pneumonia in the elderly. Bacteraemic *S. pneumoniae* pneumonia is the number one cause of mortality in community-acquired pneumonia, representing up to 80% of cases and 70% of all deaths. The short history of onset and right mid and lower lobe consolidation, seen on chest x-ray, are entirely consistent with *S. pneumoniae* versus an atypical cause.

Image source: https://commons.wikimedia.org/wiki/File:X-ray_lung_consolidation.jpg

C *Mycoplasma*

Mycoplasma presents in epidemics often associated with institutionalisation, for instance amongst recruits in an army barracks. The acute course and lobar pneumonia seen here are more consistent with *S. pneumoniae*.

B *Legionella pneumophilia*

Legionella is associated with a greater burden of confusion which may be linked to hyponatraemia, and other features such as diarrhoea and elevated transaminases.

D *Staphylococcus aureus*

S. aureus pneumonia is typically described as following an episode of influenza, and leads to focal consolidation with areas of cavitation.

E *Haemophilus influenzae*

H. influenzae pneumonia is less common in percentage terms than streptococcal infection, and is more likely to be seen in patients with pre-existing chest disease.

6337

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

You are seeing a patient on the surgical wards as a medical referral. He has had a lymph node dissected from his neck for possible recurrent head and neck cancer. On his postoperative chest X-ray, he has a moderate pleural effusion, which on aspiration drains white, milky fluid.

What is the likely diagnosis of the cause of the effusion?

Your answer was correct

A Empyema

B Mesothelioma

C Chylous effusion

D Reactive effusion

E Malignant effusion

Explanation

C Chylous effusion

A chylous effusion is suggested by the white milky fluid aspirated on thoracocentesis. Cardiothoracic and head or neck surgery, including excision of cervical lymph nodes, can result in damage to the thoracic duct resulting in a chylothorax. To confirm the diagnosis, test for pleural fluid triglycerides and cholesterol levels.

A Empyema

Aspiration of frankly purulent fluid would indicate an empyema.

B Mesothelioma

Though mesothelioma can present with a pleural effusion alone, the history in this case is not suggestive of this diagnosis.

D Reactive effusion

You would not aspirate white, milky fluid in a reactive effusion. The fluid may be cloudy or even slightly clear. Reactive effusions are principally lymphocytic infiltrates secondary to the movement of interstitial fluid. The underlying aetiology is varied, and includes congestive heart failure and collagen vascular disease.

E Malignant effusion

Malignant effusions can present as either serous or serosanginous fluid.

7179

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 35-year-old man presents with night sweats and a chronic cough. After investigations, a diagnosis of pulmonary tuberculosis is made, and he is commenced on quadruple therapy. On completion of six months of treatment, he attends for a follow-up computed tomography (CT) scan. He is noted to be asymptomatic, and there have been no identified complications of treatment so far. The CT result is shown below.



Which of the following is the most likely diagnosis?

Your answer was incorrect

A	Aspergilloma
B	Pulmonary nodule
C	Invasive aspergillosis
D	Recurrence of tuberculosis
E	*Staphylococcus aureus* empyema

Explanation

A Aspergilloma

This is the correct response, as it correlates with the asymptomatic presentation of the patient, the past history of tuberculosis and the CT appearances. Aspergilloma is a ball of fungal hyphae within a previous cavity space. The most common prior pathology creating the cavity is pulmonary tuberculosis. Patients are commonly asymptomatic, although occasionally invasion of the bronchial artery can cause haemoptysis.

C Invasive aspergillosis

This is an extremely rare condition in an immunocompetent patient. Symptoms would depend on the locus of infection but may include cough, breathlessness and haemoptysis, none of which are given in the question. A characteristic CT appearance may include the ‘halo sign’ – a haziness surrounding a nodule or infiltrate.

B Pulmonary nodule

While pulmonary nodules are asymptomatic, they do not cause a crescent-shaped air space, as shown in the image. The CT image is classical for an aspergilloma.

D Recurrence of tuberculosis

This response is incorrect, as one would expect the patient to present with symptoms congruent with the CT presentation. These may include malaise, anorexia, breathlessness and haemoptysis.

E *Staphylococcus aureus* empyema

One would expect such a large empyema to present with a symptomatic patient. *S. aureus* is a common organism for empyema development, and the previous tuberculosis infection would provide a lung cavity. An empyema would typically present with a fluid density in the lung field, sometimes with locules of gas.

71792

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A man with a history of COPD who is on home oxygen therapy is admitted to hospital with central chest pain. He suffers a cardiac arrest in the Emergency Department and is resuscitated for 23 min before return of spontaneous circulation. During resuscitation he was intubated, and he is now able to achieve adequate minute ventilation with minimal pressure support. His BP is 96/50 mmHg, HR 80, RR16, SpO_2 98% with an F_iO_2 of 0.6. Arterial blood gases are taken 3 min after ROSC.

Which of the following results are most likely to be shown on arterial blood gas analysis?

Your answer was correct

A pH 7.45 P_aO_2 7.9 kPa P_aCO_2 3.1 HCO_3^- 31 mmol/l

B pH 7.30 P_aO_2 7.9 kPa P_aCO_2 4.2 HCO_3^- 12 mmol/l

C pH 7.15 P_aO_2 12.2 kPa P_aCO_2 4.3 HCO_3^- 17 mmol/l

D pH 7.55 P_aO_2 12.2 kPa P_aCO_2 4.3 HCO_3^- 31 mmol/l

E pH 7.10 P_aO_2 7.9 kPa P_aCO_2 4.3 HCO_3^- 12 mmol/l

Explanation

C pH 7.15 P_aO_2 12.2 kPa P_aCO_2 4.3 HCO_3^- 17 mmol/l

This man has COPD requiring home oxygen. As such, he is likely to have chronic CO_2 retention and will consequently have significant renal compensation and an elevated serum bicarbonate. Given that he has suffered a significant period of cardiac arrest, it is also likely that he will have significant tissue hypoxia, leading to a build-up of products of metabolism such as lactate. This leads to the decreased pH. In patients who have suffered a cardiac arrest there is usually a significant metabolic acidosis with a low bicarbonate level. However, due to pre-existing renal compensation, the bicarbonate in this patient will not be as low as if there was no previous renal compensation. In the absence of previous renal compensation, a bicarbonate of the levels seen in B) or E), would be expected. The pressure support on the ventilator allows him to achieve adequate gas exchange.

A pH 7.45 P_aO_2 7.9 kPa P_aCO_2 3.1 HCO_3^- 31 mmol/l

In post-arrest patients we expect to see a metabolic acidosis.

B pH 7.30 P_aO_2 7.9 kPa P_aCO_2 4.2 HCO_3^- 12 mmol/l

The degree of acidaemia in this patient is likely to be worse, but the bicarbonate is likely to be higher due to previous renal compensation.

D pH 7.55 P_aO_2 12.2 kPa P_aCO_2 4.3 HCO_3^- 31 mmol/l

A metabolic alkalosis is not realistic in this scenario.

E pH 7.10 P_aO_2 7.9 kPa P_aCO_2 4.3 HCO_3^- 12 mmol/l

The bicarbonate is likely to be higher due to previous renal compensation.

21544

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A known asthmatic is admitted to A&E with wheeze and shortness of breath, although she is not yet showing signs of fatigue. She is given continued nebulised salbutamol, IV steroids and high-flow oxygen. You are asked by the nursing sister in charge whether giving IV magnesium is appropriate.

According to the BTS guidelines, which of the following would count against the need for IV magnesium at this stage?

Your answer was incorrect

A Patient is 18 years old

B Magnesium is at the lower end of the normal range

C Tachycardia with a heart rate of 120 beats/min

D Patient is treated with oral theophylline

E PEFR 240 vs 140 on admission (normal 500)

Explanation

E PEFR 240 vs 140 on admission (normal 500)

The British Thoracic Society (BTS) guidelines say that 50% magnesium sulphate at 1.2–2 g (4–8 mmol) over 20 min can be given IV to a patient with severe asthma not responding to inhaled bronchodilator or who has severe life-threatening asthma. In this case the PEFR is improving, without signs of fatigue, and it is therefore appropriate to continue with current management.

B Magnesium is at the lower end of the normal range

Magnesium concentration is irrelevant as supra-therapeutic levels are given to achieve a pharmacological response with reduced acetylcholine release and improved smooth muscle relaxation

A Patient is 18 years old

Age is irrelevant when considering the need for magnesium infusion, and there is little difference in response as age increases.

C Tachycardia with a heart rate of 120 beats/min

Heart rate >110/min is an indicator of severe asthma and more likely to drive the use of magnesium.

D Patient is treated with oral theophylline

The need for oral theophylline implies a poor response to beta-agonist therapy, and therefore a possible greater need for acute intervention with magnesium.

2332

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 45-year-old computer software firm owner presents with increasing breathlessness over the past few months. He has previously been fit and well and had installed a home gym, swimming pool and hot tub/steam room to use for personal training around 18 months ago. He is a non-smoker and has a previous history of asthma as a child. Examination reveals crackles and high-pitched wheeze throughout the lung fields. His blood pressure is 115/75 mmHg, with pulse 70 and regular. There is no ankle swelling.

Which of the following is the most appropriate intervention in this case?

Your answer was correct

A Inhaled corticosteroids

B Oral corticosteroids

C Oral antifungals

D Repainting and ventilation in the steam room

E Oral antibiotics

Explanation

D Repainting and ventilation in the steam room

Treatment of hypersensitivity pneumonitis (the diagnosis in this case) centres on allergen avoidance. *Cladosporum* is found in association with ceiling mould and is a source of ‘hot-tub lung’; as such, it is the most likely cause of his symptoms. Other moulds known to lead to hypersensitivity include micropolyspora, *Aspergillus* and thermoactinomycetes, which are associated with mouldy hay, mushroom compost and contaminated water in humidifiers. Trichosporum is associated with house dust. In this case the likely source is the steam room, and hence repainting and adequate ventilation should prevent further exposure.

A Inhaled corticosteroids

Inhaled corticosteroids are ineffective in the treatment of hypersensitivity pneumonitis; removal of exposure to the precipitant allergen is the most important intervention.

B Oral corticosteroids

Oral corticosteroids may be given for an initial period to kick-start the recovery in symptoms. However, if strict allergen avoidance is not practised, his symptoms will not completely resolve.

C Oral antifungals

Hypersensitivity pneumonitis is related to exposure to antigen rather than invasive fungal disease, and hence removal of exposure is the most important intervention.

E Oral antibiotics

Oral antibiotics have no role in the treatment of hypersensitivity pneumonitis.

32409

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Hypersensitivity Pneumonitis

Expanded
explanations

A 30-year-old man from London, with no significant past medical history, went on holiday to Peru to climb the Inca Trail. He is a non-smoker. After climbing above 3500 m he developed a headache, associated with nausea and vomiting. He also felt tired and complained of insomnia. His respiratory rate was 22/min and his pulse was 90/min. The rest of the physical examination was unremarkable. No investigations are available as the nearest hospital is over 100 km away. That night it was noticed that he was disorientated and walking with a broad-based gait, and by the following morning he was drowsy and incoherent.

How would you treat this man?

Your answer was incorrect

A	Immediate transfer down the mountain
B	Acetazolamide and continue ascent
C	Aspirin
D	Dexamethasone and continue ascent
E	Immediate antibiotics

Explanation

A Immediate transfer down the mountain

There are three types of altitude illness: acute mountain sickness (AMS), high-altitude cerebral oedema (HACE) and high-altitude pulmonary oedema (HAPE). This man has acute mountain sickness with some evidence of cerebral oedema. The exact cause is unknown, but it is thought to be secondary to cerebral oedema and raised intracranial pressure. It is more common in people who live at low altitude.

Symptoms and signs of AMS are non-specific; however, diagnosis can be made using the criteria named in 'The Lake Louise Score', namely, the setting of recent gain in altitude, the presence of headache and at least one of the following symptoms:

- Gastrointestinal – anorexia, nausea and vomiting
- Fatigue and weakness
- Dizziness or light-headedness
- Difficulty sleeping.

If he does not improve and progresses to develop ataxia and drowsiness, this is the development of high-altitude cerebral oedema, which can be considered as end-stage AMS. This is the presentation of this case.

B Acetazolamide and continue ascent

Acetazolamide may be used for the prevention and treatment of AMS. This is a carbonic anhydrase inhibitor which causes intracellular acidosis. It has a diuretic effect which can be useful in AMS/HACE and HAPE. It is also beneficial as it shifts the oxygen-dissociation curve to the right and therefore, at lower partial pressures, more oxygen is released to the tissues. In this scenario, this patient has full house symptoms of altitude sickness and is at high risk of further deterioration. Further ascent, even with acetazolamide, would be dangerous.

C Aspirin

This would not help symptoms of mountain sickness. This is a treatment for acute MI or non-haemorrhagic CVA. Aspirin, acetylsalicylic acid, may also be used in the treatment of pain, fever and inflammation.

D Dexamethasone and continue ascent

This patient has full house symptoms of altitude sickness and is at high risk of further deterioration. Further ascent even with acetazolamide would be dangerous.

E Immediate antibiotics

This is an appropriate treatment for severe infection. However, it has no role in AMS. There is no indication that this patient is suffering from infection: there are no fever, chills, rigors or skin rash.

70109

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

An HIV-positive patient is admitted acutely unwell with shortness of breath initially only on exertion, but now present even at rest. He has been prescribed HAART and pentamidine nebulisers (co-trimoxazole being unsuitable for him). However, recently he says that he has not had time to come to the clinic for treatment. His chest X-ray shows bilateral air-space shadowing and his arterial blood gas on air shows pO_2 7.9 kPa and pCO_2 5.0 kPa. His CD4 count is 164.

What is the most appropriate initial treatment option?

Your answer was incorrect

A Clindamycin and primaquine with prednisolone

B Further nebulised pentamidine

C Intravenous Pentamidine with prednisolone

D Intravenous cefotaxime and oral clarithromycin

E IV hydrocortisone

Explanation

A Clindamycin and primaquine with prednisolone

This combination is recommended for pneumocystis, where trial evidence now shows higher response rates than for IV pentamidine, (the main alternative in this case). He should also be treated with oral or intravenous steroids as his pO_2 is below the 9.3 kPa threshold.

C Intravenous Pentamidine with prednisolone

The patient has an intolerance to co-trimoxazole and has a CD4 count less than 200, so needs pentamidine nebulisers as PCP prophylaxis. If he has not been having these, he is at risk of pneumocystis pneumonia (PCP), which must be treated even before definitive histology can be obtained. The ideal treatment for him is as for severe PCP which drives clindamycin and primaquine as the correct answer because response rates are higher than for pentamidine. IV pentamidine is a fall-back option in the event that primaquine and clindamycin isn’t tolerated. He needs treatment with oral or intravenous steroids as his pO_2 is below the 9.3 kPa threshold.

B Further nebulised pentamidine

This is an appropriate prophylactic treatment, but is not the best management choice for this patient’s acute presentation.

D Intravenous cefotaxime and oral clarithromycin

This antibiotic combination therapy would be suitable if a bacterial aetiology were suspected for his presentation, and may well be appropriate as additional therapy until microbiological confirmation of the organism can be obtained, but the more likely diagnosis of PCP would make this inappropriate as sole therapy.

E IV hydrocortisone

While corticosteroids are an important adjuvant therapy, they are inadequate as monotherapy in the treatment of PCP.

71911

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 58-year-old female presents with a history of fever of one week’s duration associated with rhinorrhoea, headache and cough productive of yellowish sputum. She has had type 2 diabetes for 8 years, complicated by painful peripheral neuropathy. Her present medications include gliclazide 40 mg twice daily, metformin 500 mg three times a day and amitriptyline 75 mg at bedtime. Physical examination reveals temperature of 38°C, respiratory rate 18/min, blood pressure 110/70 mmHg, mild posterior pharynx erythema, tenderness over the maxillary sinus and fine inspiratory crackles at the right lung base.

Investigations reveal the following:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	110 g/l	115–155 g/l
White cell count (WCC)	9.0 × 10^9/l	4.0–11.0 × 10^9/l
Reticulocytes	6%	
Sodium (Na^+)	132 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Chloride (Cl^-)	98 mmol/l	98-106 mmol/l
Bicarbonate (HCO_3^-)	22 mmol/l	24–30 mmol/l
Bilirubin	40 µmol/l	2–17 µmol/l
Chest X-Ray (CXR)	Infiltrates at R. lower zone	

Which of the following drugs should be used to treat this patient?

Your answer was incorrect

A Erythromycin

B Levofloxacin

C Co-amoxiclav

D Doxycycline

E Rifampicin

Explanation

D Doxycycline

This patient has pneumonia and evidence of haemolysis (reticulocytosis and hyperbilirubinaemia). This suggests *Mycoplasma pneumoniae* as a possible aetiological agent. Macrolides, and fluoroquinolones such as levofloxacin and clarithromycin, can all be used to treat *Mycoplasma*. However, this patient is taking amitriptyline that can cause prolongation of the QT interval. Prescribing another agent that can also prolong the QT interval, such as macrolides or levofloxacin. can be hazardous. Hence, doxycycline is the best therapy for this patient. Other antimicrobial agents that can cause prolongation of the QT interval include:

- Pentamidine
- Spiramycin
- Chloroquine
- Halofantrine
- Mefloquine.

Management of drug-induced long-QT syndrome is to stop the precipitating drug, correction of any electrolyte disturbance such as hypokalaemia or hypomagnesaemia, and treatment of associated ventricular arrhythmia. First-line pharmacological therapy for drug-induced long-QT syndrome is intravenous magnesium sulphate 2 g administered as a bolus over 1–2 min, followed by another bolus in 15 min if required, or continuous infusion at a rate of 5–20 mg/min.

B Levofloxacin

Levofloxacin has been linked to QT prolongation and torsades, and should thus be avoided in the context of concomitant amitriptyline administration. Levofloxacin is active against aerobic Gram-positive bacteria, aerobic Gram-negative bacteria, anaerobic bacteria, *Chlamydophila pneumoniae*, *Chlamydophila psittaci*, *Chlamydia trachomatis*, *Legionella pneumophila*, *Mycoplasma pneumoniae*, *Mycoplasma hominis and Ureaplasma urealyticum.*

A 2019 UK government drug safety update advises that fluoroquinolones should now be avoided in non-severe or self-limiting infections and in mild to moderate infections unless other commonly recommended antibiotics are not suitable. This is due to rare reports of long lasting or potentially irreversible side effects such as tendinitis and peripheral neuropathy. Fluoroquinolones should be used with special caution in patients older than 60 years or those with renal impairment due to higher risk of these significant adverse affects.

A Erythromycin

As a macrolide, erythromycin has been linked to QT prolongation and torsades, and should thus be avoided in the context of concomitant amitriptyline administration. Erythromycin is indicated for susceptible infections in patients with penicillin hypersensitivity (e.g. respiratory tract infections (including *Legionella* infection), skin and oral infections, and *Campylobacter* enteritis).

C Co-amoxiclav

This would not represent appropriate cover against *Mycoplasma pneumoniae*, the likely pathogen in this case. Clavulanic acid is a potent inhibitor of β-lactamases produced by Gram-negative bacteria, including *Haemophilus ducreyi*, H. influenzae, *Neisseria gonorrhoeae*, *Moraxella catarrhalis*, *Bacteroides fragilis* and some

Enterobacteriaceae. It also inhibits β-lactamases produced by *Staphylococcus aureus*. Clavulanic acid reverses resistance to amoxicillin in beta-lactamase-producing strains of species otherwise sensitive, and also augments the activity of amoxicillin against numerous species not normally considered sensitive.

E Rifampicin

This would not represent appropriate cover against *Mycoplasma pneumoniae*, the likely pathogen in this case. Rifampicin, in combination with other antibacterials, is indicated for brucellosis, legionnaires' and serious Staph. infections. It's also indicated for the treatment of tuberculosis in combination with other antibacterials, and for the prevention of secondary cases of meningococcal meningitis and *Haemophilus influenzae* type B.

7435

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

National Information Program on Antibiotics
antibiotics-info.org/

(http://www.antibiotics-info.org/)

A 62-year-old retired office worker presents to the clinic with progressively increased shortness of breath on exercise and a chronic non-productive cough.

On examination, he looks tired and a little thin; his body mass index (BMI) is 21. His blood pressure (BP) is 142/88 mmHg, with pulse 78/min and regular. There are bi-basal end-inspiratory crackles on auscultation. He is clubbed and there is bilateral peripheral ankle oedema.

Investigations:

Investigation	Result	Normal value
Haemoglobin	148 g/l	135–175 g/l
White cell count (WCC)	7.9 × 10^9/l	4–11 × 10^9/l
Platelets	201 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Creatinine	141 µmol/l	50–120 µmol/l
Autoimmune profile	No specific abnormalities	
PaO_2	7.9 kPa	10.5–13.5 kPa
$PaCO_2$	3.9 kPa	4.6–6.0 kPa
CXR	Interstitial shadowing at both lung bases	
CTR	Suggests lung fibrosis	

Which of the following is the most appropriate initial treatment?

Your answer was incorrect

A N-acetylcysteine

B High-dose prednisolone

C Low-dose prednisolone

D Cyclophosphamide

E Long-term oxygen replacement therapy

Explanation

E Long-term oxygen replacement therapy

This man has strong evidence to support a diagnosis of idiopathic pulmonary fibrosis. However, there is nothing in either his occupational history or the blood results to suggest a diagnosis of connective tissue disease or occupational lung disease.

Unfortunately, trials of corticosteroids, potential disease-modifying agents such as cyclophosphamide or methotrexate and novel therapies such as *N*-acetylcysteine have been uniformly disappointing. However, these agents are often trialled, as only a small proportion of patients may respond.

He is significantly hypoxic, and thus home oxygen therapy may offer both symptom relief and affect prognosis with respect to secondary pulmonary hypertension and right heart failure. Pirfenidone and nintedanib are currently available agents which are proven to delay the progression of the disease.

As per BTS guidelines, LTOT should be ordered for patients with ILD with a resting PaO2 ≤7.3 kPa. (Grade D). LTOT should be ordered for patients with ILD with a resting PaO2 ≤8kPa in the presence of peripheral oedema, polycythaemia (haematocrit ≥55%) or evidence of pulmonary hypertension.

C Low-dose prednisolone

Prednisolone is specifically advised against in idiopathic pulmonary fibrosis.

A N-acetylcysteine

N-acetylcysteine can be used under limited circumstances for idiopathic pulmonary fibrosis, but NICE advises that its benefits are uncertain.

B High-dose prednisolone

Prednisolone is specifically advised against in idiopathic pulmonary fibrosis. However, if extrinsic allergic alveolitis (EAA) was suspected then it could be trialled. EAA would be suggested by honeycombing on CT scanning. In the absence of this feature and of any known environmental exposure, this would not be helpful.

D Cyclophosphamide

Cyclophosphamide is specifically advised against in idiopathic pulmonary fibrosis. Had the diagnosis of a rheumatoid lung condition been suspected, then this could have been considered.

21246

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

NICE. Idiopathic pulmonary fibrosis
nice.org.uk/guidance/cg163
(https://www.nice.org.uk/guidance/cg163)

A 64-year-old homeless man is admitted on the routine medical take. He has lost a significant amount of weight over the past few months and now presents with increasing shortness of breath. He is a former heavy smoker. On examination: decreased air entry over the right lower chest and dullness to percussion. Chest X-ray reveals a large, right-sided pleural effusion. Pleural aspiration is performed.

Which one of the following features, if present, would be the most specific indicator of probable underlying malignancy?

Your answer was correct

A Turbid fluid

B High pleural fluid neutrophil count

C Uniform blood staining throughout pleural fluid samples

D Abnormal liver function tests

E Abnormal pleural fluid LDH levels

Explanation

C Uniform blood staining throughout pleural fluid samples

Uniform blood staining and a protein level of >30 g/l are supportive of a diagnosis of malignancy.

pH can also be useful for interpreting pleural fluid results. Low pleural fluid pH is associated with a number of conditions, including pleural infection and empyema, inflammatory effusions such as those related to rheumatoid arthritis, tuberculosis, malignancy and oesophageal rupture.

A Turbid fluid

Turbid fluid is suggestive of an infectious cause.

B High pleural fluid neutrophil count

This is also indicative of an infective cause.

D Abnormal liver function tests

Although deranged LFTs may suggest malignancy, there are other causes such as infection (particularly atypical organisms such as *Legionella*) and chronic disease such as liver cirrhosis.

E Abnormal pleural fluid LDH levels

LDH is used to differentiate an exudate from a transudate. Raised LDH is suggestive of an exudate and therefore infectious or inflammatory cause. It needs to be measured simultaneously with serum LDH as it is the ratio of pleural fluid to serum LDH that is interpreted. A ratio >0.6 indicates an exudate.

32374

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 48-year-old builder working in the UK for over 5 years presents to your clinic with a 1-year history of wheeze and cough that is worse in the early hours of the morning. On direct questioning he admits to mild haemoptysis over the past few days with no increase or change in his sputum production, occasional mild night sweats and he has noticed a slight weight loss over the past few months due to decreased appetite. His past medical history includes previous alcohol excess and type 2 diabetes. A chest X-ray taken two years ago showed clear lung fields and no signs of previous lung pathology.

Which of the following pathologies is the most likely cause of his haemoptysis?

Your answer was correct

A Aspergilloma

B Allergic bronchopulmonary aspergillosis (ABPA)

C Semi-invasive aspergillosis

D Invasive aspergillosis

E Asthma exacerbation

Explanation

C Semi-invasive aspergillosis

This is a chronic indolent disease that affects people with mild immunosuppression, such as those with diabetes or chronic obstructive pulmonary disease or those who abuse alcohol, are elderly or on prolonged steroid use. If left untreated there is a significant associated mortality; however, it is difficult to diagnose due to its slow progression both radiologically and clinically. Symptoms include cough, wheeze, bronchitis and constitutional symptoms, and about 15% present with haemoptysis.

A Aspergilloma

This is a ball of fungus within a pre-existing cavity such as that caused by tuberculosis (TB). With a recent clear chest X-ray (2 years ago), this patient is unlikely to have any cavities in which an aspergilloma could reside. Furthermore, aspergillomas are usually asymptomatic but can cause weight loss, cough and haemoptysis, which can be massive and require emergency surgical excision. Oral antifungals can be given, but with limited success.

B Allergic bronchopulmonary aspergillosis (ABPA)

This is an IgE and IgG reaction to *Aspergillus* sp. in asthmatics. The key criteria for diagnosis include a long history of asthma, skin-prick/IgE, positive/IgG precipitins to *Aspergillus* fumigates, central bronchiectasis, elevated total serum IgE, blood/sputum eosinophilia and pulmonary infiltrates.

D Invasive aspergillosis

This patient is not acutely unwell. This condition occurs in immunosuppressed individuals and is caused by haematogenous spread of the fungus, so prompt diagnosis and treatment are required to prevent a high mortality rate.

E Asthma exacerbation

Night sweats and haemoptysis are not features of asthma exacerbation.

32444

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded
explanations

An 18-year-old woman presents with increasing shortness of breath and nasal congestion. She has a past history of recurrent chest infections.

On examination, her fingers appear as shown below. Her sputum is mucopurulent.



Her chest X-ray is shown below:



Pleural Fluid	
Appearance	Clear
Protein	35 g/l
WCC	2440 (80% lymphocytes)

What is the most likely diagnosis?

Your answer was correct

A Cystic fibrosis

B Yellow-nail syndrome

C Filariasis

D Meigs' syndrome

E Tuberculosis

Explanation

B Yellow-nail syndrome

This girl has yellow-nail syndrome which is a rare condition consisting of the triad of primary lymphoedema, recurrent pleural effusions and dystrophic yellow nails. It is associated with bronchiectasis and sinusitis. Pleural fluid is typically a clear exudate in which lymphocytes predominate. The underlying abnormality is hypoplasia of the lymphatics with impaired drainage. This results in subungual oedema, lymphoedema and pleural effusions.

A Cystic fibrosis

Cystic fibrosis would be expected to present with respiratory symptoms from infancy, is not usually associated with pleural effusions, and would be expected to cause finger-clubbing, not dystrophic nails.

C Filariasis

Filariasis is a cause of lymphoedema (most commonly of the legs and external genitals), secondary to microfilariae being introduced into the skin by mosquitoes and migrating to the lymphatics. It may present asymptomatically or with infective symptoms of fever/rigors/myalgia, and lymphangitis. It can be associated with chylous pleural effusions.

D Meigs' syndrome

This is a rare condition comprised of the triad of pleural effusions, ascites and benign ovarian tumours – usually fibromas.

E Tuberculosis

TB may present as a lymphocytic pleural effusion, but with the history of recurrent chest infections rather than an acute illness and no mention of night sweats, fevers or weight loss, alongside the examination finding of dystrophic nails, it is less likely than yellow nail syndrome.

70142

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pleural Effusion

Media

Expanded explanations

Flashcards

A 45 year old woman is referred by the GP to the respiratory care clinic with shortness of breath of unknown origin. Spirometry results: KCO 60%, FEV_1/FVC 90%.

Which of the following is most likely to be the underlying cause of these pulmonary function test results?

Your answer was incorrect

A Sarcoidosis

B Severe kyphoscoliosis

C Muscular dystrophy

D Pneumonectomy

E Bronchiectasis

Explanation

A Sarcoidosis

A FEV_1/FVC of 90% indicates a restrictive cause. All the above diseases are evidenced by a restrictive pattern, except bronchiectasis which can result in an obstructive pattern. Only sarcoidosis would reduce the transfer coefficient as there is damage to the alveolar cells themselves. Other conditions where a reduction in KCO is reduced include those such is idiopathic pulmonary fibrosis, where there is also loss of alveolar function.

D Pneumonectomy

Pneumonectomy is most likely to have been performed for chronic bronchiectasis or lung cancer, both of which would be unusual in a 45-year-old woman and wouldn't be expected to impact on transfer factor in the remaining lung.

B Severe kyphoscoliosis

Abnormalities of the thoracic cage lead to a reduction in available lung volume for gas exchange but don't impact on the transfer coefficient.

C Muscular dystrophy

Similar to kyphoscoliosis, muscular dystrophy can lead to a restrictive lung pattern but doesn’t impact on gas transfer.

E Bronchiectasis

Bronchiectasis produces an obstructive lung pattern on pulmonary function testing.

2327

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 30-year-old woman presents with sudden-onset shortness of breath and chest pain. She had a similar episode earlier in the year. At that time she had a chest X-ray taken and a needle inserted into her lung, which relieved the problem. Last year she had shortness of breath associated with ‘fluid on her lung’, which was drained off, and she was told to follow a low-fat diet. She has no other past medical history and is not on any medications.

What is the most likely diagnosis?

Your answer was incorrect

A Histiocytosis

B Tuberous sclerosis

C Neurofibromatosis type 1

D Recurrent *Pneumocystis jirovecii* infection

E Lymphangioleiomyomatosis

Explanation

E Lymphangioleiomyomatosis

Lymphangioleiomyomatosis is a rare idiopathic disease, affecting women of reproductive years. It is characterised by infiltration of immature muscle cells into the bronchiolar and alveolar walls. This results in destruction of the airways, cyst formation and progressive decline in lung function. Typical radiological findings are of interstitial lung disease, recurrent pneumothoraces and chylous effusions. Treatment is lung transplantation. Patients are asked to avoid taking the oral contraceptive pill, pregnancy and long-haul flights. Conservative management includes progesterone supplements, but the response is variable. Management of pneumothorax and chylous effusions is standard. If this occurs in men then tuberous sclerosis should be considered.

B Tuberous sclerosis

Tuberous sclerosis is an autosomal dominant disorder with hamartoma formation in the heart, kidneys, eyes, brain and skin. Patients can have adenoma sebaceum, angiofibromas and ash leaf hypomelanic macules, as well as shagreen patches. It is associated with Von Hippel–Lindau. The absence of any such skin features makes this an unlikely underlying cause.

A Histiocytosis

Histiocytosis, or Langerhans cell histiocytosis, is a sign or multi-system condition characterised by infiltrating granulomas with Langerhans cells. In the lungs this can cause pneumothoraces and pulmonary hypertension, with cysts and honeycombing on imaging. Diagnosis is by biopsy and treatment is with local excision and steroids. The presence of pleural effusions makes this unlikely in this patient, as does the absence of any guiding radiological features.

C Neurofibromatosis type 1

Neurofibromatosis type 1 is an autosomal dominant condition. Patients have café-au-lait spots, freckling and neurofibromas. Some can develop problems resulting from neurofibromas, particularly ophthalmic and neurological complications. The absence of any neurofibromas makes this an unlikely diagnosis.

D Recurrent *Pneumocystis jirovecii* infection

Pneumocystis jirovecii is an opportunistic, AIDS-defining infection associated with severe immunosuppression in HIV. It is very unlikely to cause infection without HIV and immunocompromise, but in such patients it can cause pneumothorax as a complication.

7165

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 67-year-old obese man with known mild chronic obstructive pulmonary disease (COPD) presents via ambulance. He is drowsy, and his wife is unable to rouse him. He usually has a good level of functioning, is able to walk to the local pub, uses a salbutamol inhaler only and continues to smoke five cigarettes per day.

You arrange a chest X-ray and blood gases on air, which reveal the following:

Investigation	**Result**	**Normal Value**
pH	7.2	7.35–7.45
pCO_2	8.7 kPa	4.6–6.0 kPa
pO_2	6.3 kPa	10.5–13.5 kPa
Bicarbonate (HCO_3^-)	27.7 mmol/l	24–30 mmol/l

Which of the following best describes the blood gas results?

Your answer was correct

A Mixed metabolic and respiratory acidosis

B Metabolic acidosis

C Fully compensated respiratory acidosis

D Respiratory acidosis

E Metabolic alkalosis

Explanation

D Respiratory acidosis

This is a respiratory acidosis, characterised by raised $p_a(CO_2)$, decreased pH and serum bicarbonate in the normal range. Exacerbation of COPD is a common cause, although the fact that the bicarbonate is in the normal range suggests that his deterioration has been too rapid to allow renal compensation. Pneumonia or abnormalities of the thoracic cage, such as those related to spina bifida, may also be responsible. In compliant patients, non-invasive positive pressure ventilation (NIPPV) may be an option. Alternatives include doxapram, a central respiratory stimulant. These would be combined with appropriate antimicrobial therapy. Unfortunately, many patients with COPD are unsuitable for formal intubation and ventilation.

A Mixed metabolic and respiratory acidosis

HCO3⁻ is a base in nature. This means that an acidosis is metabolic if HCO3- is lower than the normal range. This is not the case as it is increased rather than lowered, indicating a compensatory mechanism for respiratory acidosis.

B Metabolic acidosis

Metabolic acidosis can occur with an increased or normal anion gap. In an increased anion gap there is increased production or reduced excretion of organic acids, which causes HCO3⁻ to fall. In a normal anion gap there is loss of bicarbonate or ingestion of H^+ ions, such as in diarrhoea and renal tubular acidosis. In this patient, the HCO3⁻ is raised in acidosis, meaning that it is being produced in greater amounts as a compensatory mechanism.

C Fully compensated respiratory acidosis

The acidosis is not fully compensated as the pH is <7.35, and so the patient is still acidotic.

E Metabolic alkalosis

Metabolic alkalosis can occur with vomiting, potassium depletion and ingestion of a base. It is seen with a pH >7.45 and a high HCO3⁻. This patient has a low pH.

6539

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 16-year-old boy is referred to the Respiratory Clinic with a persistent cough that is productive of brown/green sputum. He occasionally produces a small amount of blood. His mother says he has always suffered from chest infections, requiring numerous courses of antibiotics. He also complains of offensive-smelling stools which are difficult to flush.

On examination, he looks small for his age. Auscultation of his chest reveals bibasal coarse crackles.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	112 g/l	135 - 175 g/l
White cell count (WCC)	10.7 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	511 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	91 fl	80 - 100 fl
Sodium (Na^+)	139 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5 - 5.0 mmol/l
Urea	5.3 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	99 μmol/l	50 - 120 μmol/l
Bilirubin	18 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	34 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	93 u/l	30 - 150 u/l
Albumin	3.9 g/dl	3.4 - 5.4 g/dl
Total Protein level	50 g/l	60 - 83 g/l

His CXR is shown below:



What is the most likely diagnosis?

Your answer was incorrect

A Cystic fibrosis

B Eosinophilic granulomatosis with polyangiitis (Churg-Strauss)

C Kartagener syndrome

D X-linked hypogammaglobulinaemia

E Young syndrome

Explanation

D X-linked hypogammaglobulinaemia

This patient has X-linked hypogammaglobulinaemia. The main clue is the low total protein level with a normal albumin level, demonstrating low levels of globulins. Patients usually present in childhood with recurrent infections, which result in bronchiectasis if untreated, and malabsorption, in a similar way to patients with cystic fibrosis. All immunoglobulin classes, as well as B and plasma cells, are reduced.

The defect is in the differentiation of pre-B cells into B cells. T cells are normal. The gene defect is on the long arm of the X chromosome, making it an X-linked condition. Treatment is with IV immunoglobulin therapy which, if started early, can prevent progression of the disease. The CXR shows bronchiectasis, parallel thickened bronchial walls with a ‘tram track’ appearance.

Both inflammatory and infectious enteritis are more common in patients with X-linked hypogammaglobulinaemia, which accounts for the appearance of the stools seen here.

C Kartagener syndrome

Kartagener syndrome is associated with bronchiectasis, sinusitis and dextrocardia. There is no dextrocardia seen on the CXR, making this less likely.

A Cystic fibrosis

Although the patient has symptoms compatible with cystic fibrosis, bronchiectasis that is visible on his CXR and the low total protein level make X-linked hypogammaglobulinaemia more likely.

B Eosinophilic granulomatosis with polyangiitis (Churg-Strauss)

An asthma/rhinitis prodrome to this condition is classical, followed by eosinophilia and systemic vasculitis. Bronchiectasis seen on the CXR and infective, rather than asthmatic, chest symptoms make hypogammaglobulinaemia a more likely diagnosis.

E Young syndrome

Young syndrome is associated with the triad of bronchiectasis, rhinosinusitis and reduced fertility, rather than with malabsorption.

73462

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Bronchiectasis

Media

Expanded explanations

Flashcards

A 68-year-old woman presents to the clinic complaining of increasing shortness of breath over the past year. Her GP has prescribed steroid and salbutamol inhalers, but they have given her little benefit. She has a past history of hypertension and has recently been prescribed furosemide for swollen ankles. In the past week or so her breathing has begun to deteriorate more rapidly. Examination reveals an increased respiratory rate at rest and finger clubbing. Auscultation of the chest reveals end inspiratory crackles at both lung bases.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	131 g/l	115–155 g/l
White cell count (WCC)	6.9 × 10^9/l	4.0–11.0 x 10^9/l
Platelets (PLT)	300 × 10^9/l	150–400 x 10^9/l
Sodium (Na^+)	140 mmol/l	135–145 mmol/l
Potassium (K^+)	5.0 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	120 μmol/l	50–120 μmol/l
pH	7.4	7.35–7.45
pO_2	7.2 kPa	10.5–13.5 kPa
pCO_2	3.7 kPa	4.6–6.0 kPa
Computed tomography (CT) thorax	Interstitial lung disease, evidence of bilateral consolidation	

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Sarcoidosis

B Idiopathic pulmonary fibrosis

C Silicosis

D Alveolar cell carcinoma

E Mycoplasma pneumonia

Explanation

B Idiopathic pulmonary fibrosis

The clinical picture here of gradual deterioration, coupled with crackles on auscultation consistent with fibrosis, finger clubbing and the CT findings, fits with the diagnosis. Patients may suffer an acute deterioration which is accompanied by consolidation, as we see here. Pulmonary histology may reveal areas of normal lung, interstitial fibrosis, inflammation and honeycomb change. Perfinidone and nintedanib are both potential interventions for idiopathic pulmonary fibrosis; they are not curative but are able to slow the decline of lung function.

C Silicosis

Silicosis occurs in patients chronically exposed to silica dust, such as stonemasons or miners. It is associated with egg cell calcification on the chest x-ray, rather than the pattern of respiratory changes seen here.

A Sarcoidosis

Sarcoidosis presents with bilateral hilar lymphadenopathy and other extrapulmonary features such as erythema nodosum.

D Alveolar cell carcinoma

Alveolar cell carcinoma presents as a peripheral lesion, the cancer arising from epithelial cells within the alveoli themselves.

E Mycoplasma pneumonia

Mycoplasma pneumonia is associated with respiratory symptoms over a few days/weeks and is characterised by wheeze, dry cough and chest pain. The time course in this case is too long for mycoplasma pneumonia to be the underlying cause for the patient’s symptoms.

18629

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

You are seeing a patient in clinic. Unfortunately, all clinic notes in the Cardiorespiratory Test Department, are being audited; however, the patient brings along a copy of her lung function results. Her date of birth is missing from the report.

Lung function results reveal the following:

Investigation	Result	Normal values
FEV_1	0.30	1.2–2.9
Total lung capacity	6.8	4.4–6.8
Corrected transfer factor	3.4	4.2–8.8

From these results, which disease do you suspect that she is most likely to have?

Your answer was incorrect

A Very severe chronic obstructive pulmonary disease (COPD)

B Severe asthma

C Moderate COPD

D Non-specific interstitial pneumonitis (NSIP)

E Sarcoidosis

Explanation

A Very severe chronic obstructive pulmonary disease (COPD)

COPD is defined as a reduced FEV_1/FVC ratio (where FEV_1 is forced expired volume in 1 second and FVC is forced vital capacity), such that FEV_1/FVC is less than 0.7. Though you are not given an FVC reading, note that the total lung capacity here is within the normal range, yet the FEV_1 is significantly reduced, this fits best with a diagnosis of COPD.

B Severe asthma

Asthma is associated with a normal or raised transfer factor.

C Moderate COPD

Predicted FEV_1 in moderate COPD is 50–79%. See table below for classification of COPD.

FEV_1 % predicted	**Severity of airflow obstruction**
> 80%	Mild**
50–79%	Moderate
30–49%	Severe
< 30%	Very severe

**If FEV_1 is ≥ 80% predicted normal, a diagnosis of COPD should only be made in the presence of respiratory symptoms, for example breathlessness or cough.

D Non-specific interstitial pneumonitis (NSIP)

Restrictive lung disease is associated with a decreased total lung capacity.

E Sarcoidosis

Like NSIP, sarcoidosis falls under the category of restrictive lung disease, and is associated with a decreased total lung capacity.

7176

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

COPD Diagnosis & Management
goldcopd.org/wp-content/uploads/2016/12/wms-GOLD-2017-Pocket-Guide.pdf
(http://goldcopd.org/wp-content/uploads/2016/12/wms-GOLD-2017-Pocket-Guide.pdf)

An overweight 55-year-old man complains of daytime sleepiness, so much so that he is at risk of losing his job. He also notes morning headaches, and his wife complains that his snoring is getting worse.

He is a heavy smoker and drinks about ten pints per night. His exercise tolerance is only about 500 m before he feels too short of breath to continue.

On examination, his body mass index (BMI) is 38 kg/m^2.

His arterial blood gas at 10 pm reveals the following:

Investigation	Result	Normal Value
PaO_2	9.3 kPa	10.5–13.5 kPa
$PaCO_2$	8.2 kPa	4.6–6.0 kPa

A repeat arterial blood gas the following morning reads:

Investigation	Result	Normal values
PaO_2	9.8 kPa	10.5–13.5 kPa
$PaCO_2$	8.4 kPa	4.6–6.0 kPa

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Obesity hypoventilation syndrome

B Central sleep apnoea

C Chronic obstructive pulmonary disease (COPD)

D Chronic thromboembolic pulmonary hypertension (CTPH)

E Motor neurone disease (MND)

Explanation

A Obesity hypoventilation syndrome

Obesity hypoventilation syndrome is the presence of awake alveolar hypoventilation in an obese individual. Patients present with daytime somnolence, loud snoring, fatigue and impaired cognition. Management is with weight loss and non-invasive ventilation at night – this can be continuous or bi-level, depending on the presence of co-existing obstructive sleep apnoea. Where peripheral sleep apnoea is the cause, as here, there are usually pointers to the diagnosis such as obesity.

B Central sleep apnoea

Central sleep apnoea is the absence of both airflow and ventilatory effort during sleep. Diagnosis is by polysomnographic sleep studies. Treatment is with CPAP. A rise in carbon dioxide in the morning is most likely to be seen in apnoea due to a peripheral cause. Where central sleep apnoea is the cause there may be pointers to an underlying neurological condition.

C Chronic obstructive pulmonary disease (COPD)

COPD is a disorder of progressive airway obstruction and can cause increases in COPD chronically, and acutely in exacerbations. Patients with COPD can retain slightly more carbon dioxide in the morning, but this would not cause such a profound rise, nor the symptoms of sleep apnoea.

D Chronic thromboembolic pulmonary hypertension (CTPH)

Following a PE most people experience complete resolution, but in some cases the embolus remains and results in pulmonary hypertension. CTPH presents initially with exertional breathlessness. Management may require surgical approaches to remove the clot. The absence of both exertional shortness of breath and previous PE makes this an unlikely diagnosis.

E Motor neurone disease (MND)

If the patient had MND such that it was affecting his diaphragm overnight, causing carbon dioxide retention, you would expect the disease to have presented itself neurologically as well, but the patient is able to walk.

7177

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 65-year-old man presents to the Emergency Department with acute breathlessness, cough and frothy, blood-stained sputum. He had been trying to save his cat from his house, which was on fire when he was overcome with smoke. The firefighters rescued him and the cat.

On examination, he is dyspnoeic, with a blood pressure of 111/62 mmHg, a pulse rate of 104 bpm and a respiratory rate of 28 breaths per minute. On auscultation, he has crackles in both lung fields. The abdominal examination is unremarkable. His oxygen saturation is 88% on air. You immediately start the patient on high-flow oxygen (15 l).

The following results are obtained on high flow oxygen:

Investigation	**Result**	**Normal value**
pH	7.31	7.35–7.45
pa(O_2)	8.0 kPa	10.5–13.5 kPa
pa(CO_2)	4.2 kPa	4.6–6.0 kPa
Bicarbonate	21 mmol/l	24–30 mmol/l

A radiological diagnosis is given in the picture (see below).



Which of the following options is the next management of choice?

Your answer was incorrect

A Give sublingual glyceryl trinitrate (GTN) spray (2 puffs)

B Intubate and ventilate the patient

C Continuous pulmonary airway pressure (CPAP)

D Bi-level pulmonary airway pressure (BiPAP)

E Ventilation/perfusion scan

Explanation

B Intubate and ventilate the patient

This patient has had an acute insult to his lungs, following smoke and carbon monoxide exposure, with his chest X-ray appearances consistent with acute respiratory distress syndrome (ARDS). Given that he is failing to maintain a $p_a(O_2) \leq 8.3$ kPa, despite maximal oxygen delivery, the next step would be intubation and ventilation. Invasive circulatory monitoring is also essential with respect to managing pulmonary capillary wedge pressure and cardiac output. There is conflicting evidence for the use of steroids; as such, they are not universally recommended in ARDS.

Image source- learningradiology.com/notes/chestnotes/ards.htm

C Continuous pulmonary airway pressure (CPAP)

This patient has had an acute insult to his lungs, following smoke and carbon monoxide exposure, with his chest X-ray appearances consistent with ARDS. CPAP would be a reasonable option in pulmonary oedema, but given that this patient most likely has ARDS and is failing to maintain his $p_a(O_2) \geq 8.3$ kPa on maximal oxygen delivery, invasive ventilation is the more appropriate option for ventilatory support.

A Give sublingual glyceryl trinitrate (GTN) spray (2 puffs)

The clinical scenario above is describing a patient who has had an acute insult to his lungs, following exposure to smoke and carbon monoxide. He has no features whatsoever of angina or acute coronary syndrome, so sublingual GTN would be inappropriate.

D Bi-level pulmonary airway pressure (BiPAP)

While the patient has a reduced pH, this is clearly metabolic acidosis as his bicarbonate level is also reduced. His carbon dioxide levels are normal, so he is not in type 2 respiratory failure and there is no indication for BiPAP.

E Ventilation/perfusion scan

This patient has had an acute insult to his lungs, following smoke and carbon monoxide exposure, with his chest X-ray appearances consistent with ARDS. As such, he is not maintaining optimal oxygen saturation despite maximal oxygen delivery, so a decision must be made immediately regarding additional ventilator support. Carrying out a scan at this point will delay treatment and will not benefit the patient. This investigation would be more appropriate in a stable patient in whom pulmonary embolism is suspected but who would not be appropriate for computed tomography pulmonary angiography (eg a pregnant woman or a patient in acute kidney injury who is at high risk of contrast-induced nephropathy).

32402

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Acute Respiratory Distress Syndrome

Expanded explanations

A 23-year-old African-Caribbean patient develops painful red nodules on both her shins over 2 days and has noticed painful knees, ankles and wrists. She also complains of mild sweats and weight loss over the past 2 months. Her chest x-ray is shown below.



Given the likely clinical diagnosis, what is the most likely outcome for her?

Your answer was incorrect

A Complete remission after a course of steroids

B Spontaneous resolution with no long-term sequelae

C Spontaneous resolution followed by further relapses

D Peripheral lung infiltrates and shortness of breath

E Increasing weight loss and neurological impairment

Explanation

B Spontaneous resolution with no long-term sequelae

This patient has acute sarcoidosis with erythema nodosum, polyarthralgia and mild constitutional symptoms. Her chest x-ray shows bilateral hilar lymphadenopathy (BHL) but clear lung fields. Around 90% patients have BHL at presentation. In 10–20% of cases, symptoms progress to shortness of breath, dry cough and chest pain. The majority of patients (80%) either stabilise or regress, so in the first instance treatment is not indicated. Non-pulmonary manifestations include neuropathy, Bell's palsy, hepatosplenomegaly, uveitis, conjunctivitis, lupus pernio, subcutaneous nodules, cardiomyopathy, arrhythmias, hypercalcaemia and renal stones. Treatment of acute sarcoidosis is usually supportive, including non-steroidal anti-inflammatory drugs (NSAIDs) for polyarthralgia, because most recover spontaneously.

Image source: https://commons.wikimedia.org/wiki/File:Hilar_Adenopathy_from_Sarcoidosis.jpg (https://commons.wikimedia.org/wiki/File:Hilar_Adenopathy_from_Sarcoidosis.jpg)

A Complete remission after a course of steroids

Sarcoidosis usually resolves spontaneously and steroids at first presentation would not be indicated. Steroids should be commenced if: symptoms worsen or cause limitation to daily activities, pulmonary function deteriorates, CXR signs worsen (e.g. development of cavitation or fibrosis) or if other organ system involvement develops.

C Spontaneous resolution followed by further relapses

Most patients recover spontaneously, and in 90% this is within 2 years. Less than 10% develop relapses after this point. Relapses and chronic disease are more likely if spontaneous remission does not occur within the first 2 years.

D Peripheral lung infiltrates and shortness of breath

The extent of infiltrates on CXR is used in staging of the disease and is a guide to prognosis, with the likelihood of relapse increasing as the extent of infiltrate worsens. This option is incorrect because it is uncommon.

E Increasing weight loss and neurological impairment

Neurological involvement does occur with sarcoidosis, but this is rare, occurring in 5-15% of patients. Neurological involvement tends to be present at presentation and often in disease with other significant extra-pulmonary features. This option is incorrect because it is an uncommon outcome.

31663

Rate this question:

Next Question

Previous Question Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 62-year-old man presents with a cough and shortness of breath which has increased in severity over the course of the past four days. The cough is productive with rusty-coloured/pus-stained sputum. He has a past history of mild asthma, for which he takes a salbutamol and beclometasone inhaler. He is pyrexial at 38.2 °C. His pulse is 89 bpm and regular, and his blood pressure is 105/72 mmHg. There are signs of right-sided consolidation on auscultation. You notice a cold sore on his right upper lip.

Investigations:

Investigations	**Results**	**Normal Values**
Haemoglobin (Hb)	132 g/l	135–175 g/l
White cell count (WCC)	14.4×10^9/l	$4–11 \times 10^9$/l
Platelets (PLT)	288×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine	110 µmol/l	50–120 µmol/l
C-reactive protein (CRP)	189 mg/l	0–10 mg/l
Glucose	6.1 mmol/l	3.5–5.5 mmol/l

Chest X-ray is shown below:



Which of the following is the most likely diagnosis?

Your answer was incorrect

A *Haemophilus influenzae*

B *Klebsiella*

C *Legionella pneumophilia*

D *Streptococcus pneumoniae*

E *Staphylococcus aureus*

Explanation

D *Streptococcus pneumoniae*

Pneumococcal pneumonia is the most common cause of community-acquired pneumonia in percentage terms and is often associated, as in this case, with the presence of cold sores. The chest x-ray here shows right upper lobe pneumonia, The treatment of choice is use of co-amoxiclav and clarithromycin in combination, as per the British Thoracic Society (BTS) guidelines.

Image source: https://commons.wikimedia.org/wiki/File:ZP_bronchopneumonie_-_sed_(7).jpg

B *Klebsiella*

Klebsiella is a cause of cavitating pneumonia in patients who are immunocompromised, eg with a history of chronic alcohol abuse. It does not fit with the scenario here.

A *Haemophilus influenzae*

H. influenzae is characteristically seen as a cause of infectious exacerbations of chronic obstructive pulmonary disease. It does not fit with the rusty-coloured sputum and cold sores here.

C *Legionella pneumophilia*

Legionella infection is associated with atypical pneumonia, including marked pyrexia, diarrhoea, abnormal liver function and hyponatraemia, which are not features seen here.

E *Staphylococcus aureus*

Like *Klebsiella*, *S. aureus* infection is associated with cavitating pneumonia. It occurs most commonly after influenza infection.

38393

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 41-year-old woman who is known to have brittle asthma was admitted to the Emergency Department with increased shortness of breath and a severe cough over the past few days, productive of yellow/green sputum. She was pyrexial 38.4°C, pulse was 90/min and regular, BP was 85/60 mmHg on admission, but improved with a fluid challenge. You review her on the ward some 36 h later, with her blood culture results. Her BP is now 105/70 mmHg, with pulse 75/min and regular. There are right basal crackles consistent with consolidation. She is still spiking fevers of 39.2°C every few hours.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	129 g/l	135 - 175 g/l
White cell count (WCC)	15.2× 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	203 × 10^9/l	150 - 400 × 10^9/l
C-reactive protein (CRP)	215 mg/l	< 10 mg/l
Sodium (Na^+)	132 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.1 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	162 µmol/l	50 - 120 µmol/l
Urea	14.1 mmol/l	2.5 - 6.5 mmol/l
Blood cultures	*Pseudomonas*	

Which of the following is the most appropriate antibiotic choice?

Your answer was incorrect

A IV colistin (polymyxin E)

B IV gentamicin

C IV meropenem

D IV tazocin

E Oral ciprofloxacin

Explanation

D IV tazocin

The choice here is between oral quinolone and IV therapy for *Pseudomonas*. Given that this patient is still spiking fevers and has an elevated urea, IV therapy with tazocin is most appropriate.

C IV meropenem

Carbapenems are an option for the treatment of *Pseudomonas*, although inducible resistance is a concern. For this reason, tazocin is the preferred option.

A IV colistin (polymyxin E)

Colistin is generally reserved for serious pseudomonal infections that are resistant to therapy with other agents.

B IV gentamicin

Gentamicin monotherapy is not recommended for the treatment of respiratory tract infections.

E Oral ciprofloxacin

Although oral ciprofloxacin is an option for pseudomonal infection, the severity of this patient's pneumonia drives us more towards IV tazocin as the preferred therapy. It is also now recommended that quinolone treatment is avoided wherever an effective alternative is available due to a small but significant risk of debilitating adverse effects.

70490

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 45-year-old man of African descent presents to the Respiratory Clinic with increased shortness of breath and a nagging dry cough, not relieved by a salbutamol inhaler. He has also been feeling non-specifically unwell over the past few weeks with fevers, general fatigue and lack of appetite. He has no past medical history of note and works as a solicitor.

Examination reveals a blood pressure of 122/72 mmHg and a pulse of 75 bpm and regular. There are scattered crackles on auscultation of the chest. His blood results and chest X-ray are shown below.

Investigations:

Investigations	Results	Normal Values
Haemoglobin (Hb)	130 g/l	135–175 g/l
White cell count (WCC)	10.0 × 10^9/l	4–11 × 10^9/l
Platelets (PLT)	200 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Creatinine	103 µmol/l	50–120 µmol/l
Erythrocyte sedimentation rate (ESR)	71 mm/first hour	0–10 mm/first hour



Given these results, which of the following is the most useful next investigation?

Your answer was incorrect

A Computed tomography (CT) chest

B Lung function testing

C Positron emission tomography (PET) scan

D Serum angiotensin-converting enzyme (ACE)

E Transbronchial biopsy

Explanation

A Computed tomography (CT) chest

Given the clinical features seen here, coupled with a CXR that shows hilar lymphadenopathy, sarcoidosis should be ruled out. The best initial investigation to do this is a high-resolution CT scan. This is likely to demonstrate pulmonary fibrosis and/or alveolitis, with good correlation with the diagnostic yield seen on transbronchial biopsy. Only 15% of patients with sarcoid have persistent pulmonary disease which requires steroid therapy.

B Lung function testing

Lung function testing may only see a relatively minor restrictive defect at presentation with sarcoid and is therefore not useful in confirming the diagnosis.

C Positron emission tomography (PET) scan

PET scanning is useful to demonstrate active inflammation and can differentiate benign from malignant lymphadenopathy. It is not a current first-line investigation.

D Serum angiotensin-converting enzyme (ACE)

While serum ACE can be elevated in sarcoidosis, elevated levels are only seen in 60% of patients. As such, it is not sensitive enough to confirm the diagnosis.

E Transbronchial biopsy

Although transbronchial biopsy will confirm granulomas consistent with sarcoidosis, it is not the next investigation of choice. It is only likely to be instigated in the event the diagnosis is in doubt following non-invasive investigations. Additionally, there may be more easily accessible biopsy sites if one is required.

38296

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 45-year-old farmer comes to the Respiratory Clinic for review. He has been diagnosed by the GP as having asthma, for which he is treated with high-dose fluticasone/salmeterol and salbutamol ,without significant improvement in his symptoms. Examination reveals a BP of 122/72 mmHg, with pulse 70/min and regular. There is scattered wheeze and coarse crackles on auscultation of the chest. Respiratory function tests show an obstructive picture. A thoracic HRCT shows bilateral centrally dilatated thickened airways with ‘signet rings’.

Which of the following tests would be most useful to confirm the suspected diagnosis?

Your answer was incorrect

A IgE

B IgG

C ANCA

D ANA

E C3 and C4

Explanation

A IgE

The HRCT changes described are classical of proximal bronchiectasis. The diagnosis is therefore allergic bronchopulmonary aspergillosis (ABPA). The total IgE will be raised in ABPA, especially during a flare-up, and therefore represents the preferred investigation.

D ANA

ANA is raised in connective tissue disease such as SLE, although there are no other features to suggest connective tissue disease as the underlying diagnosis.

B IgG

IgG can be low in bronchiectasis secondary to hypogammaglobulinaemia, although presentation for the first time at an earlier age would be expected because of recurrent respiratory tract infection.

C ANCA

ANCA is most likely to be elevated in vasculitis. Eosinophilic vasculitis (Churg–Strauss) is associated with more features than just worsening of asthma, including skin lesions and mononeuritis, which aren't seen in this scenario.

E C3 and C4

Complement factors are consumed most when there is active tissue inflammation, such as with SLE. The absence of other features such as glomerulonephritis counts against C3 and C4 as being useful here.

2329

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 62-year-old patient presents complaining of shortness of breath which has increased over the past 2–3 months. He smokes 10 cigarettes per day. Examination reveals a BP of 132/82, with pulse 74 (AF). He has hyper-expanded lung fields and bilateral poor air entry. There is quiet wheeze and bilateral pitting oedema of the ankles. BMI is 26.

Blood gas investigation reveals the following:

Investigation	**Result**	**Normal Value**
pCO_2	7.2 kPa	4.6–6.0 kPa
pO_2	9.1 kPa	10.5–13.5 kPa
pH	7.33	7.35–7.45
Bicarbonate (HCO_3^-)	29 mmol/l	24–30 mmol/l

Lung function results reveal the following:

Investigation	**Result**
RV	120%
TLC	110%
KCO	70% (as a percentage of the expected values)

His chest x-ray shows the following:



Which of the following diseases fits best with the above presentation?

Your answer was incorrect

A	Emphysema
B	Asthma
C	Obesity hypoventilation syndrome
D	Interstitial lung disease
E	Lymphangiomyomatosis

Explanation

A Emphysema

Emphysema gives a chronic CO_2-retaining picture, with an increased residual volume (RV), slightly high total lung capacity (TLC) and high RV/TLC percentage indicative of obstructive lung pathology. It's consistent with the compensated respiratory acidosis and evidence of right heart failure. The chest x-ray shows mild overexpansion with slightly flattened diaphragm, in keeping with emphysema.

Image source: https://commons.wikimedia.org/wiki/File:Chest_x-ray_-_posteroanterior_view.jpg

C Obesity hypoventilation syndrome

Obesity hypoventilation is associated with reduced TLC, and would only be associated with hypercapnia in severe disease complicated by pulmonary hypertension and right heart failure.

B Asthma

Although acute asthma is associated with increased residual volume, this is not really a feature of chronic controlled asthma, nor is hypercapnia.

D Interstitial lung disease

Although interstitial lung disease is associated with hypoxia, hypercapnia is unexpected unless the patient has a superimposed acute respiratory tract infection.

E Lymphangiomyomatosis

This usually presents in women of child-bearing age with shortness of breath and recurrent pneumothoraces.

2318

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

You are on call on Boxing Day and you are referred a 61-year-old lady with shortness of breath. Her only other complaint is of recent diarrhoea. She is a smoker of 30/day.

She tells you that she consumed a significant volume of alcohol the previous day, and does not remember going to bed; she woke up on the floor.

On examination she looks unwell. Her temperature is 38.5°C. On auscultation of her chest she has coarse crackles at the right base. She has a distended abdomen with dullness in the flanks.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	111 g/l	135 - 175 g/l
White cell count (WCC)	12.3 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	9.5 × 10^9/l	1.5 - 7.0 × 10^9/l
Platelets (PLT)	88 × 10^9/ll	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	101 fl	80 - 100 fl
Sodium (Na^+)	130 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.4 mmol/l	3.5 - 5.0 mmol/l
Urea	7.8 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	77 μmol/l	50 - 120 μmol/l
International normalized ratio (INR)	1.4	< 1.1
Bilirubin	16 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	73 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	300 u/l	30 - 150 u/l
Albumin	30 g/l	35 - 55 g/l
Gamma glutamyl-transferase (GGT)	100 u/l	< 50 u/l
Creatinine kinase (CK)	700 u/l	22 - 198 u/l

Her chest X-ray is shown below:



What is the most likely diagnosis?

Your answer was correct

A	Mycoplasma pneumonia
B	Legionella pneumonia
C	Aspiration pneumonia
D	Streptococcal pneumonia
E	Klebsiella pneumonia

Explanation

C Aspiration pneumonia

This is aspiration pneumonia. This lady has signs of chronic alcohol excess (thrombocytopenia, macrocytosis, raised INR and abnormal liver function tests – in particular the gamma GT). She has consumed too much alcohol on Christmas Day, fallen (note creatine kinase) and aspirated. The chest X-ray shows a right lower lobe pneumonia consistent with an aspiration pneumonia.

A Mycoplasma pneumonia

Mycoplasma pneumonia usually affects young adults and is associated with dry cough, fever, headache and malaise.

B Legionella pneumonia

Legionella pneumonia is associated with poorly maintained water systems (hotel air-conditioning, etc.), and presents with symptoms such as cough, dyspnoea, fever and myalgia. In addition, while Legionella can cause hepatic dysfunction, the clinical findings are compatible with ascites and her bloods are consistent with chronic rather than acute liver disease.

D Streptococcal pneumonia

The history is suggestive of aspiration pneumonia rather than classical community-acquired streptococcal pneumonia, which would be expected to present with a longer history of productive cough, fever, etc.

E Klebsiella pneumonia

Klebsiella pneumonia is associated with lung destruction, including cavitation and abscess formation.

70082

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 71-year-old man returns from a veterans' convention in Spain. A few days later he becomes unwell with high fevers, rigors, a dry cough and severe headaches. On examination his BP is 105/60 mmHg, pulse is 92/min and regular and his temperature is 38.°9C. There is evidence of left basal consolidation and dullness suggestive of a pleural effusion, and his O_2 saturation is 93% on air.

Which one of the following features might point towards *Legionella* infection if found in this patient?

Your answer was incorrect

A Elevated transaminases

B Onset following influenza illness

C Cavitation on chest X-ray

D History of contact with birds

E Cold agglutinins

Explanation

A Elevated transaminases

Legionella has an incubation period of 2–10 days, and presents with fever, rigors, headache, myalgia, dry cough, progressively increasing respiratory distress and confusion. Diarrhoea and vomiting are common, as are non-specific liver function test abnormalities. Relative bradycardia may be seen in some patients, in contrast to other acute pneumonias. Pleural effusion may be seen in up to 50% and chest X-ray appearance may take up to four months to return to normal. Diagnosis is made with serology, or with PCR on urinary Legionella antigen. Treatment is with macrolides as first-line therapy. Complications include pericarditis, encephalitis and kidney disease.

C Cavitation on chest X-ray

Cavitation on X-ray may imply necrotising pneumonia, as can occur with any pneumonia. It would occur following sever sepsis, especially with pneumococcal pneumonia. Cavitating lesions can also occur with tuberculosis. The acute onset makes this a less likely diagnosis for this patient.

B Onset following influenza illness

Post influenza pneumonia is most typically associated with Staphylococcus aureas. There is no link with legionella.

D History of contact with birds

Contact with birds can lead to infection with *Chlamydia sittaci*, which causes psittacosiss following spread from birds, mostly parrots. It presents with headache, fever, dry cough, arthralgia and diarrhoea and vomiting. The chest X-ray shows patchy consolidation in this condition.

E Cold agglutinins

Cold agglutinins are seen in patients with infectious mononucleosis. Infective mononucleosis is caused by EBV. Following spread is by droplets or saliva and incubation over four weeks. In early childhood it is nearly asymptomatic, but when adolescents are infected they can develop fever, sore throat, malaise and lympho-adenopathy. Blood films show lymphocytosis and atypical lymphocytes.

32372

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 34-year-old woman is referred to the Respiratory Clinic. She has had a long history of asthma since childhood and is maintained on combination inhaler therapy with fluticasone 250 µg/salmeterol 50 mg BD. On the advice of her doctor she moved away from London to the countryside and was well. Around 2 years ago she began keeping a horse and since then has noticed a gradual deterioration in her asthma control, with increased coughing up of mucus plugs. She has been started on high-dose oral corticosteroids by her GP, although she feels her symptoms have not significantly improved.

On examination in the clinic there was clear evidence of bilateral wheeze. Her peak flow was measured at 350 l/min.

Investigations reveal the following:

Investigation	**Result**	**Normal values**
Immunoglobulin E (IgE)	1100 ng/ml	
Peripheral blood	Eosinophil count 0.6 × 10^9/l	0.04–0.4 × 10^9/l
IgG	Precipitins to *Aspergillus*	
Skin prick	+ to *Aspergillus*	
Chest X-ray (CXR)	Evidence of pulmonary infiltration	

Which of the following represents the optimal next step in this patient?

Your answer was incorrect

A Increased dose of inhaled steroids

B Addition of oral montelukast

C Stop oral steroids and start itraconazole

D Continue oral corticosteroids alone

E Add oral itraconazole to her regime

Explanation

E Add oral itraconazole to her regime

There is evidence that this patient is suffering from bronchopulmonary aspergillosis. The factors that promote evolution from atopic asthma to this disease are unknown, but it appears that the type of CD4 response to aspergillus antigen is crucial. Management involves the use of oral corticosteroids, and there is evidence that additional prolonged courses of itraconazole (up to 4 months) may diminish the steroid requirement. Aggressive therapy with oral corticosteroids is known to reduce the risk of bronchiectasis that results from mucus plug impaction and subsequent localised infection.

C Stop oral steroids and start itraconazole

Without oral corticosteroids ABPA does not improve on itraconazole alone; hence combination therapy is required.

A Increased dose of inhaled steroids

Inhaled corticosteroids do not relieve the inflammation seen here, even at high dose.

B Addition of oral montelukast

Oral montelukast is utilised in the treatment of adult asthma for patients who fail to gain control of symptoms on a combination of inhaled corticosteroids and long-acting beta agonist.

D Continue oral corticosteroids alone

Itraconazole has additional anti-inflammatory and anti-fungal effects with respect to the treatment of ABPA, and leads to more effective resolution of the condition.

18556

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A 45 year old man with acute pancreatitis and developing ARDS is referred to the Intensive Therapy Unit. On examination his BP is 135/85 mmHg; pulse is 95/min and regular. There are scattered crackles on auscultation of the chest and O_2 saturation is only 86% on high flow oxygen. CXR shows patchy shadowing. The patient is intubated.

Which of the following is correct with respect to the ventilator settings?

Your answer was incorrect

A Low PEEP, Low tidal volume, respiratory rate up to 35 breaths/min

B Low PEEP High tidal volume, respiratory rate up to 25 breaths/min

C High PEEP, High tidal volume, respiratory rate up to 35 breaths/min

D High PEEP High tidal volume, respiratory rate up to 25 breaths/min

E High PEEP low tidal volume, respiratory rate up to 35 breaths/min

Explanation

E High PEEP low tidal volume, respiratory rate up to 35 breaths/min

The answer is High PEEP, low tidal volume, respiratory rate up to 35 breaths/min -

Low tidal volumes are thought to improve survival in ARDS by reducing excessive lung stretch; hence it is the preferred option here. High PEEP is thought to reduce progression of ARDS versus lower levels. Respiratory rate up to 35 breaths / min is required to deliver the predicted minute volume needed for most patients.

http://www.ncbi.nlm.nih.gov/pubmed/23740697 (http://www.ncbi.nlm.nih.gov/pubmed/23740697)

C High PEEP, High tidal volume, respiratory rate up to 35 breaths/min

High PEEP, High tidal volume, respiratory rate up to 35 breaths/min, is incorrect because high tidal volume may accentuate damage to lung tissue, although high PEEP is in fact has a positive impact on reducing ARDS progression.

A Low PEEP, Low tidal volume, respiratory rate up to 35 breaths/min

Low PEEP, Low tidal volume, respiratory rate up to 35 breaths/min, is incorrect because low levels of PEEP do not reduce the progression of ARDS versus higher levels.

B Low PEEP High tidal volume, respiratory rate up to 25 breaths/min

Low PEEP High tidal volume, respiratory rate up to 25 breaths/min, is incorrect because high tidal volume may accentuate stretch damage to lung tissue.

D High PEEP High tidal volume, respiratory rate up to 25 breaths/min

High PEEP High tidal volume, respiratory rate up to 25 breaths/min is incorrect because high tidal volume may accentuate damage to lung tissue, although high PEEP is in fact has a positive impact on reducing ARDS progression.

40408

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Acute Respiratory Distress Syndrome

Expanded
explanations

A 17-year-old man presents to the Respiratory Clinic for review. He has suffered recurrent otitis media, gastrointestinal infections and, most recently, four episodes of pneumonia over the past year. He tells you that he has always been sickly as a child and, as such, is below the 25th centile with respect to height.

On examination, his blood pressure is 115/82 mmHg; pulse is 80 bpm and regular. You cannot hear his heart sounds on auscultation and there are coarse bibasal on listening to his chest.

Investigations:

Investigation	Result	Normal value
Haemoglobin (Hb)	131 g/l	135–175 g/l
White cell count (WCC)	8.2×10^9/l	$4–11 \times 10^9$/l
Platelets (PLT)	203×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	95 µmol/l	50–120 µmol/l
Electrocardiogram (ECG)	Right axis deviation, positive QRS in aVR, poor R wave progression	

His chest x-ray is shown below.



Which of the following is the most likely diagnosis?

Your answer was incorrect

A Allergic bronchopulmonary aspergillosis

B Idiopathic nasal polyposis

C IgA deficiency

D IgG deficiency

E Primary ciliary dyskinesia

Explanation

E Primary ciliary dyskinesia

Classically, patients present with respiratory tract infections including pneumonia, upper respiratory tract infections, otitis and sinusitis. Situs inversus is the reversal of the internal organs which is in keeping with the quiet heart sounds, ECG appearances and chest x-ray making the diagnosis of primary ciliary dyskinesia a strong possibility. The chest x-ray also shows The signs of bronchiectasis including tram lines which is commonly found alongside situs inversus.

Image source:
https://upload.wikimedia.org/wikipedia/commons/f/f0/Situs_inversus_due_to_Kartagener%27s_syndrome.png

C IgA deficiency

IgA deficiency is usually associated with increased risk of gastrointestinal infections and immunoglobulin deficiencies are not consistent with dextrocardia per se.

A Allergic bronchopulmonary aspergillosis

Patients with allergic bronchopulmonary aspergillosis usually present with uncontrollable asthma despite treatment. A chest X-ray in these patients would typically show changes in keeping with bronchiectasis, such as ‘tram lines’ and ‘gloved finger shadows’.

B Idiopathic nasal polyposis

The recurrent infections and chest X-ray findings are not consistent with nasal polyposis. Patients with nasal polyposis may complain of nasal fullness or stuffiness along with difficultly sleeping due to nasal obstruction.

D IgG deficiency

Isolated IgG deficiency rarely causes such recurrent infections and immunoglobulin deficiency is not consistent with dextrocardia. Patients with combined IgA and IgG deficiency are at much greater risk of recurrent infections.

39348

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Bronchiectasis

Media

Expanded explanations

Flashcards

A 47-year-old homeless man is brought to the Emergency Department with fever, cough with purulent expectoration, breathlessness and chest pain. He states that his chest pain is on the right side and is aggravated by deep inspiration. His symptoms have worsened over the past seven days.

He has poor dental hygiene and is tachypnoeic at rest. On examination, his blood pressure is 100/60 mmHg, heart rate 112 bpm, respiratory rate 28 breaths per minute and temperature 38.9°C. Respiratory examination revealed decreased air entry over the right hemithorax.

Which of the following findings on pleural fluid analysis would mandate placement of an intercostal drain?

Your answer was incorrect

A Gram-positive organism in the pleural fluid

B Pleural fluid glucose 3.6 mmol/l

C Pleural fluid lactate dehydrogenase (LDH) 960 U/l

D Pleural fluid pH 7.3

E Presence of turbid pleural fluid

Explanation

A Gram-positive organism in the pleural fluid

Identification of organisms on Gram staining, the presence of frank pus on needle aspiration and identification of an organism on culture of the pleural fluid are absolute indications for intercostal tube placement and drainage. This patient likely has an empyema due to a mixed aerobic and anaerobic infection, and initiation of broad-spectrum antibiotics is the best initial step, followed by placement of an intercostal drain.

C Pleural fluid lactate dehydrogenase (LDH) 960 U/l

In the absence of availability of pleural fluid pH, a pleural fluid LDH level of > 1000 U/l is considered to be an indication for drainage of pleural fluid.

B Pleural fluid glucose 3.6 mmol/l

A pleural fluid glucose level of < 3.3 mmol/l is considered to be indicative of a complicated parapneumonic effusion and would mandate drainage.

D Pleural fluid pH 7.3

A pleural fluid pH of < 7.2 is an indication for drainage.

E Presence of turbid pleural fluid

The presence of frank pus, rather than turbid pleural fluid, is an indication for intercostal tube drainage. 72783

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Empyema

Expanded explanations

A 34-year-old woman presents to the Respiratory Clinic with a gradual onset of exertional dyspnoea over the previous 18 months. She has no cough, sputum or haemoptysis. She describes recent fatigue and several episodes of presyncope. She is a non-smoker and has no significant past medical or family history.
On examination, she has a respiratory rate of 19 breaths per minute, an oxygen saturation of 94% on air, a BMI of 36 kg/m^2 and a loud pulmonary component to the second heart sound.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	160 g/l	115–155 g/l
White cell count (WCC)	6.0 × 10^9/l	4–11 × 10^9/l
Platelets	250 × 10^9/l	150–400 × 10^9/l
Plasma D-dimers	130 (upper limit of normal = 285)	
ECG	Sinus rhythm, right axis deviation, right ventricular hypertrophy	
Chest X-ray	Enlarged hilar bilaterally, oligaemic peripheral lung fields	

Which investigation would be most helpful in establishing the diagnosis?

Your answer was incorrect

A 24-hour ECG monitoring

B Dobutamine stress echocardiography

C Exercise ECG

D Respiratory function tests

E Ventilation/perfusion lung scan

Explanation

E Ventilation/perfusion lung scan

The diagnosis is either primary pulmonary hypertension (PPH) or multiple recurrent pulmonary emboli. PPH is a rare disorder of unknown aetiology, which typically affects young women. There is an average delay of two years between symptom onset and diagnosis, by which time cor pulmonale usually develops. Echocardiography shows right ventricular enlargement, and cardiac catheterisation will demonstrate a raised pulmonary artery pressure. A ventilation/perfusion (V/Q) scan may show patchy filling defects that are different from those seen in pulmonary embolism and remains part of the diagnostic work-up.

Multiple pulmonary emboli are a possibility, given this patient's obesity. It is possible for an accumulation of small pulmonary emboli to eventually lead to secondary pulmonary hypertension and right heart failure. Diagnosis is with V/Q scan or CT pulmonary angiography, which will demonstrate multiple perfusion defects. If V/Q is negative, then PPH becomes a more likely possibility.

C Exercise ECG

An exercise ECG would help to pick up an exercise-induced arrhythmia but would not be of any diagnostic utility in this patient's case.

A 24-hour ECG monitoring

Ambulatory ECG monitoring might help to uncover an arrhythmia causing intermittent dyspnoea and symptoms of palpitations, but it will not add any additional diagnostic insight into this patient's case.

B Dobutamine stress echocardiography

A dobutamine stress echocardiography will help ascertain how cardiac function performs under physiological stress, but it will not help to diagnose either of the above two differentials.

D Respiratory function tests

Pulmonary function testing will give us no more additional information that will help us to reach a diagnosis. The diffusing capacity for carbon monoxide may be reduced, given the reduced capacity for oxygen transfer due to pulmonary hypertension. However, pulmonary function testing will not help make either of these diagnoses.

73466

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pulmonary Embolism

Media

Expanded explanations

Flashcards

A woman presents to clinic in her eighth week of pregnancy. She is concerned about her husband's family history of cystic fibrosis (CF), as her husband's brother has cystic fibrosis and her husband has been confirmed as a carrier. The patient does not wish to know about her risk of developing CF and is reluctant to be tested herself, but is concerned about the health of her child. There is a 1:20 prevalence of CF carriage in the area where she lived as a child and husband and wife are not consanguineous.

What is the chance of her child having cystic fibrosis?

Your answer was correct

A 1/4

B 1/40

C 1/80

D 1/160

E 1/320

Explanation

C 1/80

CF is an autosomal recessive disease, so two abnormal alleles are required for the disease to be manifest. As a carrier (with one abnormal gene) the husband has a 1:2 chance of passing the abnormal gene to his offspring. If one assumes that the woman is of the same genetic stock as those in the area of her birth, she has a 1:20 chance of being a carrier, so the chance of the child receiving an abnormal allele from both parents is 1/2 (male risk) × 1/20 (female carrier risk) × 1/2 (female risk of passing a gene on) = 1/80.

A 1/4

1 in 4 is the risk of a child being affected by CF if both of the parents are confirmed to be carriers for the disease.

B 1/40

1 in 40 would be the risk of offspring suffering from a recessive condition if one partner was a homozygote for the abnormal gene, and the carrier frequency was 1 in 20.

D 1/160

This would be the correct answer if the carrier frequency was 1 in 40, not 1 in 20 as it is.

E 1/320

This would be the correct answer if the carrier frequency was 1 in 80.

9835

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Cystic Fibrosis

Media

Expanded explanations

Flashcards

An 18-year-old male attends his GP complaining of fever and a productive cough present for 2 weeks. He initially thought it was flu but the symptoms have been getting worse and not better. He is concerned since he is the main breadwinner at home for his elderly parents, who have just returned from a trip to India. On further questioning he admits to having lost his appetite recently, and at night-time suffers from terrible sweating, drenching the bed sheets. On the basis of the history and his clinical examination, the GP suspects a diagnosis of tuberculosis. An urgent referral is made to the local Chest Clinic to confirm the diagnosis, and tests confirm that the diagnosis is correct. The patient is commenced on standard treatment for pulmonary tuberculosis. He returns to the GP several weeks later, stating that he has noticed deterioration in his eyesight since his diagnosis.

Which of the following is most likely to be responsible for this side effect?

Your answer was incorrect

A Ethambutol

B Rifampicin

C Isoniazid

D Pyrizinamide

E Streptomycin

Explanation

A Ethambutol

The standard medical treatment of pulmonary tuberculosis consists of an initial phase of 2 months on rifampicin, isoniazid, pyrazinamide and ethambutol. This is followed by a further 4-month continuation phase on isoniazid and rifampicin only. There are some combination preparations that may also be used, but this depends on their availability. Prior to treatment, it is imperative to stress the importance of compliance. Also, both liver and renal function should be assessed prior to commencing therapy. Visual acuity, colour vision and visual fields should also be tested, preferably by an ophthalmologist, both before and during treatment with ethambutol since ocular toxicity may occur.

C Isoniazid

This is associated with peripheral neuropathy in high doses, GI upset, agranulocytosis and haemolytic anaemia.

B Rifampicin

This is associated with orange discoloration of secretions such as tears and urine, GI symptoms and alteration of hepatic function.

D Pyrizinamide

This is associated with GI upset and hepatotoxicity.

E Streptomycin

This is associated with vestibular and renal toxicity.

6806

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 77-year-old woman was referred to the Respiratory Outpatient Clinic with increasing shortness of breath. Her chest X-ray is shown below.



What was her underlying diagnosis?

Your answer was incorrect

A Adenocarcinoma

B Pneumothorax

C Small cell lung cancer

D Tuberculosis

E Chronic spontaneous pneumothorax

Explanation

D Tuberculosis

This question is a ‘spot diagnosis’. Clues that this is a thoracoplasty are that there is evidence of rib resection in conjunction with loss of lung volume. The CXR shows a thoracoplasty. This lady had tuberculosis in the 1940s, prior to the use of anti-tuberculosis chemotherapy. She had a thoracoplasty, which was one of the main surgical treatments for tuberculosis between 1930 and 1955. Several ribs were resected, which reduced the thoracic volume and collapsed the underlying lung. The aim was to close the tuberculous cavity and ‘rest the lung’. Control of pulmonary tuberculosis and survival was good. However, patients were left with severe chest deformity and the associated respiratory compromise, a restrictive defect.

B Pneumothorax

Symptoms of pneumothorax would include sudden onset of sharp, unilateral chest pain with shortness of breath. On a chest X-ray there would be an air rim and collapse of lung. However, there would not be rib resection as in this case.

A Adenocarcinoma

Adenocarcinoma would present as a mass on the chest X-ray. There would not be rib resection and loss of lung volume. Other symptoms and signs may include anaemia and weight loss.

C Small cell lung cancer

As with adenocarcinoma, chest X-ray would show a mass rather than loss of lung volume. Small cell lung cancer is a fast-growing type of lung cancer, spreading more quickly than non-small cell lung cancer and therefore it is the most aggressive form of lung cancer.. Small cell lung cancer is more common in men than women. Almost all cases of small cell lung cancer are due to cigarette smoking. It is very rare in people who have never smoked.

E Chronic spontaneous pneumothorax

The rib resection counts against this being a chronic pneumothorax and fits with a thoracoplasty for tuberculosis.

70127

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 30-year-old man from London with no significant past medical history is engaged on a holiday to Peru to climb the Inca Trail. He is a non-smoker. After climbing above 3500 m he has developed a headache, associated with nausea and vomiting. He also feels tired and complains of insomnia. You are the expedition medic. His respiratory rate is 22/min and his pulse is 90/min. Chest is clear on auscultation. No investigations are available as the nearest hospital is over 100 km away.

What is the most likely diagnosis?

Your answer was incorrect

A Exhaustion

B Hangover

C Migraine

D Acute mountain sickness

E Hypothermia

Explanation

D Acute mountain sickness

There are three types of altitude illness: acute mountain sickness (AMS), high-altitude cerebral oedema (HACE) and high-altitude pulmonary oedema (HAPE). This man has acute mountain sickness with some evidence of cerebral oedema. The exact cause is unknown but it is thought to be secondary to cerebral oedema and raised intracranial pressure. It is more common in people who live at low altitude.

Symptoms and signs of AMS are non-specific; however, diagnosis can be made using the criteria named in 'The Lake Louise Score', namely, the setting of recent gain in altitude, the presence of headache and at least one of the following symptoms:

- Gastrointestinal – anorexia, nausea and vomiting
- Fatigue and weakness
- Dizziness or light-headedness
- Difficulty sleeping.

He does not improve and progresses to develop ataxia and drowsiness. This is the development of high-altitude cerebral oedema, which can be considered as end-stage AMS.

Treatment is immediate descent, oxygen and intravenous dexamethasone.

Acetazolamide may be used for prevention and treatment of AMS. It is a carbonic anhydrase inhibitor that causes intracellular acidosis. It has a diuretic effect which can be useful in AMS/HACE and HAPE. It is also beneficial as it shifts the oxygen-dissociation curve to the right and therefore at lower partial pressures, more oxygen is released to the tissues.

C Migraine

This would be a likely differential diagnosis if he was not at altitude, although it would be unlikely to present for the first time at this age and there is no history of migraine triggers reported as part of the scenario.

A Exhaustion

This is more than exhaustion, since he has symptoms suggestive of altitude sickness. The severe headache, nausea and vomiting are possible pointers to cerebral oedema.

B Hangover

There is no history to suggest alcohol consumption, and it is extremely unlikely the patient would be able to climb to above 3500 m if he had consumed significant amounts of alcohol.

E Hypothermia

He has classical symptoms of mountain sickness, and hypothermia is not the most likely diagnosis here. Significant signs consistent with attempts to maintain core temperature, such as shivering, would be expected and are not reported here.

70106

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

High Altitude Medicine

Expanded
explanations

A 39-year-old man presented to his GP with a 3-month history of fever, rhinitis and weight loss. Two weeks ago he had an exacerbation of asthma and was treated with a course of erythromycin. He also complained of weakness in his hands and legs. He has a past medical history of angina. He is an ex-smoker and stopped 5 years ago.

On examination he has a rash on his shins, shown below.



Auscultation of his chest reveals a mild expiratory wheeze. Examination was otherwise unremarkable. Urinalysis was normal.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	121 g/l	135 - 175 g/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
White cell count (WCC)	9.4×10^9/l	$4.0 - 11.0 \times 10^9$/l
Neutrophils	5.3×10^9/l	$1.5 - 7.0 \times 10^9$/l
Lymphocytes	1.8×10^9/l	$1.0 - 4.0 \times 10^9$/l
Monocytes	0.3×10^9/l	$0.0 - 0.8 \times 10^9$/l
Basophils	0.08×10^9/l	$< 0.1 \times 10^9$/l
Eosinophils	$1.9 \times \times 10^9$/l	$0.04 - 0.4 \times 10^9$/l
Platelets (PLT)	411×10^9/l	$150 - 400 \times 10^9$/l
Erythrocyte sedimentation rate (ESR)	82 mm/h	1 - 20 mm/h

Sodium (Na^+)	138 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Urea	4.1 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	111 µmol/l	50 - 120 µmol/l

What is the diagnosis?

Your answer was incorrect

A Microscopic polyangiitis

B Granulomatosis with polyangiitis

C Eosinophilic pneumonia secondary to erythromycin

D Eosinophilic granulomatosis with polyangiitis (EGPA)

E Allergic bronchopulmonary aspergillosis

Explanation

D Eosinophilic granulomatosis with polyangiitis (EGPA)

The patient has systemic symptoms, asthma, eosinophils $>1.5 \times 10^9$/l and evidence of vasculitis in two or more non-lung organs (Lanham's criteria):

- Cutaneous vasculitis (see picture)
- Mononeuritis multiplex
- A cardiac history.

The American Criteria of Rheumatology require four out of six of the following to be present for a diagnosis of Eosinophilic granulomatosis with polyangiitis (EGPA):

- The presence of asthma
- Eosinophils >10% in the peripheral blood
- Evidence of a neuropathy in a vasculitic pattern, e.g. mononeuritis multiplex
- Transient pulmonary infiltrates
- A history of sinus disease
- Evidence of intravascular eosinophils on biopsy.

Renal involvement in EGPA syndrome is very rare.

B Granulomatosis with polyangiitis

Granulomatosis with polyangiitis is associated with glomerulonephritis, which has not been demonstrated here.

A Microscopic polyangiitis

Microscopic polyangiitis is associated with renal disease. The absence of any evidence of glomerulonephritis counts against this as the underlying diagnosis.

C Eosinophilic pneumonia secondary to erythromycin

The symptoms preceded erythromycin administration, making eosinophilic pneumonia unlikely.

E Allergic bronchopulmonary aspergillosis

Allergic bronchopulmonary aspergillosis usually occurs in asthmatics, but would not produce the signs and symptoms of vasculitis as in this patient, including the obvious skin lesions.

70091

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219

Responses Total:	301
Responses - % Correct:	27%

Eosinophilic Granulomatosis with Polyangiitis

Expanded
explanations

A 51-year-old woman is admitted with worsening fevers and night sweats over the past 2 months, associated with joint pain and fatigue. She has been previously well, other than a recent attendance at her general practitioner (GP) for a long-standing productive cough some 5 months earlier. Travel of note includes a spell working at a church mission in India during the past year.

A chest X-ray, as shown, is undertaken and sputum samples are awaited.



Other investigations are as follows:

Haemoglobin (Hb)	101 g/l	(135–155 g/l)
White cell count (WCC)	3.4×10^9/l (neutrophils 1.98)	($4–11 \times 10^9$/l)

What is the most appropriate management step with regard to isolation for this patient?

Your answer was incorrect

A Isolate in a negative pressure room

B Isolate in a positive pressure room

C Isolate in a standard patient bay

D Isolate in a standard side room

E No isolation required

Explanation

A Isolate in a negative pressure room

The chest X-ray and symptoms are highly suggestive of active miliary tuberculosis (TB). The National Institute for Health and Care Excellence (NICE) guidelines recommend managing patients with a high risk for TB in a negative pressure room. A negative pressure room is one where the air from the room is sucked out into dedicated ducting through a filter and into the outside air, at a distance from all other air intakes.

B Isolate in a positive pressure room

The chest X-ray and symptoms are highly suggestive of active miliary TB. NICE guidelines recommend managing patients with a high risk for TB in a negative pressure room, rather than a positive pressure one. In a positive pressure room, ventilation allows air to flow out of the room into the general environment, instead of into the room. This is useful in immunocompromised patients. Here the patient has mild neutropenia; however, this would not generally be an indication for positive pressure.

C Isolate in a standard patient bay

Source-isolating a patient in a bay is less effective than in a side room and is usually only undertaken when side rooms or other facilities are not available. This would not be an appropriate step to manage isolation of miliary TB.

D Isolate in a standard side room

The chest X-ray and symptoms are highly suggestive of active miliary TB. NICE guidelines recommend managing patients with a high risk for TB in a negative pressure room. Patients who are low risk for TB may be managed in a side room.

E No isolation required

The chest X-ray and symptoms are highly suggestive of active miliary TB. Managing the patient in a general bay with other patients risks transmission of TB.

71775

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

National Institute for Health and Care Excellence. 2016. Tuberculosis. NICE guideline [NG33].
nice.org.uk/guidance/ng33/chapter/recommendations#managing-active-tb-in-all-age-groups
(https://www.nice.org.uk/guidance/ng33/chapter/recommendations#managing-active-tb-in-all-age-groups)

An 18-year-old man with cystic fibrosis is admitted to the hospital with an infective exacerbation of his condition. His latest sputum grew *Burkholderia cepacia*. His sister also has cystic fibrosis (CF) and has problems with abdominal pains. He has previously been relatively well, plays for a local football team and is in the first year of a law course at university. He has maintained his weight so far.

Which of the following would you most expect to find?

Your answer was incorrect

A Severe restrictive pattern on spirometry

B Exocrine pancreatic insufficiency

C Diabetes mellitus

D Low thiamine and vitamin B_{12}

E Symptoms of chronic constipation

Explanation

B Exocrine pancreatic insufficiency

Exocrine pancreatic insufficiency is a feature of cystic fibrosis and occurs long before evidence of dysglycaemia and frank diabetes mellitus. Pancreatic exocrine supplements and fat-soluble vitamins (A, D, E and K) are a key feature of treatment.

C Diabetes mellitus

Endocrine pancreatic dysfunction leading to diabetes mellitus occurs later in the disease and is associated with weight loss and symptoms of dysglycaemia.

A Severe restrictive pattern on spirometry

There is good evidence of maintained respiratory function seen here (continuing to play in a football team). An obstructive picture is more likely to be seen on spirometry in CF in this situation.

D Low thiamine and vitamin B_{12}

Small bowel malabsorption leading to low thiamine and B_{12} is unlikely, given pancreatic enzyme supplementation.

E Symptoms of chronic constipation

Although patients with cystic fibrosis may present with symptoms of constipation, the tendency to constipation in adults is modulated by pancreatic enzyme insufficiency.

71073

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 62-year-old cleaner comes to the Respiratory Clinic complaining of increasing shortness of breath and a persistent cough. She used to work sweeping up for a plumber and boiler maker. She has begun to get tired while walking the dog, which is spoiling the enjoyment of her time off. She smokes five cigarettes per day.

On examination, she is finger-clubbed and you can hear fine end-inspiratory crackles.

Investigations reveal the following:

Investigation	Result	Normal value
Haemoglobin (Hb)	128 g/l	115–155 g/l
White cell count (WCC)	4.5 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	210 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135–145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	140 µmol/l	50–120 µmol/l
Forced expiratory volume in 1 s (FEV_1)	2.7	
Forced vital capacity (FVC)	3.1	
Chest X-ray (CXR)	Shown below	



Which of the following is the most likely diagnosis?

Your answer was correct

A	Asthma
B	Solderer’s lung
C	Asbestosis
D	Chronic obstructive pulmonary disease (COPD)
E	Idiopathic pulmonary fibrosis

Explanation

C Asbestosis

The clue is very clear that, in her work at a plumbers, this woman is likely to have had significant exposure to asbestos. The chest X-ray findings of bilateral pleural plaques also point towards asbestos exposure. High-resolution computed tomography (CT) is important to confirm pulmonary fibrosis, and pulse oximetry normally reveals a degree of hypoxaemia. Despite extensive trials, neither corticosteroids nor second-line immunomodulatory agents have any significant impact on the course of the disease.

A Asthma

The chest X-ray findings here are consistent with early pulmonary fibrosis related to asbestos exposure, and FEV_1/FVC is 87%.

B Solderer’s lung

Solderer’s lung is a cause of occupational asthma but does not result in the pleural plaques seen on chest X-ray here, nor the reticulo-nodular pulmonary shadowing.

D Chronic obstructive pulmonary disease (COPD)

COPD leads to obstruction on pulmonary function testing and is not associated with the chest X-ray changes seen here.

E Idiopathic pulmonary fibrosis

Idiopathic pulmonary fibrosis is a diagnosis of exclusion, and this patient’s job working with a plumber is associated with a high risk of asbestos exposure. The pleural plaques on chest X-ray do confirm previous exposure to asbestos fibres.

21028

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 68-year-old man with a 10-year history of chronic obstructive pulmonary disease (COPD) presents with a 6-month history of increasing exertional dyspnoea and fatigue. He also has a cough productive of yellow sputum, episodes of chest pain, nausea and anorexia. He stopped smoking 2 years ago.

On examination, he had raised jugular venous pressure (JVP), right parasternal heave, a loud second heart sound and an ejection systolic murmur at the left second intercostal space. On chest auscultation, he had scattered expiratory wheeze and bi-basal crepitations. He had pitting leg oedema.

Investigations reveal the following:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	180 g/l	135–175 g/l
White cell count (WCC)	8.0 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	220 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	135 mmol/l	135–145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	110 μmol/l	50–120 μmol/l
Bilirubin	16 μmol/l	2–17 μmol/l
Urea	4.6 mmol/l	2.5–6.5 mmol/l
Alkaline phosphatase (ALP)	50 IU/l	30–130 IU/l
Gamma-glutamyltransferase (**γ**GT)	100 IU/l	5–30 IU/l
Electrocardiogram (ECG)	Right ventricular hypertrophy, peaked P waves	
Chest X-ray (CXR)	Shown below	



Which of the following therapies is most likely to improve long-term prognosis?

Your answer was correct

A	High-dose furosemide
B	Long-term oxygen therapy
C	Angiotensin-converting enzyme (ACE) inhibitor
D	Regular venesection
E	Daily amoxicillin

Explanation

B Long-term oxygen therapy

This man has developed cor pulmonale, which is defined as right heart failure secondary to pulmonary hypertension. Causes include any chronic lung disease, pulmonary vascular disease, hypoventilation neuromuscular disorders and thoracic cage abnormalities. Long-term home oxygen therapy for at least 15 hours per day can reduce hypoxia, which lowers pulmonary resistance and can improve symptoms and increase life expectancy.

A High-dose furosemide

Diuretics can be used for right ventricular failure, but care should be taken to avoid excessive fluid depletion. In heart failure, these do not improve long-term prognosis.

C Angiotensin-converting enzyme (ACE) inhibitor

In contrast to left ventricular impairment, ACE inhibitors do not have any prognostic benefit and may, in fact, make matters worse. In systolic heart failure, they are of prognostic benefit and very useful.

D Regular venesection

This patient has polycythaemia secondary to chronic hypoxia as a mechanism of compensation. Venesection improves prognosis in haemochromatosis and can also be used for primary polycythaemia.

E Daily amoxicillin

There is no role for antibiotic prophylaxis in either cor pulmonale or COPD.

7713

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

An HIV-positive patient attends A+E with a history of a dry cough, shortness of breath and fever. On investigation, his CD4 count is 50 cells/mm^3 and his oxygen levels drop after exercise. On further questioning, he admits that he has not been taking his medications recently.

Investigations reveal the following:

Investigation	Result	Normal Value
Haemoglobin (Hb)	145 g/l	135–175 g/l
CD4	50/mm^3	500–1600/mm^3
Platelets (PLT)	210 × 10^9/l	150–400 × 10^9/l
pO2 on room air	9.5 kPa	

His X-ray is shown below



Which of the following is the first-line treatment for this patient?

Your answer was correct

A Prednisolone

B Co-trimoxazole

C Aciclovir

D Rifampicin, Isoniazid, Ethambutol and Pyrizinamide quadruple therapy

E Ganciclovir

Explanation

B Co-trimoxazole

Clinically he is infected with *Pneumocystis jirovecii*, which often presents with a dry cough in HIV-positive patients and other immunosuppressed patients, particularly after organ transplantation and prolonged use of immunosuppressants, and when the CD4 count is very low as here. On examination there is often tachycardia and tachypnoea, and there may be signs of cyanosis; however, the chest may even reveal no abnormality on examination in some patients. The chest X-ray shows subtle fine reticular interstitial changes in keeping with the diagnosis. If there is significant infiltration then crackles will be found on auscultation. Diagnosis is from pulmonary secretions or lung biopsy, looking for organisms using immunofluorescence or methenamine silver staining. The first-line treatment is with co-trimoxazole, but prednisolone can be added if the oxygen level is < 9.3 kPa.

Image source: https://commons.wikimedia.org/w/index.php?curid=77414546

A Prednisolone

The first-line treatment is with co-trimoxazole but prednisolone can be added as adjuvant therapy.

C Aciclovir

This is used to treat herpes virus infections. Herpes simplex infection is associated with oral and genital mucous membrane ulceration. It can be associated with headache and neurological symptomatology in the case of simplex encephalitis/meningitis.

D Rifampicin, Isoniazid, Ethambutol and Pyrizinamide quadruple therapy

This is the treatment for tuberculosis. It is associated with cough, fever, weight loss and night sweats, which are often described as ‘drenching’ in severity.

E Ganciclovir

This is the treatment for CMV infections. CMV retinitis, leading to visual symptoms, is the most common manifestation of systemic CMV infection in HIV-positive patients. Other features may include GI upset, or changes in cognition/consciousness in the case of CMV encephalitis.

7164

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 22-year-old woman with cystic fibrosis attends the chest clinic with her partner. They want to start a family before her health deteriorates. She has had several admissions to hospital over the past few years, but currently she feels relatively well. She works at a local estate agent. During a recent admission to hospital with an exacerbation of her cystic fibrosis, a sputum culture revealed a heavy growth of *Pseudomonas*, sensitive to ceftazidime.

Investigations:

Investigation	Result	Normal value
Haemoglobin	124 g/l	115–155 g/l
White cell count (WCC)	11.2 × 10^9/l	4–11 × 10^9/l
Platelets	197 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine	110 μmol/l	50–120 μmol/l
FEV_1	72% of that predicted	

Which of the following is the correct advice when it comes to getting pregnant?

Your answer was correct

A She should be advised to never get pregnant

B After treating the acute infection she still has a very high risk of dying/premature delivery of her child

C After treating the acute infection she has a good chance of a successful pregnancy

D She is unlikely to conceive due to thickening of the cervical mucus

E The fact her periods started late indicates subfertility

Explanation

C After treating the acute infection she has a good chance of a successful pregnancy

Evidence is difficult to evaluate with respect to cystic fibrosis and successful pregnancy. Around 98% of men with CF will be infertile, but the effect of CF on fertility in women is not so clear-cut. The first successful pregnancy in a woman with cystic fibrosis occurred in 1960, and since then increasing number of children have been born to mothers with CF.

A She should be advised to never get pregnant

Pregnancy can be well tolerated in those with mild disease, but in patients whose lung function is poor there is increased likelihood of infant prematurity and death. There may also be loss of maternal lung function with accelerated decline following pregnancy.

B After treating the acute infection she still has a very high risk of dying/premature delivery of her child

Observational studies suggest that patients with an FEV1 <60% predicted have a much worse outcome both with respect to their own health status and the viability of the pregnancy, and that chronic *Pseudomonas* infection and multiple exacerbations do much worse.

D She is unlikely to conceive due to thickening of the cervical mucus

Despite the fact that cervical mucus is thicker, this does not appear to impact significantly on ability to become pregnant. CFTR is found in large quantities on the cervix, which means that cervical mucus remains thick throughout the menstrual cycle.

E The fact her periods started late indicates subfertility

Even though menstruation tends to begin later in CF, this does not appear to impact significantly on ability to get pregnant.

21536

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Women with cystic fibrosis and their potential for reproduction
thorax.bmj.com/content/56/8/649
(http://thorax.bmj.com/content/56/8/649)

A 54-year-old man with chronic obstructive pulmonary disease (COPD) attends for smoking cessation advice.

Which of the following methods is likely to be associated with the greatest chance of quitting?

Your answer was incorrect

A Bupropion

B Nicotine replacement lozenges

C Willpower

D Nicotine replacement gum

E Hypnosis

Explanation

A Bupropion

Studies have shown success in smoking cessation in 20–30% of patients involved in bupropion studies at 1 year, with marginally higher quit rates seen when bupropion and nicotine replacement were combined (35%). Studies have shown that varenicline also aids smoking cessation in combination with motivational support, resulting in higher quit rates when higher doses were used (44% success rate).

C Willpower

Willpower quit rates are thought to be around only 2–4% at 1 year in reality.

B Nicotine replacement lozenges

Nicotine replacement is associated with a quit rate of only around 10% in clinical trials.

D Nicotine replacement gum

Like option B, nicotine replacement is associated with a quit rate of only around 10% in clinical trials.

E Hypnosis

Hypnosis aims to improve willpower in the subconscious state, and trials have showed no greater abstinence rate.

32442

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 68-year-old retired teacher presents with a 3-month history of recurrent haemoptysis. She had not sought medical advice at first as it had only been tiny, but the last few times there was a large amount of blood, which alarmed her. She had also noticed a persistent cough with some malaise and weight loss. She had a past medical history of tuberculosis when aged 19 years. She was an ex-smoker who stopped 15 years ago and did not drink alcohol. Physical examination was unremarkable.

The chest X-ray is shown below.



What is the most likely diagnosis?

Your answer was incorrect

A Apical lung carcinoma

B Aspergilloma

C Reactivation of pulmonary tuberculosis

D Lung abscess

E Mesothelioma

Explanation

B Aspergilloma

Image: *By Yale Rosen from USA (Aspergilloma X-rayUploaded by CFCF) [CC BY-SA 2.0 (https://creativecommons.org/licenses/by-sa/2.0)], via Wikimedia Commons*

This is a ball of *Aspergillus* fungus which forms inside an area of damaged lung tissue, most commonly secondary to TB but occasionally bronchiectasis, neoplasm or abscess cavity. The chest X-ray typically shows a round opacity in the upper zone, with a surrounding halo of air. It is usually asymptomatic, but may cause massive haemoptysis. Treatment is usually surgical excision of the affected lung segment. Other diseases caused by *Aspergillus fumigatus* are allergic bronchopulmonary aspergillosis and invasive aspergillosis.

C Reactivation of pulmonary tuberculosis

Reactivation of TB is a possible diagnosis, but the X-ray does not match with this diagnosis. TB would show a cavitating lesion or consolidation.

A Apical lung carcinoma

The appearance on the chest X-ray does not fit with a cancerous growth. Malignant growths in the lung are speculated on imaging, as the malignant cells follow lymphatic and vascular systems to grow from a central point.

D Lung abscess

The absence of recent infection and fever makes a lung abscess unlikely. An abscess would also have a denser appearance due to accumulation of fluid within it.

E Mesothelioma

Mesothelioma is rare without exposure to asbsestos, and when it does occur there are multiple plaques in the pleura, which are not present on this X-ray.

9153

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

This 66-year-old man presented with several months history of dyspnoea and chest pain which has not responded to inhalers or to medication for angina. He is a retired boilermaker, who smoked 40 cigarettes per day until 4 years ago. On examination his BP is 152/92, with pulse 80 and regular. There is wheeze on auscultation of the chest, and pitting oedema of both ankles.

His chest radiograph is shown below.



What is the most likely cause for the main abnormalities shown?

Your answer was correct

A Cigarette smoking

B Tuberculosis

C Asbestos exposure

D Silica exposure

E Previous trauma

Explanation

C Asbestos exposure

The chest radiograph shows a mesothelioma at the left apex which is invading the underlying rib. There are also calcified pleural plaques on the surface of both hemi-diaphragms. These appearances are consistent with asbestos exposure. Extensive occupational history is essential to establish any entitlement to compensation, as mesothelioma can develop up to 40 years after the initial exposure (often presenting with a unilateral pleural effusion). It is rarely suitable for surgical resection and is not responsive to chemotherapy. Palliative radiotherapy after thoracoscopic pleurectomy may be indicated. Only pemetrexed and cisplatin as chemotherapy has demonstrated a positive response. The prognosis is extremely poor, with an average survival of 1–2 years from diagnosis.

A Cigarette smoking

Cigarette smoking could not account for the mesothelioma seen in the left upper zone on CXR, nor the calcified pleural plaques.

B Tuberculosis

This leads to fibrotic changes and lymphadenopathy, but could not account for the left upper zone lesion or the calcified plaques seen here.

D Silica exposure

This leads to ‘eggshell’ calcification rather than pleural plaques, and is not associated with mesothelioma.

E Previous trauma

Previous trauma is more likely to be associated with evidence of old rib fractures and calcification, rather than the pleural changes seen here. It would also not account for the mesothelioma.

9151

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 45-year-old man presents to the Emergency Department with the sensation of something stuck in his throat. He reports to have been eating a steak dinner meal before a piece of meat became lodged when swallowing. On further questioning, he reports several episodes of dysphagia and has been told it is due to him not chewing properly. He occasionally takes over-the-counter antacids for heartburn but has no other past medical history of note.

He is referred for an oesophagogastroduodenoscopy (OGD) and the food bolus is removed. Images are captured of his oesophagus:



What is the most likely diagnosis?

Your answer was correct

A Achalasia

B Eosinophilic oesophagitis

C Gastro-oesophageal reflux disease

D Oesophageal adenocarcinoma

E Pharyngeal pouch

Explanation

B Eosinophilic oesophagitis

This patient likely has eosinophilic oesophagitis, given the history of repeated dysphagia and the presentation of an obstructed food bolus. Eosinophilic oesophagitis is an allergen-mediated condition characterised by oesophageal dysfunction including dyspepsia, dysphagia, vomiting and abdominal pain, along with eosinophilic infiltration in the oesophageal epithelium on biopsy. Oesophagogastroduodenoscopy (OGD) and biopsy are necessary for diagnosis, and findings include oesophageal oedema, oesophageal rings and strictures with white plaques or exudates representing eosinophilic micro-abscesses. The endoscopic image shown here shows multiple oesophageal rings, which supports the underlying diagnosis.

Image source: By Samir, CC BY-SA 3.0, https://commons.wikimedia.org/w/index.php?curid=5161861

A Achalasia

Achalasia is characterised by motor dysfunction of the oesophagus that causes symptoms of dysphagia and regurgitation due to loss of oesophageal peristalsis. Unlike oesophageal carcinoma, dysphagia for solids and liquids can present initially as the lower oesophageal sphincter fails to relax on swallowing. Nevertheless, OGD is indicated in the investigation of achalasia due to the concern of oesophageal carcinoma, and findings range from being normal to a dilated oesophagus with a tight lower oesophageal junction.

C Gastro-oesophageal reflux disease

Gastro-oesophageal reflux disease (GORD) typically presents with dyspepsia and regurgitation. It is associated with Barrett’s oesophagus, oesophageal strictures and oesophageal carcinoma. Typically, GORD can present similarly to eosinophilic oesophagitis and, in some cases, both conditions can be present with GORD worsening eosinophilic oesophagitis by worsening the oesophageal mucosal wall. OGD is an initial investigation to request with findings ranging from being normal to showing oesophagitis with erosions, ulcerations and strictures. The oesophageal rings on this endoscopy make eosinophilic oesophagitis the most likely diagnosis.

D Oesophageal adenocarcinoma

Oesophageal adenocarcinoma is an important differential diagnosis to consider, given the history of dysphagia. However, additional symptoms tend to include odynophagia, weight loss and a progressive dysphagia from solids to liquids. The findings of OGD include oesophageal obstruction due to an oesophageal mass.

E Pharyngeal pouch

A pharyngeal pouch is caused by a herniation between the thyropharyngeus and cricopharyngeus muscles that causes symptoms of dysphagia, regurgitation, chronic cough, aspiration and halitosis. It is most commonly seen in men aged above 70 years. Barium swallow is preferred over OGD due to the risk of perforating the pouch.

73484

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 62-year-old heavy smoker comes to the Emergency Department for review. He has suffered from progressively worsening headaches, shortness of breath and chest pain over the past three months. He has suffered from a dry cough, which is occasionally productive of bloodstained sputum, for about six months. A past history of chronic obstructive pulmonary disease (COPD) and ischaemic heart disease is noted.

On examination, his blood pressure is 155/95 mmHg, his pulse is 88 bpm and regular and he looks distressed. There is facial oedema and oedema of both arms, with a blue discoloration to the skin of the face. He has injected conjunctivae and optic discs are shown below Auscultation of the chest reveals bilateral wheeze and coarse crackles consistent with COPD, and decreased breath sounds on the right hand side.



Figure 1A Figure 1b

Investigations:

Investigations	Results	Normal Values
Haemoglobin (Hb)	135 g/l	135–175 g/l
White cell count (WCC)	8.9 × 10^9/l	4–11 × 10^9/l
Platelets (PLT)	201 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potasium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine	122 µmol/l	50–120 µmol/l
Glucose	5.4 mmol/l	3.5–5.5 mmol/l*

Chest X-ray: Large spiculated right hilar mass

Which of the following is the most effective and appropriate intervention in this situation?

Your answer was incorrect

A Chemotherapy

B Dexamethasone

C Endovascular stenting

D Radiotherapy

E Surgical reconstruction

Explanation

C Endovascular stenting

The oedema, facial discolouration and bilateral optic disc swelling, in a patient with an underlying tumour is indicative of superior vena cava (SVC) obstruction. In this situation, while corticosteroids are the mainstay of initial treatment, they are thought to be less effective when SVC obstruction is due to bronchial carcinoma, compared to SVC obstruction due to an underlying lymphoid malignancy. As such, where it is available, interventional radiology is the best intervention. Stenting also leads to a rapid improvement in symptoms, whereas with corticosteroids or radiotherapy, the improvement is gradual and modest.

Image source: https://commons.wikimedia.org/wiki/File:Behcet_fundus.JPG

D Radiotherapy

Radiotherapy is an alternative to stenting in the event that interventional radiology is not available or is not technically feasible. The disadvantage is that it can cause significant damage to the vascular anatomy, making future stenting technically more difficult.

A Chemotherapy

The response to chemotherapy in bronchial carcinoma is likely to be modest and relatively slow, compared to stenting. This makes it not a preferred option for symptom relief here.

B Dexamethasone

Steroids result in some reduction in local oedema and may impact on symptoms, although the effect is almost certainly going to be inferior to endovascular stenting. Where on-call interventional radiology is available, it is the intervention of choice, endorsed by the National Institute for Health and Care Excellence (NICE) guidance, for SVC obstruction.

E Surgical reconstruction

Surgical reconstruction would not likely be an option in this situation due to underlying advanced cancer, and the coexisting COPD makes him a very poor prospect for aggressive intervention.

39513

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 46-year-old woman presents to the Sleep Clinic with complaints of daytime sleepiness, irritability and morning headaches that have been getting worse over the past six months. She is accompanied by her partner who states that she snores during the night and he has observed a few episodes of apnoea occurring each night that occasionally awakens her up from sleep. She smokes five cigarettes per day.

On examination, her BP is 154/92 mmHg and her heart rate is 90 bpm. Her body mass index is 34 kg/m^2. A blood gas analysis is done at the clinic and the results are shown below. Chest auscultation revealed an occasional wheeze.

Investigation	Result	Normal value
pH	7.34	7.35–7.45
$p_a(O_2)$	10.9 kPa	11–14.4 kPa
$p_a(CO_2)$	6.3 kPa	4.6–6 kPa
HCO_3^-	27 mmol/l	22–26 mmol/l



Which of the following is the most likely diagnosis?

Your answer was incorrect

A Central sleep apnoea

B Chronic obstructive pulmonary disease

C Interstitial lung disease

D Obstructive sleep apnoea

E Obesity hypoventilation syndrome

Explanation

E Obesity hypoventilation syndrome

The diagnosis is obesity hypoventilation syndrome. The diagnostic criteria for obesity hypoventilation syndrome are:

- Body mass index (BMI) > 30 kg/m^2
- daytime $p_a(CO_2) > 6$ kPa
- absence of other causes of hypoventilation, eg opioid overdose.

Since this patient fits all of the above criteria, the diagnosis of obesity hypoventilation syndrome can be made. In addition, nocturnal oximetry can be done to identify coexisting sleep apnoea.

Image Source: Dr Yale Rosen, CC BY 2.0 via Wikimedia Commons

C Interstitial lung disease

Interstitial lung disease is not associated with hypercapnia and auscultation is likely to reveal bilateral crepitations.

A Central sleep apnoea

Central sleep apnoea is unlikely to be the diagnosis in this patient with daytime hypercapnia.

B Chronic obstructive pulmonary disease

In this patient with obesity and a normal chest X-ray, as well as with auscultation revealing only a mild wheeze, the diagnosis of obesity hypoventilation syndrome is much more likely than that of chronic obstructive pulmonary disease.

D Obstructive sleep apnoea

Obstructive sleep apnoea may coexist with obesity hypoventilation syndrome. However, nocturnal oximetry is required to demonstrate sleep apnoea.

72780

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 64-year-old woman comes to the Respiratory Clinic for review. She has recently been diagnosed with moderate COPD and is about to commence therapy for the condition. She also has a history of hypertension and chronic stable angina for which she receives ramipril, indapamide and diltiazem. On examination her BP is 132/82 mmHg, with pulse 70/min and regular. There is bilateral poor air entry with quiet wheeze on auscultation of the chest.

Which of the following is likely to have the greatest short-term impact on her quality of life, (in the next 2-3 months)?

Your answer was correct

A Ipratropium

B Long-term oxygen therapy (LTOT)

C Pulmonary rehabilitation

D Salbutamol

E Tiotropium

Explanation

C Pulmonary rehabilitation

Pulmonary rehabilitation has been shown to improve exercise tolerance in COPD, and will also improve this patient's exercise tolerance in relation to her chronic stable angina. Over a relatively short time, (about 6-8 weeks), these improvements have been shown to lead to quality of life (QoL) benefit.

A Ipratropium

Short-acting anti-cholinergics do not significantly impact on quality of life in COPD.

B Long-term oxygen therapy (LTOT)

LTOT reduces progression of pulmonary hypertension and right heart failure over the longer term. Although this translates into QoL benefit over the longer term, in the short term pulmonary rehabilitation will have a greater impact.

D Salbutamol

Short acting beta-agonists do not have significant impact on QoL in COPD.

E Tiotropium

Although tiotropium did show an impact on the St George’s respiratory quality of life score, the effect was moderate and was only seen in 3 out of 4 pivotal studies. Hence over the short term, pulmonary rehabilitation carries greater benefit.

70500

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 72-year-old man is referred to the Respiratory Outpatients Clinic with a 2-month history of increasing shortness of breath and cough. Over the last few weeks, he has coughed up a few streaks of blood. He has also noticed difficulty getting up from a chair, and that his clothes are looser recently without trying to lose weight. He is taking regular inhalers, but is unsure of their names.

On examination, he is clubbed. Auscultation of his chest reveals reduced expansion and breath sounds on the left. Heart sounds are normal. He has a 2 cm hepatomegaly on palpation of his abdomen. He has reduced power (4/5) on hip flexion bilaterally and generally reduced tendon reflexes. Plantars both show a flexor response; proprioception and sensation are normal.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	109 g/l	135 - 175 g/l
White cell count (WCC)	12.2×10^9/l	$4.0 - 11.0 \times 10^9$/l
Platelets (PLT)	491×10^9/l	$150 - 400 \times 10^9$/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	58 mm/h	1 - 20 mm/h
Sodium (Na^+)	126 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.6 mmol/l	3.5 - 5.0 mmol/l
Urea	5.9 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	125 μmol/l	50 - 120 μmol/l
Bilirubin	49 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	120 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	441 u/l	30 - 150 u/l
Albumin	32 g/l	35 - 55 g/l
Corrected calcium (Ca^{2+})	2.68 mmol/l	2.2 - 2.7 mmol/l
Phosphate (PO_4^{3-})	0.81 mmol/l	1.1 - 1.45 mmol/l

His chest X-ray is shown below:



What is the most likely underlying diagnosis?

Your answer was correct

A	Adenocarcinoma of the lung
B	Squamous carcinoma of the lung
C	Small-cell carcinoma of the lung
D	Mesothelioma
E	Bronchioalveolar carcinoma

Explanation

C Small-cell carcinoma of the lung

The chest X-ray shows left lower lobe collapse. There is loss of volume of the left hemi-thorax and the classical 'sail sign' behind the heart.

In addition to this suggestive CXR, this man also has clinical features suggestive of a lung malignancy weight loss, haemoptysis, dyspnoea and hypercalcaemia.

His investigations and clinical examination are suggestive of a paraneoplastic component to this malignancy:

- hyponatraemia secondary to the syndrome of inappropriate antidiuretic hormone secretion

- the proximal weakness and hyporeflexia, which, along with autonomic features, suggest Lambert-Eaton myasthenic syndrome.

These are seen more frequently with small cell lung cancer, making this the most likely diagnosis in this case. Ectopic ACTH production is also associated but rare.

Squamous cell carcinoma is associated with:

- gynaecomastia
- hyperthyroidism
- hypertropic pulmonary osteoarthropathy (HPOA) finger-clubbing, periostitis and arthritis
- hypercalcaemia usually secondary to secretion of PTH-related peptide.

Common to both:

- Clubbing
- Smoking: he is likely to be a smoker or ex-smoker and have COPD as he is on inhalers.

A Adenocarcinoma of the lung

Suspected lung malignancy with hyponatraemia and Lambert-Eaton myasthenic syndrome would be more compatible with small cell lung cancer than adenocarcinoma.

B Squamous carcinoma of the lung

Suspected lung malignancy with hyponatraemia and Lambert-Eaton myasthenic syndrome would be more compatible with small cell lung cancer than squamous cell carcinoma. Although the calcium is elevated, this minor elevation could well be consistent with dehydration rather than PTHrp

D Mesothelioma

Mesothelioma classically presents as progressive pleural thickening ± pleural effusion. This is not seen here, and there are no pleural or diaphragmatic calcified plaques, or features in the history, to suggest previous asbestos exposure.

E Bronchioalveolar carcinoma

This is a subtype of adenocarcinoma but with different characteristics and management. It presents more commonly in non-smokers, and has no gender predominance (unlike other forms of lung cancer which show a male predominance). Patients may present asymptomatically in up to 50% of cases. The paraneoplastic features make small cell lung cancer more likely than bronchioalveolar carcinoma in this case.

70133

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 62-year-old woman with a history of 30 pack-years of smoking presents with a cough and haemoptysis. She has been feeling increasingly lethargic over the past few months. On examination, there is no lymphadenopathy and she looks thin. Auscultation of the chest reveals wheeze and occasional coarse crackles consistent with a COPD picture.

Investigations reveal the following:

Investigation	**Result**	**Normal values**
Haemoglobin (Hb)	115 g/l	115–155 g/l
White cell count (WCC)	8.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	224 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	134 mmol/l	135–145 mmol/l
Potassium (K^+)	5.0 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	100 μmol/l	50–120 μmol/l
Chest X-ray (CXR)	Right upper lobe tumour	
FEV_1	2.0 l	
Staging CT	3.5 cm lesion with ipsilateral hilar lymphadenopathy; PET uptake consistent with ipsilateral lymph node involvement only	
Calcium level	3 mmol/L	2 2.6 mmol/L

Which of the following represents the likely appropriate management?

Your answer was incorrect

A Palliative chemotherapy

B Palliative radiotherapy

C Radical radiotherapy

D Surgery

E Radical chemoradiotherapy

Explanation

D Surgery

This patient is having changes suggestive of lung cancer with hypercalcemia which point towards squamous cell cancer of the lung. The N1 staging seen here, with ipsilateral hilar lymphadenopathy, is not a contraindication to surgery, with the PET scanning reassuring with respect to the possibility of metastases. This woman's lung function also suggests that she would be a potential surgical candidate. Guidelines suggest that $\text{FEV}_1 > 2.0$ l is the cut-off for pneumonectomy. N2 lymph nodes (ipsilateral mediastinal or sub-carinal involvement), particularly when they are bulky, do contraindicate surgery.

A Palliative chemotherapy

Palliative chemotherapy for non-small cell lung cancer is thought to at best extend life by around 2 months. Given this patient's young age, more aggressive intervention is warranted.

B Palliative radiotherapy

Palliative radiotherapy is considered for stage III and IV non-small cell lung cancer where surgical resection isn't an option.

C Radical radiotherapy

Radical radiotherapy is considered for patients with potentially curable lung cancer who are not fit to undergo surgical resection, for example on grounds of poor lung function or severe cardiovascular disease.

E Radical chemoradiotherapy

Here, with a curable lesion, surgery is the most appropriate intervention. With regards to immunotherapy in lung cancer, PD-1 and PDL-1 inhibitors, which help T cells to recognise cancer cells, are gaining ground in the treatment of the condition.

If she did have more advanced lymph node involvement and was not a surgical candidate, then combined chemo- and radiotherapy would be the most appropriate option.

21215

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

You are asked to review a 71-year-old man on the Respiratory Ward where he was admitted with a lower respiratory tract infection some five days earlier. Sodium levels and osmolalities on admission were consistent with syndrome of inappropriate antidiuretic hormone (SIADH). The nurses are concerned that, despite fluid restriction over the past 72 hours, his sodium level is still low at 125 mmol/l and he is drowsy and confused. Examination reveals a blood pressure of 143/85 mmHg and a pulse of 80 bpm and regular. He has chest signs consistent with chronic obstructive pulmonary disease (COPD). The abdomen is soft and non-tender, and there are no signs of fluid overload. He is drowsy and believes he is being held against his will in a prison hospital.

Investigations:

Investigations	Results	Normal Values
Haemoglobin (Hb)	131 g/l	135–175 g/l
White cell count (WCC)	7.1 × 10^9/l	4–11 × 10^9/l
Platelets (PLT)	181 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	125 mmol/l	135–145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine	100 μmol/l	50–120 μmol/l
C-reactive protein (CRP)	85 mg/l (down from 190 on admission)	< 5 mg/l

Which of the following is the most appropriate next step?

Your answer was incorrect

A	Continued fluid restriction only
B	Demeclocycline
C	Furosemide
D	Intravenous (IV) 0.9% saline
E	IV 1.8% saline

Explanation

B Demeclocycline

Given that this man's fluid restriction will have been well observed by nursing staff and, if anything, his fluid consumption is likely to have been reduced further by his drowsiness and confusion, continuing it is unlikely to drive rapid enough recovery of his serum sodium levels. Demeclocycline is the preferred option because it leads to partial nephrogenic diabetes insipidus and therefore drives the correction of serum sodium levels more rapidly than fluid restriction alone.

C Furosemide

Furosemide may be of value but only really drives improvement if urine osmolality is over twice plasma osmolality. It may also drive excess volume depletion and is therefore not the preferred option here.

A Continued fluid restriction only

Given this patient has already had a number of days of fluid restriction and his sodium level is still low at 125 mmol/l, further action is required, with demeclocycline as the preferred intervention here.

D Intravenous (IV) 0.9% saline

Normal saline (0.9%) may actually worsen hyponatraemia when urine osmolality exceeds plasma osmolality and so should be avoided here.

E IV 1.8% saline

IV saline solutions of 1.8% and 3% should only be considered in the event of life-threatening hyponatraemia (for instance, when there is significant neurological depression and/or seizures). Concentrated saline must be given in a High Dependence setting. In severe hyponatraemia, correction of > 8–10 mmol/day risks central pontine myelinolysis.

38377

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

An 18-year-old patient of short stature presents with fevers, thick nasal mucus and frontal headache. He is found to have bilateral crackles and clubbing of his fingers. He also gives a past history of chronic diarrhoea and has long-term problems with maintaining his weight.

Which of the following results is most closely related to the underlying diagnosis?

Your answer was incorrect

A Hypercalcaemia

B Eosinophilia

C A high number of macrophages on bronchoalveolar lavage

D A forced expiratory volume in 1 second (FEV_1) that improves > 12% following administration of salbutamol

E Inadequate pancreatic exocrine function

Explanation

E Inadequate pancreatic exocrine function

Cystic fibrosis gene mutation leads to defective chloride secretion and increased sodium absorption across the airway epithelium, which can result in bronchiectasis and deficient pancreatic exocrine function.

C A high number of macrophages on bronchoalveolar lavage

A high number of macrophages are often seen in smokers' lungs and are considered to be a non-specific finding.

A Hypercalcaemia

The constellation of clinical signs is in keeping with a diagnosis of cystic fibrosis. A raised calcium level could be associated with a number of conditions (such as sarcoidosis, hyperparathyroidism, Paget's disease and bone metastases), none of which would be consistent with the above findings.

B Eosinophilia

Multiple conditions may lead to a state of eosinophilia. Common causes include allergic conditions such as hay fever and asthma, as well as hypersensitivity due to medications. Eosinophilia may also be indicative of allergic bronchopulmonary aspergillosis (ABPA), which would, however, not lead to clubbing.

D A forced expiratory volume in 1 second (FEV_1) that improves > 12% following administration of salbutamol

A forced expiratory volume in 1 second (FEV_1) that improves > 12% following administration of salbutamol (Option D) is incorrect.

On spirometry, reversible FEV_1 values following the administration of bronchodilators (such as salbutamol) are typical of a diagnosis of asthma.

2331

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 28-year-old woman comes to the Respiratory Clinic for review. She has been suffering from asthma, confirmed by her general practitioner (GP) on peak flow testing, for the past four years. She has progressively increased her therapy to cope with symptoms of wheeze, shortness of breath on exercise and nocturnal coughing, but she is still not controlled. A past history of paroxysmal atrial fibrillation (AF) is noted, although she is currently not using any medication for prophylaxis. Current asthma medication includes inhaled beclometasone 400 mg twice daily and montelukast 10 mg orally (she originally responded well to the introduction of the oral therapy). On examination, her blood pressure is 115/85 mmHg and pulse 75 bpm and regular. There is bilateral wheeze on auscultation. Peak flow is 390 litres/minute (500 litres/minute predicted).

Which of the following is the most appropriate next step?

Your answer was incorrect

A Inhaled Tiotropium

B Inhaled salmeterol

C Oral Prednisolone

D Oral Sodium chromoglicate

E Oral Theophylline

Explanation

B Inhaled salmeterol

The National Institute for Health and Care Excellence (NICE) asthma guidelines recommend montelukast at step 2 (as here), and in the event patients are not controlled, a long-acting beta agonist (LABA), such as salmeterol, is a logical next step. Where patients do not see an improvement initially with montelukast, a switch to salmeterol should be accelerated.

C Oral Prednisolone

In this case, with multiple other potential interventions for asthma, moving to oral corticosteroids would not be considered positive from the point of view of benefit–risk. LABA, theophylline and anti-immunoglobulin E (IgE) monoclonal antibodies are all potential options before reaching for prednisolone.

A Inhaled Tiotropium

Tiotropium is a long-acting muscarinic antagonist (LAMA), used in the treatment of chronic obstructive pulmonary disease. Both dual therapy with LABA/LAMA and high-dose inhaled corticosteroids and LABA are endorsed by current guidelines.

D Oral Sodium chromoglicate

Sodium cromoglicate appears to be most effective in children with asthma and in those with predominant exercise-induced symptoms.

E Oral Theophylline

Assuming this patient’s inhaler technique is adequate, there is little added value in oral beta agonist therapy and it may increase the risk for AF in a patient with previous episodes of paroxysmal AF.

39176

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Asthma: diagnosis, monitoring and chronic asthma management
nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment
(https://www.nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment)

Asthma

Media

Expanded explanations

Flashcards

A 36-year-old man presents with increasing shortness of breath and a cough, which is worse at night. He has no significant past medical history. He has never smoked. He works as a car mechanic, restoring run-down cars. He notes that his symptoms are often better at the weekend.



Investigations:

FEV_1	70% of normal
FVC	85% of normal
PEFR	450 l/min
Histamine challenge test	20% fall in FEV_1 with 2 micromol/l histamine (normal >4)

What is the most likely diagnosis?

Your answer was incorrect

A Carbon monoxide poisoning

B Occupational asthma

C Silicosis

D Extrinsic Allergic Alveolitis

E Hyperactive airways

Explanation

B Occupational asthma

This man has occupational asthma secondary to isocyanates, which are found in paints used to spray cars. Occupational asthma is characterised by airway obstruction and/or bronchial hyper-reactivity induced by a person's occupation. His spirometry reveals an obstructive picture with a hyper-responsive histamine challenge test**.** Diagnosis is confirmed by asking the patient to do 2 h peak flows both at his place of work and at home, and ideally during a period of leave from work, to look for significant differences and worsening of PEFR at work.

Other occupations commonly associated with occupational asthma that may come up in the MRCP exam include: bakers, vets, hairdressers, pharmaceutical workers, welders, roofers and textile workers.

Image source: https://commons.wikimedia.org/wiki/File:Normal_AP.JPG

C Silicosis

Silicosis is associated with occupations such as foundry work, hard rock mining and civil engineering, and is due to inhalation of silicon dioxide. The history does not suggest this and the chest x-ray usually shows upper lobe fibrosis, unlike the normal film seen here.

A Carbon monoxide poisoning

The chest x-ray in carbon monoxide poisoning is usually normal, as here, however imaging can sometimes show a ground-glass appearance and alveolar oedema. It would usually present as a flu-like illness with malaise and fatigue, unlike in this case. There can be dyspnoea on exertion and chest pain with palpitations can occur. As exposure progresses, lethargy, hallucinations, agitation and confusion occur, eventually leading to coma. Patients may also experience headache, abdominal pain with nausea, vomiting and diarrhoea.

D Extrinsic Allergic Alveolitis

There is no history of allergen exposure to suggest EAA, which usually presents as a flu-like illness with dyspnoea soon after exposure to the offending allergen. The chest x-ray in EAA can be normal, especially in early disease, with later disease showing fibrosis, typically of the upper lobes, volume loss and cardiomegaly.

E Hyperactive airways

The patient does have hyperactive airways, but the 20% drop in FEV_1 on histamine challenge confirms a diagnosis of asthma as the underlying cause.

70144

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

You admit a 73-year-old man to the Emergency Department. He has had influenza over the past few days and has a worsening cough productive of purulent, blood-stained sputum. Observations: BP 122/72 mmHg, pulse 90/min and regular, respiratory rate 28/min. There are signs of right lower lobe pneumonia. A routine blood screen including arterial blood gas is performed.

Which one of the following features on admission is likely to be associated with a poorer prognosis when symptoms, signs and investigations are reviewed?

Your answer was incorrect

A Age

B Respiratory rate 28/min

C Serum urea 6 mmol/l

D Albumin 37 g/l

E ALT slightly outside the normal range

Explanation

A Age

Age >65 years is associated with a poorer prognosis in acute pneumonia. Other features not present here and which are associated with poorer prognosis include:

- confusion;
- urea >7 mmol/l;
- respiratory rate >30/min;
- blood pressure <90 systolic or <60 diastolic.

These features plus age make up the CURB-65 criteria. Where all five features are present, mortality from pneumonia is >50%, whereas 0-1 features carries a mortality rate of <3%.

C Serum urea 6 mmol/l

Normal serum urea is 2.5-7 mmol/l. This is a normal result and not of clinical significance.

B Respiratory rate 28/min

The normal respiratory rate is 12-20 breaths per minute. It is likely to be elevated in the context of pneumonia, due both to the lung pathology and systemic effects of the infection.

D Albumin 37 g/l

Normal serum albumin levels are >35-50 g/l. This is therefore normal and not of prognostic significance.

E ALT slightly outside the normal range

Deranged liver function tests tend to occur in atypical pneumonias such *Legionella* or *Mycoplasma*. Elevated ALT can indicated a poor prognosis, though a mild elevation is unlikely to be of clinical significance.

32371

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BTS management of CAP
brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-mana...

(https://www.brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/)

A 64-year-old man presents to the Emergency Department with nausea, vomiting and polyuria over the past few days. By the time he reaches the hospital he is drowsy and confused. Past medical history of note includes COPD, for which he takes high-dose seretide, a previous anterior myocardial infarction (MI), and he continues to smoke 30 cigarettes per day. On examination his BP is 115/75 mmHg, with pulse 90/min and regular. There is bilateral poor air entry on auscultation of the chest, with quiet wheeze. Abdomen is soft and non-tender.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	131 g/l	135 - 175 g/l
White cell count (WCC)	9.2 × 10^9 /l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	203 × 10^9 /l	150 - 400 × 10^9/l
Sodium (Na^+)	145 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.1 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	151 µmol/l	50 - 120 µmol/l
Glucose	6.9 mmol/l	3.9 - 7.1 mmol/l
Corrected calcium (Ca^{2+})	3.31 mmol/l	2.2 - 2.7 mmol/l
Albumin	38 g/l	35 - 55 g/l
Alanine aminotransferase (ALT)	122 u/l	7 - 55 u/l
Alkaline phosphatase (ALP)	191 u/l	30 - 150 u/l
Phosphate (PO_4^{3-})	0.7 mmol/l	1.1 - 1.45 mmol/l

Chest x-ray shown below.



Which of the following is the most likely diagnosis?

Your answer was incorrect

A	Adenocarcinoma of the bronchus
B	Bronchial carcinoid
C	Non-Hodgkin’s lymphoma
D	Small cell carcinoma of the bronchus
E	Squamous cell carcinoma of the bronchus

Explanation

E Squamous cell carcinoma of the bronchus

The chest x-ray reveals a peripheral mass in the left upper lobe. The most likely diagnosis in this case with hypercalcaemia and phosphate below the lower limit of the normal range, is squamous cell carcinoma of the bronchus associated with production of parathyroid hormone-related protein (PTHrP). If intact PTH is measured, then this is found to be below the lower limit of normal.

Image source: https://radiopaedia.org/cases/cavitating-lung-cancer?lang=gb

B Bronchial carcinoid

Bronchial carcinoid generally occurs at a younger age and is not associated with smoking; lesions are often peripheral. It is most likely to be associated with secretion of serotonin, although tumours secreting ACTH and growth hormone are also seen.

A Adenocarcinoma of the bronchus

Bronchial adenocarcinoma is rarely associated with paraneoplastic syndromes, although some cases of polymyositis are documented.

C Non-Hodgkin’s lymphoma

In NHL more extensive lymphadenopathy would be expected, and although PTHrP secretion is seen in NHL, it is much less frequently seen than in bronchial carcinoma.

D Small cell carcinoma of the bronchus

This is more likely to be associated with syndrome of inappropriate ADH secretion (SIADH) and hyponatraemia.

70499

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

This 72-year-old woman presents with several months' history of dyspnoea, cough and weight loss. There is a history of COPD for which she takes seretide BD, and coarse crackles and wheeze consistent with the diagnosis are evident on auscultation.

Her Postero-anterior (PA) and lateral chest radiographs are shown below:





Which of the following investigations is most appropriate to confirm the diagnosis?

Your answer was incorrect

A Fibre-optic bronchoscopy ± transbronchial biopsy

B CT guided fine-needle aspiration biopsy

C Respiratory function tests

D Sputum for cytology

E Mediastinoscopy

Explanation

A Fibre-optic bronchoscopy ± transbronchial biopsy

The combination of the history with a solitary lung nodule on chest X-ray is highly suggestive of bronchial carcinoma. Bronchoscopy enables the tumour to be directly visualised and samples taken for cytology and histology. The extent of carinal involvement can also be seen – if the tumour involves the first 2 cm of either main bronchus, it is deemed inoperable.

C Respiratory function tests

The combination of the history with a solitary lung nodule on chest X-ray is highly suggestive of bronchial carcinoma. Respiratory function tests are not useful in diagnosis, but are essential if pneumonectomy is being considered, as a forced expiratory volume (FEV_1) of <2.0 l means that there will in all likelihood be insufficient respiratory reserve post surgery.

B CT guided fine-needle aspiration biopsy

The combination of the history with a solitary lung nodule on chest X-ray is highly suggestive of bronchial carcinoma. CT-guided FNA carries more risk than bronchoscopy, and the lesion should be easily accessible via the bronchoscopy route. CT scanning of the thorax is however still useful in assessing involvement of the mediastinum and lymph nodes. If surgery is being considered, it should be extended to the abdomen and brain to exclude distant metastases.

D Sputum for cytology

The combination of the history with a solitary lung nodule on chest X-ray is highly suggestive of bronchial carcinoma. This option is incorrect, because sputum cytology is a low-yield diagnostic tool. It is unlikely to be able to provide enough malignancy cells from a sputum sample in order to make a histological diagnosis. In addition, the patient may not even be producing sputum. If an infective diagnosis is being considered, sputum induction could be carried out in order to obtain a sample.

E Mediastinoscopy

The combination of the history with a solitary lung nodule on chest X-ray is highly suggestive of bronchial carcinoma. Mediastinoscopy is an invasive procedure that involves making an incision in the anterior chest wall in order to introduce the endoscope. It is used to obtain mediastinal lymph node biopsies, for example, for staging lymph nodes involved in lung cancer or for the diagnosis of sarcoidosis or lymphoma. However, on reviewing the above chest radiographs, the mass should be easily accessible using the less invasive technique of bronchoscopy, so this should preclude mediastinoscopy.

32408

Rate this question:

Next Question

Previous Question Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 55-year-old woman presented with a 6-month history of recurrent chest infections. She had had a few episodes of wheeze and haemoptysis in the last few weeks. She also noticed intermittent hot flushes, which her GP had put down to the menopause. These could occur at any time but on occasions were precipitated by alcohol. She had no significant past medical history and she had never smoked.

On examination, she was anxious. Examination of her chest revealed reduced expansion of the upper right zone. Cardiovascular examination revealed a pansystolic murmur. Blood results were unremarkable.

Her chest X-ray is shown below:



What investigation is most likely to reveal the diagnosis?

Your answer was incorrect

A CT thorax

B Metacholine challenge test

C Urinary 5-hydroxyindoleacetic acid

D Urinary catecholamines

E Sputum culture

Explanation

C Urinary 5-hydroxyindoleacetic acid

This patient has bronchial carcinoid. It commonly presents between the ages of 50 and 70 and is unrelated to smoking. Bronchial carcinoid tumours are the most indolent form of a spectrum of neuroendocrine tumours of the lung that include small cell lung cancer as the most malignant. They are thought to arise from Kulchitsky cells in the bronchial mucosa. They usually secrete serotonin and arise from the large bronchus. Patients may present with a mass that is usually centrally located (as in this case – CXR shows right lower and mid lobe collapse), or with recurrent chest infections, haemoptysis, chest pain and wheeze. Flushing (which may be precipitated by alcohol, food ingestion, stress or emotion) and diarrhoea may occur. However, patients are commonly asymptomatic and the tumour is found incidentally.

Carcinoid syndrome occurs when there are secondary tumours in the liver that release serotonin into the systemic circulation. However, symptoms of carcinoid syndrome may rarely occur in bronchial carcinoid in the absence of metastases, as the bronchial tree drains into the systemic circulation.

Diagnosis is by detection of 5-hydroxyindoleacetic acid (the metabolite of serotonin) in the urine, and bronchial carcinoid is usually seen and diagnosed by bronchoscopy and biopsy.

The treatment of choice for bronchial carcinoid is surgical resection. Flushing may be controlled with somatostatin analogues.

Pellagra may occur due to tumour uptake of tryptophan, the precursor of nicotinic acid.

Image source: https://radiopaedia.org/cases/bronchus-intermedius-carcinoid?lang=gb (https://radiopaedia.org/cases/bronchus-intermedius-carcinoid?lang=gb)

B Metacholine challenge test

This is a test for asthma, which is an unlikely diagnosis based on the clinical features seen here.

A CT thorax

CT may confirm upper lobe collapse and help visualise any obstructing lesion, but would not give a definitive diagnosis.

D Urinary catecholamines

This is a test for phaeochromocytoma. Phaeochromocytoma is more likely to present with paroxysms of hypertension, severe headaches and dysglycaemia.

E Sputum culture

Sputum culture may reveal pulmonary infection due to an obstructed right upper lobe bronchus, but will not confirm the underlying diagnosis.

70102

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 39-year-old man presented to his general practitioner (GP) with a 3-month history of fever, rhinitis and weight loss. Two weeks ago he had an exacerbation of asthma and was treated with a course of erythromycin. He also complained of weakness in his hands and legs. He has a past medical history of angina. He is an ex-smoker and stopped five years ago.

On examination, he has a rash on his shins, shown below.



Auscultation of his chest reveals a mild expiratory wheeze. Examination was otherwise unremarkable. Urinalysis was normal.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	121 g/l	135 - 175 g/l
White cell count (WCC)	9.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	411 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	138 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	111 μmol/l	50 - 120 μmol/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
Neutrophils	5.3 × 10^9/l	1.5 - 7.0 × 10^9/l
Lymphocytes	1.8 × 10^9/l	1.5 - 4.0 × 10^9/l
Monocytes	0.3 × 10^9/l	0.0 - 0.8 × 10^9/l

Basophils	0.08 × 10^9/l	< 0.1 × 10^9/l
Eosinophils	1.9 × 10^9/l	0.04 - 0.4 × 10^9/l
Erythrocyte sedimentation rate (ESR)	82 mm/h	1 - 20 mm/h
Urea	4.1 mmol/l	2.5 - 6.5 mmol/l
Urine	Blood –, protein –	

Which investigation would help confirm the diagnosis?

Your answer was incorrect

A c-ANCA, proteinase 3 antibody positive

B c-ANCA, myeloperoxidase antibody positive

C p-ANCA, proteinase 3 antibody positive

D p-ANCA, myeloperoxidase antibody positive

E *Aspergillus* serology

Explanation

D p-ANCA, myeloperoxidase antibody positive

This patient has Eosinophilic granulomatosis with polyangiitis (EGPA) with systemic symptoms: asthma, eosinophils >1.5 × 10^9/l and evidence of vasculitis in two or more non-lung organs (Lanham's criteria):

- Cutaneous vasculitis (see picture)
- Mononeuritis multiplex
- A cardiac history.

The American Criteria of Rheumatology require four out of six of the following to be present for a diagnosis of EGPA:

- The presence of asthma
- Eosinophils >10% in the peripheral blood
- Evidence of a neuropathy in a vasculitic pattern, e.g. mononeuritis multiplex
- Transient pulmonary infiltrates
- A history of sinus disease
- Evidence of extravascular eosinophils on biopsy.

Renal involvement in EGPA is very rare.

C p-ANCA, proteinase 3 antibody positive

p-ANCA, proteinase 3 antibody positive is not usually seen as a combination of positive tests.

A c-ANCA, proteinase 3 antibody positive

c-ANCA, proteinase 3 antibody positive is associated with granulomatosis with polyangiitis

B c-ANCA, myeloperoxidase antibody positive

c-ANCA, myeloperoxidase antibody positive is not usually seen as a combination of positive tests, and c-ANCA positivity is inconsistent with a diagnosis of Eosinophilic granulomatosis with polyangiitis (EGPA), (where p-ANCA positivity would be more likely).

E *Aspergillus* serology

Aspergillus serology is used to diagnose *Aspergillus* lung disease, which would be inconsistent with the features of cutaneous vasculitis seen here.

70092

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Eosinophilic Granulomatosis with Polyangiitis

Expanded explanations

A 60-year old man presents with dull, right-sided chest ache, progressive shortness of breath and weight loss. He smokes 30 cigarettes per day and is a retired plumber with known asbestos exposure. On examination, BP is 125/72 mmHg with pulse 80/min and regular. There is reduced chest expansion on the right, with dullness to percussion and reduced air entry in the same area. Chest X-ray confirms the presence of a right-sided pleural effusion.

Which of the following is the optimal way to confirm the diagnosis?

Your answer was correct

A Blind pleural biopsy

B CT thorax

C Pleural fluid aspiration

D Thoracoscopy and biopsy

E Ultrasound guided biopsy

Explanation

D Thoracoscopy and biopsy

This patient most likely has mesothelioma, which can be difficult to diagnose. Biopsy is essential, but this needs to be done under imaging guidance to reduce the risk of tract spread, which is seeding of the tumour cells along the route of the biopsy needle. Thoracoscopy with biopsy under direct vision is the investigation of choice to confirm the diagnosis, and therefore this option is correct.

Mesothelial cells line the body cavities of the pleura, peritoneum, testis and pericardium. Malignancies involving these cells can be diffuse or localised. Most but not all pleural mesothelioma is associated with asbestos exposure. The most potent one is crocidolite (blue) followed by amosite (brown). There are three main histological types: mixed, sarcomatous and epithelial. The tumour arises from the visceral or parietal pleura and expands to encase the lung and spread to the oesophagus, superior vena cava and other nearby structures. The latent period from exposure is usually over 30 years. The higher the exposure dose the higher the risk. Some patients present up to 60 years after asbestos exposure. Where exposure is very high, presentation may occur as little as 10 years. Median survival after diagnosis is 11 months.

A Blind pleural biopsy

Although pleural biopsy is necessary for the diagnosis of mesothelioma, this method is relatively inaccurate at identifying the correct location for biopsy, and there is high risk of tract spread. CT-guided biopsy is better than blind biopsy with an Abraham's needle, since it increases the chance of biopsy of an affected area; however, it still carries the same risk of tract spread.

B CT thorax

CT is often used in the early stages of diagnosis of mesothelioma. On CT, mesothelioma appears as a nodular mass that spreads along pleural fissures. CT is a less sensitive test and would require subsequent biopsy to confirm the diagnosis.

C Pleural fluid aspiration

More than 90% of patients present with pleural effusion and although pleural fluid aspiration may be necessary, it is not frequently diagnostic. Pleural fluid tends to have low leukocyte levels, low LDH and elevated protein.

E Ultrasound guided biopsy

USS guidance is now recommended for all intrathoracic biopsies and fluid sampling. The superiority of thoracoscopy and biopsy under direct vision makes this option incorrect.

32377

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 31-year-old woman presents to the Emergency Department with abdominal pain and shortness of breath. She has had six attendances in the last six months, including gastritis, alcohol-related injuries and an upper gastrointestinal (GI) bleed, the latter from which she self-discharged from hospital one month ago.

She feels generally unwell, and on examination, her respiratory rate is 30 breaths/minute. Oxygen saturations are 81% on air, and she is apyrexial. She is cyanosed and has widespread fine inspiratory crackles on auscultation. Her abdomen is diffusely tender, with the pain moving to her back.

The results of her initial blood tests are as follows:

Haemoglobin (Hb)	103 g/l	(115–135 g/l)
White cell count (WCC)	14.1 × 10^9/l	(4–11 × 10^9/l)
Adjusted calcium	2.71 mmol/l	(2.20–2.60 mmol/l)
Alkaline phosphatase (ALP)	250 iu/l	(30–130 iu/l)
Alanine aminotransferase (ALT)	31 iu/l	(5–30 iu/l)

Her chest X-ray is shown below.



What is the most likely cause of this presentation?

Your answer was correct

A	Acute respiratory distress syndrome (ARDS)
B	Flash pulmonary oedema secondary to heart failure
C	Pulmonary haemorrhage
D	Pulmonary sepsis
E	Transfusion reaction

Explanation

A Acute respiratory distress syndrome (ARDS)

This patient has presented with a history of alcohol-related disease and abdominal pain. The nature of the pain, the blood results and the patient's risk factors all point to a diagnosis of pancreatitis. The chest X-ray shows diffuse bilateral coalescent opacities, which fits with **ARDS** , a known severe complication of pancreatitis. In

ARDS, pulmonary oedema is caused by lung damage of various potential sources, resulting in fluid leakage into the alveoli. Mortality may be as high as 75%, and early Intensive Therapy Unit (ITU) support is indicated.

B Flash pulmonary oedema secondary to heart failure

The X-ray demonstrates diffuse bilateral coalescent opacities, for which a reasonable differential diagnosis may be severe pulmonary oedema in the context of heart failure. However, the clinical stem is much more strongly suggestive of a pancreatitis picture, given the patient's age, lack of cardiac risk factors, blood results and history of alcohol excess.

C Pulmonary haemorrhage

Pulmonary haemorrhage may well be a reasonable differential for this X-ray. However, the major causes of pulmonary haemorrhage, such as vasculitides, are not suggested in the stem.

D Pulmonary sepsis

It would be important to consider sepsis in this patient, who has raised inflammatory markers and haemodynamic instability. However, the chest X-ray reveals more than just pneumonia, with ARDS as the most likely diagnosis.

E Transfusion reaction

This patient has had a previous upper GI bleed, and therefore, it is possible that they recently had a blood transfusion. However, while ARDS may be a manifestation of a transfusion reaction, this would tend to appear much more quickly than in this stem. ARDS can develop over 12–24 hours.

71770

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Faculty of Intensive Care Medicine. 2018. Guidelines on the management of ARDS.
ficm.ac.uk/news-events-education/news/guidelines-management-ards
(https://www.ficm.ac.uk/news-events-education/news/guidelines-management-ards)

Acute Respiratory Distress Syndrome

Expanded
explanations

A 19-year-old woman presents to the Emergency Department with shortness of breath and increasing purulent respiratory tract secretions. She has cystic fibrosis and takes nebulised tobramycin for *Pseudomonas* colonisation. On examination, she is pyrexial at 37.8 °C; there are bilateral coarse crackles affecting both bases, more marked on the right than on the left and wheeze throughout.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	110 g/l	135–175 g/l
White cell count (WCC)	12.3 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	181 × 10^9/l	400 × 10^9/l
C-reactive protein (CRP)	188 mg/l	0–10 mg/l
Chest X-ray (CXR)	Right basal consolidation	
Sputum microscopy, culture and sensitivity (MCS)	*Pseudomonas*species	

Which of the following is the most appropriate next step with respect to antibiotic therapy?

Your answer was incorrect

A One week IV co-amoxiclav

B One week IV gentamicin

C One week IV linezolid

D Two weeks oral ciprofloxacin

E Six months oral azithromycin

Explanation

E Six months oral azithromycin

Guidelines from the Cystic Fibrosis Trust recommend a prolonged course of oral azithromycin for patients who are chronically colonised with *Pseudomonas*. Two studies, one over three months and the other over six months, showed a positive impact on the forced expiratory volume in one second, exacerbations, body weight and quality of life.

C One week IV linezolid

Linezolid is an ineffective antibiotic against *Pseudomonas* infection. It is effective against Gram-positive bacteria such as *streptococci*, vancomycin-resistant enterococci and methicillin-resistant *Staphylococcus* aureus.

A One week IV co-amoxiclav

While useful for a broad spectrum of infections, including most respiratory pathogens, co-amoxiclav is not effective against *Pseudomonas* and is, therefore, not appropriate here.

B One week IV gentamicin

While gentamicin is effective against *Pseudomonas*, a prolonged course commonly results in adverse effects. A prolonged course of azithromycin has been shown to have more positive effects on respiratory function and quality of life in this population.

D Two weeks oral ciprofloxacin

Ciprofloxacin is incorrect. While it is effective against *Pseudomonas* infection, it is a less effective option, compared to azithromycin, and has a much higher risk of adverse effects, including musculoskeletal and neurological side effects.

40422

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Cystic Fibrosis Trust recommendations
cysticfibrosis.org.uk/sites/default/files/2020-11/Anitbiotic%20Treatment.pdf
(https://www.cysticfibrosis.org.uk/sites/default/files/2020-11/Anitbiotic%20Treatment.pdf)

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 24-year-old man with a known history of cystic fibrosis (CF) presents as a self-admission to the ward. He has increased volume of purulent sputum, more shortness of breath and left-sided chest ache. On reviewing his history, you note that he is under consideration for the transplant list and has been colonised with *Pseudomonas* for some years. On examination he is pyrexial, with a temperature of 38.2°C, and appears very unwell with FEV_1 (forced expiratory volume in 1 s) having fallen to <75% of the previous recorded value. He is hypotensive and tachycardic, with a respiratory rate of 32/min. Gram-negative rod-shaped bacilli are isolated from his sputum.

Which of the following micro-organisms would fit best with this pattern of deterioration?

Your answer was incorrect

A *Haemophilus influenzae*

B *Staphylococcus aureus*

C *Streptococcus pneumoniae*

D *Burkholderia cepacia*

E *Stenotrophomonas maltophilia*

Explanation

D *Burkholderia cepacia*

Cepacia syndrome is associated with a severe worsening of respiratory symptoms in a small number of patients with cystic fibrosis. Thankfully the majority of patients colonised with *B. cepacia* do not, however, suffer a deterioration in their underlying condition. Cepacia is known to demonstrate resistance to multiple antibiotics including colomycin and aminoglycosides, and hence intravenous regimes involving two or more antibiotics such as ceftazidime and an aminoglycoside should be employed. Other possible combinations include ceftazidime and chloramphenicol or chloramphenicol and minocycline. Approximately 2.5–3.0% of cystic fibrosis individuals are thought to be colonised with *B. cepacia*, and measures are usually taken to limit colonisation where possible.

C *Streptococcus pneumoniae*

S. pneumoniae is a Gram-positive coccus that does cause community-acquired pneumonia. However, the picture here with rapid deterioration and rod-shaped bacilli on sputum sampling is more consistent with *B. cepacia* syndrome.

A *Haemophilus influenzae*

H. influenzae colonisation is frequently seen in patients with CF, where they are visible as Gram-negative cocci in sputum samples. Colonisation may be associated with acute exacerbations of cystic fibrosis.

B *Staphylococcus aureus*

S. aureus is most likely to be associated with lower respiratory tract infection post-influenza. There is no apparent preceding episode of influenza reported here.

E *Stenotrophomonas maltophilia*

S. maltophilia colonises equipment used in mechanical ventilation and urinary catheters and is a rare cause of opportunistic infection in these settings.

32425

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 49-year-old male head teacher presents with a progressive shortness of breath on exertion for the last six months. He has also suffered from joint pains over the last few months, and an erythematous rash affecting sun-exposed areas during the summer months. He has previously been fit and has had no medical consultations for 25 years. He does not smoke. On examination his BP is 155/82 mmHg, pulse is 82/min and regular, BMI 22 and there are occasional crackles on auscultation of his chest. Full blood count and renal function are within normal limits, but autoantibodies are positive for anti-nuclear antibodies.
Lung function test results are as follows:

	Actual	Predicted
FEV_1 (L/min)	1.4	2.0 - 2.8
VC (L)	1.6	2.4 - 3.4
FEV_1/VC	88%	
TLCO (ml/min/mmHg)	2.8	3.5 - 7.3
KCO	0.9	1.2 - 1.8

Which of the following is the most likely cause of his abnormal lung function?

Your answer was incorrect

A Asthma

B Systemic lupus erythematosus (SLE)

C Pulmonary haemorrhage

D Idiopathic pulmonary fibrosis (IPF)

E Inherited thrombophilia with multiple pulmonary emboli

Explanation

B Systemic lupus erythematosus (SLE)

The results show reduced lung volumes with normal FEV_1/VC ratio and impaired gas transfer. The diagnosis is interstitial lung disease, of which the most common in this age group are idiopathic fibrosing alveolitis, and fibrosing alveolitis secondary to connective tissue disease, sarcoidosis and pneumoconiosis. Spirometry shows reduced lung volumes with FEV_1/FVC ratio >75%. Gas transfer is also impaired (low KCO). The history of polyarthritis, photosensitive rash and positive anti-nuclear antibodies fits best with a diagnosis of SLE.

C Pulmonary haemorrhage

A more acute deterioration would be expected in patients with pulmonary haemorrhage, and elevation in low carbon monoxide transfer factor (TLCO) due to intra-alveolar haemoglobin would be expected.

A Asthma

Asthma is associated with an obstructive lung defect rather than the restrictive picture seen here.

D Idiopathic pulmonary fibrosis (IPF)

IPF is inconsistent with the other features of connective tissue disease seen here.

E Inherited thrombophilia with multiple pulmonary emboli

Multiple thromboemboli may result in reduced gas transfer, but they wouldn't result in the restrictive picture on pulmonary function testing seen here. In addition, given the features of SLE seen here, if pulmonary emboli were detected, anti-cardiolipin antibodies would be a more likely cause.

32430

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 79-year-old man with a previous history of tuberculosis comes to the Emergency Department for review because of increased shortness of breath. He has a past history of tuberculosis (TB), for which he underwent a surgical procedure many years ago, and has hypertension and ischaemic heart disease, but is otherwise relatively well.

Examination reveals a blood pressure of 138/85 mmHg; pulse is 80 bpm and regular. He has reduced expansion on the right-hand side of his chest, and decreased breath sounds over the upper lobe.

His chest X-ray is shown below:



Which procedure has been performed?

Your answer was incorrect

A	Oleothorax
B	Pleurodesis
C	Plumbage
D	Upper lobectomy
E	Thoracoplasty

Explanation

E Thoracoplasty

Thoracoplasty involves removal of ribs to a collapsed area of lung. Given the chest X-ray appearances, this is the most likely procedure that has been performed here. It was largely discontinued after the advent of anti-microbial therapy for tuberculosis as well as the operative mortality rate being high.

D Upper lobectomy

A right upper lobectomy would result in an elevated right hemidiaphragm and displacement of the right hilum as opposed to the chest X-ray changes here.

A Oleothorax

Oleothorax involves instillation of paraffin into the pleural cavity and is used in pleurodesis. It does not fit with the clinical history above.

B Pleurodesis

Pleurodesis can be performed medically or surgically and is indicated for recurrent pneumothorax or pleural effusions, for example, caused by incurable cancer. Medical pleurodesis involves the insertion of an irritant, such as talc, into the pleural space to cause the pleural to adhere to the lung. There is no clinical indication in the vignette to suggest its use here.

C Plumbage

Plumbage involves instillation of lucite balls to permanently collapse the lung and was used as treatment option in patients with TB. However, the chest X-ray appearances are not consistent with this diagnosis.

39655

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 30-year-old man from London, with no significant past medical history, went on holiday to Peru to climb the Inca Trail. He is a non-smoker. After climbing above 3500 m, he developed a headache, associated with nausea and vomiting. He also felt tired and complained of insomnia. His respiratory rate was 22 breaths per minute and his heart rate was 90 bpm.

Investigations:

Investigation	Result	Normal Value
White cell count (WCC)	10.1×10^9/l	$4.0 - 11.0 \times 10^9$/l
Platelets (PLT)	223×10^9/l	$150 - 400 \times 10^9$/l
Urea	10 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	70 μmol/l	50 - 120 μmol/l
Sodium (Na^+)	135 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5 - 5.0 mmol/l

Which of these factors, if present, should indicate the need for immediate descent?

Your answer was incorrect

A Ataxia

B Dizziness

C Headache

D Insomnia

E Vomiting

Explanation

A Ataxia

There are three types of altitude illness: acute mountain sickness (AMS), high-altitude cerebral oedema (HACE) and high-altitude pulmonary oedema (HAPE). This man has AMS, with some evidence of cerebral oedema. The exact cause is unknown; however, it is thought to be secondary to cerebral oedema and raised intracranial pressure. It is more common in people who live at low altitudes.

Symptoms and signs of AMS are non-specific; however, diagnosis can be made using the criteria according to the ‘Lake Louise Score’, namely, the setting of the recent gain in altitude, the presence of headache and at least one of the following symptoms:

- gastrointestinal – anorexia, nausea and vomiting
- fatigue and weakness
- dizziness or light-headedness
- difficulty sleeping.

Acetazolamide may be used for the prevention and treatment of AMS. It is a carbonic anhydrase inhibitor that causes intracellular acidosis. It has a diuretic effect which can be useful in AMS/HACE and HAPE. It is also beneficial, as it shifts the oxygen dissociation curve to the right and therefore, at lower partial pressures, more oxygen is released to the tissues.

If the patient does not improve and progresses to developing ataxia and drowsiness, this is the development of HACE, which can be considered end-stage AMS. Treatment is immediate descent, oxygen and intravenous dexamethasone.

C Headache

Headache is part of the Lake Louise Score criteria for recognising mild disease due to high altitude. The presence of headache may indicate AMS. The approach, if this develops, should be to stop, rest, treat once diagnosed and descend whenever necessary. It is not an indication for immediate descent.

B Dizziness

Dizziness is another criterion in the Lake Louise Score for AMS and is not a trigger for immediate descent.

D Insomnia

Insomnia is also one of the criteria within the Lake Louise Score. Acetazolamide can be used as prophylaxis to prevent AMS.

E Vomiting

Vomiting is another of the Lake Louise Score criteria for AMS and is not a trigger for immediate descent. 72976

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Difficult

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

High Altitude Medicine

Expanded explanations

A 73-year-old man is sent to the Emergency Department by ambulance after a General Practitioner home visit. He says he has been feeling unwell with a 3-day history of shortness of breath and a cough productive of rusty-coloured sputum. His past medical history includes hypertension and hypercholesterolaemia.

On examination, he has a heart rate of 90 bpm and blood pressure of 101/59 mmHg. There is bronchial breathing on auscultation. His respiratory rate is 22 breaths per minute and oxygen saturations 94% on air. He has herpetic lesions at the corner of his mouth.

Investigations:

Investigation	Result	Normal value
Haemoglobin	159 g/l	135–175 g/l
White cell count (WCC)	15.5 × 10^9/l	4–11 × 10^9/l
Platelets	450 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135–145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5–5.0 mmol/l
Urea	7.4 mmol/l	2.5–6.5 mmol/l
Creatinine	145 µmol/l	50–120 µmol/l

Given this patient’s presentation, what is the recommended management plan?

Your answer was incorrect

A Admission to High Dependency Unit (HDU)

B Admit to Medical Ward for intravenous antibiotics

C Discharge home with aciclovir and early follow-up

D Discharge home with oral antibiotics

E Urgent Respiratory referral

Explanation

A Admission to High Dependency Unit (HDU)

The British Thoracic Society community-acquired pneumonia severity score (CURB-65) uses five variables, each of which scores one point – it is a mortality score.

The variables are:

- confusion (abbreviated mental test (AMT) score of ≤ 8)

- urea concentration > 7 mmol/l
- respiratory rate ≥ 30 breaths per minute
- systolic blood pressure ≤ 90 mmHg or diastolic blood pressure ≤ 60 mmHg
- age > 65.

A score of 3 is associated with a mortality risk of 17% from community-acquired pneumonia; with a score of 5, the mortality risk rises to 57%. These data are extrapolated to give a guide as to whether or where the patient should be admitted – those with a score of 0 or 1 may be managed at home safely; those with a score of 2 should receive hospital care, whereas management in the High Dependency Unit (HDU) or Intensive Therapy Unit should be considered for patients with a score of 3 or more. This is a guide and should not replace clinical judgement.

D Discharge home with oral antibiotics

This patient has signs and symptoms consistent with bacterial pneumonia (most likely *Streptococcus pneumoniae*). It is important to calculate the CURB-65 score in such patients to help guide the most appropriate management plan. This patient has a CURB-65 score of 3 and patients with a CURB-65 score of 3 or above should be considered for admission to the HDU, given the significantly increased mortality risk.

B Admit to Medical Ward for intravenous antibiotics

Admission to a Medical Ward is a reasonable consideration. However, given the CURB-65 score of 3, considerations should be made regarding admission to a level 2 place of care (eg HDU) for more cautious monitoring.

C Discharge home with aciclovir and early follow-up

It would be inappropriate to discharge this patient home on oral aciclovir, as this will not treat his bacterial pneumonia. A patient with a CURB-65 score of 3 or more has > 10% mortality risk.

E Urgent Respiratory referral

An urgent Respiratory referral is not an immediately necessary step in the management of this patient. Rather, admission for intravenous antibiotics and fluid resuscitation are key.

72944

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society (BTS): BTS Pleural Disease Guideline 2010 Management of spontaneous pneumothorax
thorax.bmj.com/content/65/Suppl_2/ii18
(https://thorax.bmj.com/content/65/Suppl_2/ii18)

Pneumonia

Media

Expanded explanations

Flashcards

A 27-year-old professional football player is admitted with a fracture of his right tibia, which he sustained during the course of a match. He is put on traction in the ward and he is scheduled for theatre two days later. On the eve of the procedure he is seen by the house officer because of an episode of chest pain. Clinical examination is unremarkable except petechial rash on chest. His electrocardiogram (ECG) and chest X-ray are reported as normal, and the patient is prescribed some analgesia.
The following morning the patient is taken to theatre as planned. While in the anaesthetic room he is noted to have a respiratory rate of 25/min and oxygen saturations of 90% on room air. His heart rate is 100 beats/min regular and his blood pressure is 90/65 mmHg. His heart sounds are normal and there are coarse crackles on auscultation of his chest bilaterally. His jugular venous pressure is raised to 5 cm. There is no peripheral oedema.

His ECG shows right axis deviation with prominent R waves on leads V1–V2.

Which one of the following measures would be the appropriate next step in this patient’s management?

Your answer was incorrect

A IV furosemide

B IV GTN infusion

C IV fluids

D Subcutaneous low-molecular weight heparin

E IV thrombolysis

Explanation

C IV fluids

This patient has symptoms and signs suggestive of acute pulmonary oedema which s extremely unlikely to be cardiac in origin in a young athlete. In this case the non cardiac pulmonary oedema is most likely caused by fat embolism syndrome. The syndrome classically presents with three main components: a respiratory picture indistinguishable from ARDS, neurological features (ranging from acute confusion, reduced consciousness to seizure and focal deficit) and a petechial rash. Usually chest signs appear first, and not all patients will have all three system features. Treatment is supportive; maintain circulatory pressure with IV fluids if needed, and to remove the underlying cause if possible.

D Subcutaneous low-molecular weight heparin

There is no evidence that heparin is of benefit in the management of fat embolism.

A IV furosemide

IV furosemide is not recommended in the acute situation because of the importance of maintaining circulatory pressure.

B IV GTN infusion

Like furosemide, GTN infusion may significantly reduce blood pressure with a consequent negative impact on outcomes.

E IV thrombolysis

Although thrombolysis is of benefit in massive pulmonary embolism, it is not of use here as the symptoms are related to ARDS, not to an embolus.

32276

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Acute Respiratory Distress Syndrome

Expanded
explanations

A 39-year old Irish accountant was referred to the clinic with a 4-month history of progressive breathlessness. He noticed this mainly on climbing the stairs. Over the last few months he had been using some eye drops from the local pharmacy for dry eyes. He was also complaining of fatigue, night sweats and polyuria. He was a non-smoker and drank 20 units of alcohol per week.

On examination he had conjunctival injection, but examination of his eyes was otherwise unremarkable. He had no rash or joint swelling. On auscultation of his chest he had fine inspiratory crackles. Heart sounds were normal. He had no organomegaly on palpation of the abdomen. Urinalysis was unremarkable.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	101 g/l	135 - 175 g/l
White cell count (WCC)	6.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	201 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	100 µmol/l	50 - 120 µmol/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	45 mm/h	1 - 20 mm/h
Urea	7.2 mmol/l	2.5 - 6.5 mmol/l
Bilirubin	23 µmol/l	1 - 22 µmol/l
Asparate aminotransferase (AST)	51 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	91 u/l	30 - 150 u/l
Albumin	45 g/l	35 - 55 g/l
Tuberculin test	-	
FEV1	75% predicted	
FVC	70% predicted	
TLCO	80% predicted	

His chest X-ray is shown below.



Which of the following is most likely to give you a definitive diagnosis?

Your answer was incorrect

A	Serum angiotensin-converting enzyme
B	Sputum for microbiology
C	Bronchoscopy and transbronchial biopsies
D	Bronchoscopy and endobronchial biopsies
E	Kveim test

Explanation

C Bronchoscopy and transbronchial biopsies

The most likely diagnosis is sarcoidosis. This is a relatively young patient with breathlessness, dry eyes, fatigue, night sweats and polyuria. He has a normocytic anaemia, a raised ESR and slightly deranged liver function tests.

Sarcoid is a multi-system granulomatous disorder of unknown aetiology with a number of different presentations. It usually presents in the under-forties, and is more common in women and persons of West Indian and African descent. Presentation may be with erythema nodosum, bilateral hilar lymphadenopathy on chest radiography with fever or arthralgia (Löfgren syndrome) or, more insidiously, with symptoms such as dyspnoea, dry cough, fever, malaise and weight loss. The main differential diagnoses are TB and lymphoma.

The diagnosis is suggested from a chest X-ray and high-resolution CT. High-resolution CT characteristically shows features such as hilar and mediastinal lymphadenopathy, peribronchovascular nodules, ground-glass shadowing, parenchymal bands, cysts and fibrosis. Pulmonary involvement can be classified according to the radiographic stage of the disease:

Stage 0: Clear chest radiograph
Stage 1: Bi-hilar adenopathy
Stage 2: Bi-hilar adenopathy and interstitial infiltrates
Stage 3: Diffuse interstitial disease
Stage 4: Advanced fibrosis.

Transbronchial biopsy is the investigation with the highest yield, as positive results are seen in 90% of patients with pulmonary sarcoidosis. Endobronchial biopsies are also useful, but less sensitive. Histology typically shows non-caseating granulomas composed of macrophages, lymphocytes and epithelioid histiocytes, which fuse to form multinucleate giant cells.

The lung function tests depend on severity of the disease. The tuberculin test is negative in 80% of patients with sarcoidosis, but clinicians must be mindful that anergy can also be seen in HIV-positive patients or patients who have overwhelming TB.

Treatment is rarely required in stage 0 or 1 disease. Corticosteroids should be given to patients with stage 2 or higher disease with deterioration in lung function or in patients with ocular sarcoid, hypercalcaemia, severe or persistent erythema nodosum, myocardial and neurological manifestations of sarcoidosis. Prednisolone dose is 0.5 mg/kg for first 4 weeks of treatment and is then down-titrated to the lowest maintenance dose that controls the disease. Treatment withdrawal should be considered after 6–24 months. Steroid-sparing agents such as methotrexate or azathioprine may be used for relapsing disease.

Common clinical characteristics of sarcoidosis include:

- Skin:
  Erythema nodosum – caution, not always on the shins
  Lupus pernio – red crusty lesions, often around the nose
  Annular lesion
- Polyuria:
  Secondary to hypercalciuria/hypercalcaemia (can also cause renal calculi and nephrocalcinosis)
  Secondary to central diabetes insipidus
- Cardiac:
  Ventricular arrhythmias
  Conduction defects
  Cardiomyopathy
  Congestive cardiac failure
- Neurological:
  Involvement of CNS occurs in 2%
  Cranial diabetes insipidus
- Ocular:
  Anterior uveitis
  Conjunctivitis
  Retinal lesions
  Keratoconjunctivitis sicca and lacrimal gland enlargement
  Optic neuritis

- Metabolic:
  Hypercalcaemia and hypercalciuria
- Bone and joints:
  Arthralgia
  Bone cysts

Others:
Hepatosplenomegaly
Löfgren syndrome (acute sarcodosis) – triad of bi-hilar lymphadenopathy, arthritis and erythema nodosum.

B Sputum for microbiology

This is not a history of infection. Sputum culture will be negative. Culture may be useful in the event that TB is suspected, although there is no history of potential TB exposure.

A Serum angiotensin-converting enzyme

Serum ACE is not a specific test for sarcoidosis. Raised ACE levels are also seen in pulmonary TB, asbestosis, silicosis and lymphoma. It is elevated in about 75% of patients with untreated sarcoid. The value of using serum ACE to monitor disease activity remains unclear – currently inflammatory markers are thought to be more useful.

D Bronchoscopy and endobronchial biopsies

Endobronchial biopsies may give a diagnosis of sarcoidosis, but transbronchial biopsies in addition to endobronchial biopsies will give a higher yield.

E Kveim test

The Kveim test is now obsolete because of potential infection risk. It involves using material from the spleen of a patient suffering from sarcoidosis.

70084

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 67-year-old non-smoker who was previously a plumber was referred to the Respiratory Consultant with a 3-month history of progressive shortness of breath. She was also complaining of right-sided chest pain, which she described as a constant severe ache. She had lost about 5 kg in the last month and felt lethargic. She had initially put the symptoms down to the stress of recently being widowed. Her husband, who also worked as a plumber, died of a 'lung problem'. She has two daughters who are fit and well.

On examination, she looked unwell and was breathless at rest. Examination of her chest revealed reduced expansion, percussion note, vocal fremitus and breath sounds throughout the right hemithorax.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	102 g/l	135 - 175 g/l
White cell count (WCC)	10.8 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	471 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	136 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.7 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	77 μmol/l	50 - 120 μmol/l
Mean corpuscular volume (MCV)	81 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	69 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	44 mg/l	< 10 mg/l
Urea	6.6 mmol/l	2.5 - 6.5 mmol/l
Bilirubin	34 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	21 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	97 u/l	30 - 150 u/l
Albumin	37 g/l	35 - 55 g/l

Her chest X-ray is shown below:



Which of the investigations is most likely to lead to a definitive diagnosis?

Your answer was incorrect

A	Video-assisted thoracoscopic surgery biopsy
B	CT chest
C	Chest drain and fluid sent for cytology
D	CT-guided percutaneous needle biopsy
E	Blind pleural biopsy with an Abraham’s needle

Explanation

A Video-assisted thoracoscopic surgery biopsy

Both this patient and her husband worked as plumbers for most of their adult life, and, therefore, there is a high possibility that they were both exposed to asbestos. She has systemic symptoms as well as a pleural effusion, making the most likely diagnosis malignant mesothelioma.

Video-assisted thoracoscopy surgery is the investigation of choice as it has the highest diagnostic yield. Prophylactic radiotherapy should be given to the operation site.

The chest X-ray shows pleural thickening and effusion on the right. There is a holly-leaf pleural plaque on the left (directly above the heart, to the left of the aortic knuckle), consistent with previous asbestos exposure. Incidentally, there is pericardial calcification.

C Chest drain and fluid sent for cytology

It is unclear on chest radiography alone whether this woman has a pleural effusion. The shadowing seen may just be due to pleural thickening. A thoracic ultrasound should be performed prior to any pleural fluid aspiration, and certainly before chest drain insertion for pleural effusion (as per BTS guidelines). Additionally, pleural fluid cytology is often non-diagnostic in mesothelioma.

B CT chest

A CT may show features in keeping with a high likelihood of mesothelioma, but histology is required for a definitive diagnosis.

D CT-guided percutaneous needle biopsy

In practice, radiologically guided pleural biopsies are used more in diagnosis of mesothelioma than VATS biopsy. However, the question asks which investigation is most likely to lead to a definitive diagnosis and the answer to this is VATS biopsy.

E Blind pleural biopsy with an Abraham’s needle

Blind pleural biopsy will give a lower diagnostic yield than CT-guided or VATS biopsy.

70097

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 46-year-old man presents with a persistent dry cough and shortness of breath. There has been a history of fatigue over the past few months and he has recently lost his job. On examination there are diffuse crackles on auscultation of the lung fields. He also has a nodular rash on his shins. There is a diffuse increase in globulins.

His chest x-ray is shown below.



Which of the following represents the best initial step for this patient?

Your answer was incorrect

A Oral prednisolone

B Oral methotrexate

C Chemotherapy for myeloma

D Bronchoscopy to look for mycobacteria

E Oral azathioprine

Explanation

A Oral prednisolone

This patient has sarcoidosis, a chronic granulomatous disorder of unknown origin, characterised histologically by the presence of non-caseating granulomas. Work-up involves chest X-ray and biopsy of easily accessible affected tissue. Here, the chest x-ray shows bilateral hilar enlargement and a lesion in the left lower zone. Corticosteroids, namely oral prednisolone, remain the initial mainstay of treatment, although they are reserved for patients with significant symptoms (in this case, cough and shortness of breath) or severe changes on CXR. Steroid-sparing agents such as methotrexate or hydroxychloroquine may be added if required. Most patients achieve remission within 2 years, though as yet there are no curative therapies and relapses are common.

Image source: https://radiopaedia.org/cases/sarcoidosis-57?lang=gb (https://radiopaedia.org/cases/sarcoidosis-57?lang=gb)

C Chemotherapy for myeloma

Myeloma tends to present in older patients, with symptoms such as bone pain, fatigue secondary to anaemia, recurrent infections, hypercalcaemia and renal symptoms. The persistent cough and CXR findings are more in keeping with sarcoidosis than myeloma.

B Oral methotrexate

In patients who are unresponsive to, or dependent on, steroids, methotrexate can be considered as a steroid-sparing option. It should not be considered a first-line option in patients requiring treatment.

D Bronchoscopy to look for mycobacteria

Mycobacterium tuberculosis is the organism responsible for TB. There is no mention in the history of night sweats or weight loss; the history and examination findings here are more in keeping with sarcoid than TB.

E Oral azathioprine

In patients who are unresponsive to, or dependent on, steroids, azathioprine can be considered as a steroid-sparing option. It should not be considered a first-line option in patients requiring treatment.

6607

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

40-year-old lady is admitted to the Emergency Department following shortness of breath of 3 days' duration. She is a patient known to have pulmonary sarcoidosis and had a percutaneous left upper lung biopsy a week ago. Now she presents with left-sided chest pain. She tells you that the pain worsens on inspiration and she has a cough that is not productive. Her past medical history includes: pulmonary fibrosis, asthma and irritable bowel syndrome. She takes dexamethasone 4 mg od and is not allergic to any medication. On examination, she is slightly dyspnoeic at rest with a blood pressure of 135/79 mmHg, pulse of 90/min and respiratory rate of 20/min. Her oxygen saturation is 94% on air. Her trachea is central but her chest expansion is minimally reduced on the left side. Breath sounds are reduced of the left side. Her apex beat is in the 5th intercostal space in the mid-clavicular line. Her cardiovascular and abdominal examination is normal.

Investigations:

Investigation	**Result**	**Normal value**
ECG	Normal sinus rhythm	
pH	7.40	7.35–7.45
pa(O_2)	9.1 kPa	10.5–13.5 kPa
pa(CO_2)	3.8 kPa	4.6–6.0 kPa
Bicarbonate	28 mmol/l	24–30 mmol/l
Sodium	137 mmol/l	135–145 mmol/l
Potassium	3.7 mmol/l	3.5–5.0 mmol/l
Urea	7.0 mmol/l	2.5–6.5 mmol/l
Creatinine	111 μmol/l	50–120 μmol/l
Haemoglobin	113 g/l	115–155 g/l
Platelets	233 x 10^9/l	150–400 × 10^9/l
Mean corpuscular volume (MCV)	83 fl	76–98 fl
White cell count (WCC)	10.2 x 10^9/l	4–11 × 10^9/l

X-ray shown below:



How would you treat her pneumothorax?

Your answer was incorrect

A Proceed to intercostal chest tube drainage

B Discharge with review after 48 h in the clinic

C 60% oxygen

D Start her on oxygen and just observe her

E Arrange for a bronchoscopy

Explanation

A Proceed to intercostal chest tube drainage

This is a secondary pneumothorax, which could have resulted either from the percutaneous lung biopsy or secondary to her pulmonary sarcoidosis. From the blood gases, this lady is also hypoxic and the pneumothorax is large on CXR (a large pneumothorax approximates to a 50% loss of lung volume). Since this is a large secondary pneumothorax it is appropriate to proceed to intercostal drain insertion. Oxygen replacement therapy is also advisable, as is suitable analgesia.

Image source-https://radiopaedia.org/cases/pneumothorax-6

C 60% oxygen

All patients who are hypoxic with a pneumothorax should be given oxygen. Patients who have a secondary pneumothorax that is less than 2 cm in size and are not breathless can be managed with high-flow oxygen and observation for 24 h. This is contraindicated in those who are oxygen sensitive. The size of this patient's pneumothorax makes this inappropriate.

B Discharge with review after 48 h in the clinic

If a patient has a unilateral pneumothorax that is less than 2 cm, is not breathless and the pneumothorax is primary rather than secondary, then the patient can be discharged with outpatient review. As the pneumothorax is secondary and large, this would be inappropriate.

D Start her on oxygen and just observe her

This option is not appropriate. High-flow oxygen should be given in pneumothorax as it helps to diffuse the nitrogen from the pneumothorax and thereby making it easier to reabsorb. As this patient has a large pneumothorax, this is inappropriate.

E Arrange for a bronchoscopy

Bronchoscopy would be useful in obtaining a biopsy of the bronchus and sometimes in tissue near the bronchus. It is not indicated for this patient.

32401

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BTS Pleural Disease Guideline 2010
thorax.bmj.com/content/65/Suppl_2.toc#BTSPleuralDiseaseGuideline2010
(http://thorax.bmj.com/content/65/Suppl_2.toc#BTSPleuralDiseaseGuideline2010)

Pneumothorax

Media

Expanded explanations

Flashcards

A 35-year-old, HIV-positive, East African patient presents to A&E with shortness of breath and night sweats. On examination he is thin and tachypnoeic, and has a right base that is stony dull to percussion with reduced air entry. On his chest X-ray there is a moderate-sized pleural effusion. His past medical history includes tuberculosis (TB) as a child for which he can’t remember whether he had treatment. His initial three sputum samples do not stain for acid-fast bacilli on Ziehl–Neelsen (ZN) stain. You want to carry out a test with the highest pick-up rate for TB diagnosis.

Which of the following would be the most useful?

Your answer was correct

A Pleural fluid aspirate culture

B Pleural biopsy culture

C Pleural biopsy histology

D CD4 count

E Heaf test

Explanation

B Pleural biopsy culture

Tuberculous effusion occurs in about 30% of patients with TB; this is comparable to HIV-positive and -negative patients. ZN stains are positive in about 75% of cases. Indications for pleural biopsy are exudative recurrent pleural effusion of unknown aetiology and pleural mass or thickening.

A Pleural fluid aspirate culture

ZN stains of pleural fluid are positive in 10–20% of tuberculous effusions.

C Pleural biopsy histology

Although it may demonstrate granulomas, it cannot give bacterial sensitivities.

D CD4 count

A low CD4 count not only predisposes to TB. CD4 cell counts give an indication of the health of the immune system. Monitoring CD4 cell counts is less important while taking HIV treatment than before starting. The count should go up when you take HIV treatment. Patients with higher CD4 counts present similar to those who are HIV-negative with TB.

E Heaf test

This test has been discontinued since 2005 and has been replaced by the Mantoux test – a person's medical risk factors determine at which increment (5, 10 or 15 mm) of induration the result is considered positive. Other techniques used include bronchoscopy/endobronchial ultrasound in those who have a non-productive cough/ unhelpful sputum culture and a high index of clinical suspicion.

Adenosine deaminase (ADA) levels are useful when TB is suspected in those with negative pleural fluid or biopsy cultures. However, raised ADA levels can be observed in other conditions leading to a false-positive diagnosis of TB, such as in empyema, rheumatoid effusion and malignancy.

Adenosine deaminase (ADA) levels are useful when TB is suspected in those with negative pleural fluid or biopsy cultures. However, raised ADA levels can be observed in other conditions leading to a false-positive diagnosis of TB, such as in empyema, rheumatoid effusion and malignancy.

32446

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301

Responses - % Correct: 27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 58-year-old man presents to the Emergency Department with approximately 500 ml of haemoptysis. He has a past medical history of previous tuberculosis (TB) infection, chronic obstructive pulmonary disease (COPD) and heart failure. His mobility is noted to be poor, and he is normally confined to a wheelchair.

An urgent computed tomography (CT) scan was carried out, and the result is shown below.



Which of the following is the most definitive management option in this case?

Your answer was correct

A Bronchial artery embolisation

B Tranexamic acid

C Blood transfusion

D Lobectomy

E Oral itraconazole

Explanation

A Bronchial artery embolisation

The image shown in the CT is classical for an aspergilloma. This would correlate with the patient’s past medical history of previous pulmonary TB, leaving a cavity for the aspergilloma to develop. In some patients, the aspergilloma can invade the bronchial artery. As the patient has undergone massive haemoptysis, urgent intervention is necessary. Given the patient’s comorbidities and performance status, he is unfit for a lobectomy, leaving embolisation as the next most appropriate management option.

B Tranexamic acid

While tranexamic acid would form part of the immediate management of the patient, it is not part of the definitive management of the patient. This would be bronchial artery embolisation.

C Blood transfusion

While a blood transfusion would form part of the immediate management of the patient, it is not part of the definitive management of the patient. This would be bronchial artery embolisation.

D Lobectomy

A lobectomy would be the most definitive management if the patient were otherwise fit and had a good performance status. However, in this case, research has shown the most appropriate definitive management as bronchial artery embolisation.

E Oral itraconazole

Oral itraconazole has been shown to successfully treat aspergillomas in up to 60% of patients. However, the patient presented is in a life-threatening scenario, and definitive management must focus on preventing further haemoptysis.

71793

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A 40-year-old nurse is admitted to the Medical Unit with a 2-day history of wheeze and exacerbation of asthma during peak pollen season. She has had this diagnosis for four years, since moving from her home in West Africa to the UK. She also has a history of hay fever since moving to the UK, and her daughter has developed asthma too. She has no other past medical history and is a non-smoker.

Investigations:

Investigation	Result	Normal value
Haemoglobin	135 g/l	115–155 g/l
White cell count (WCC)	10 × 10^9/l	4–11 × 10^9/l
Eosinophils	1.5 × 10^9/l	0–0.4 × 10^9/l
Platelets	350 × 10^9/l	150–400 × 10^9/l

Her lung fields on the chest X-ray show minor patchy shadowing.

Which additional test would you opt to do to best exclude causes, other than asthma, of her wheeze and current exacerbation?

Your answer was correct

A Antinuclear antibody (ANA)

B IgE

C *Aspergillus*-specific IgE radioallergosorbant (RAST) level

D Peak flow reading

E Stool culture for ova, cysts and parasites

Explanation

C *Aspergillus*-specific IgE radioallergosorbant (RAST) level

Allergic bronchopulmonary aspergillosis (ABPA) should always be excluded in people with asthma who have a raised eosinophil count. ABPA is a hypersensitivity reaction to *Aspergillus* in people with asthma. This can be done by sputum culture or testing for atopy to the fungi with RAST, skin-prick tests or IgE specific to the fungi. *Aspergillus* precipitins testing looks for the presence of the antibody in the serum; if the result is highly positive, the patient may have invasive aspergillosis.

A Antinuclear antibody (ANA)

Although eosinophil count can also be raised in Churg–Strauss syndrome, a small-to-medium-vessel vasculitis that presents usually in adults with asthma-like symptoms, this results in a positive pANCA, not ANA.

B IgE

A high eosinophil count may be due to her atopy and the high pollen count, so her total IgE would also be raised.

D Peak flow reading

Peak flow readings will likely be low in respiratory diseases like asthma, ABPA and Churg–Strauss, and therefore not good at excluding causes of wheeze and exacerbation.

E Stool culture for ova, cysts and parasites

Stool culture may be positive in a parasitic cause of eosinophilia pneumonia, but this is unlikely given the very minor chest X-ray changes.

32448

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A 30-year-old man from London, with no significant past medical history, went on holiday to Peru to climb the Inca Trail. He is a non-smoker. After climbing above 3500 m he developed a headache, associated with nausea and vomiting. He also felt tired and complained of insomnia. His respiratory rate was 22/min and his pulse was 90/min. The remainder of physical examination was unremarkable.

Investigations reveal:

Investigation	Result	Normal Values
Urea	10 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	70 μmol/l	50 - 120 μmol/l
Sodium (Na^+)	135 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5 - 5.0 mmol/l

That night it was noticed that he was disorientated and walking with a broad-based gait, and by the following morning he was drowsy and incoherent.

What is the most likely cause?

Your answer was incorrect

A Cerebellar infarction

B Cerebral haemorrhage

C High-altitude pulmonary oedema

D Meningococcal septicaemia

E High-altitude cerebral oedema

Explanation

E High-altitude cerebral oedema

There are three types of altitude illness: acute mountain sickness (AMS), high-altitude cerebral oedema (HACE) and high-altitude pulmonary oedema (HAPE). This man has acute mountain sickness with some evidence of cerebral oedema. The exact cause is unknown, but it is thought to be secondary to cerebral oedema and raised intracranial pressure. It is more common in people who live at low altitude.

Symptoms and signs of AMS are non-specific; however, diagnosis can be made using the criteria named in 'The Lake Louise Score', namely, the setting of recent gain in altitude, the presence of headache and at least one of the following symptoms:

- Gastrointestinal – anorexia, nausea and vomiting
- Fatigue and weakness
- Dizziness or light-headedness
- Difficulty sleeping.

He does not improve and progresses to develop ataxia and drowsiness. This is the development of high-altitude cerebral oedema, which can be considered as end-stage AMS.

Treatment is immediate descent, oxygen and intravenous dexamethasone.

Acetazolamide may be used for prevention and treatment of AMS. This is a carbonic anhydrase inhibitor which causes intracellular acidosis. It has a diuretic effect which can be useful in AMS/HACE and HAPE. It is also beneficial as it shifts the oxygen-dissociation curve to the right and therefore at lower partial pressures, more oxygen is released to the tissues.

C High-altitude pulmonary oedema

There is nothing in the history to suggest pulmonary oedema. There is no history of breathlessness.

A Cerebellar infarction

There is no history of unsteady gait and, given that the patient is at altitude, mountain sickness is more likely. Cerebellar infarction presents with non-specific symptoms such as dizziness, nausea and vomiting, unsteady gait and headache.

B Cerebral haemorrhage

There are other more likely diagnoses here. He has no neurological symptoms other than headache. Cerebral haemorrhage will present with neurological symptoms and signs depending on where the haemorrhage is located. For example, a cerebellar infarct will present with unsteady gait.

D Meningococcal septicaemia

There is no suggestion of fever, meningism or rash. Meningococcal infection is the most common cause of bacterial meningitis in the UK. There are several strains of meningococcal bacteria, such as A, B and C.

70108

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 57-year-old male patient with a 40 pack-year smoking history attends his GP with incessant daytime sleepiness, making it difficult for him to even attend work. His initial examination reveals a height of 180 cm, weight of 120 kg and neck circumference of 44 cm. He is referred for sleep studies, which show hypnoea/apnoea episodes consistent with the diagnosis of moderate to severe obstructive sleep apnoea (OSA).

Which of the following is the most appropriate treatment option?

Your answer was incorrect

A Overnight continuous positive airways pressure (CPAP)

B Amitriptyline

C Long-term oxygen therapy

D Nocturnal lorazepam

E Intra-oral device to prevent snoring

Explanation

A Overnight continuous positive airways pressure (CPAP)

The treatment of choice has been shown in studies to be effective in reducing OSA symptoms, and is first line in moderate–severe disease requiring intervention, but does not prevent the cardiovascular mortality associated with OSA. Hypnotics worsen OSA, as do alcohol, smoking and obesity, and so patients should alter their lifestyle appropriately, with weight loss and alcohol and smoking cessation as first-line treatment. Long-term oxygen therapy has not been shown to have any long-term benefit in OSA patients. Modafinil has been shown in some small studies to have beneficial effects in use with CPAP, but not as an alternative. Intra-oral devices may be useful for mild OSA, but not in daytime sleepiness.

B Amitriptyline

This is an inappropriate treatment option, as hypnotics can worsen OSA.

C Long-term oxygen therapy

This can be used as an adjunct to CPAP therapy or in cases where patients are unable to adhere to CPAP, but should not be offered first line in preference to CPAP.

D Nocturnal lorazepam

This is an inappropriate treatment option, as hypnotics can worsen OSA.

E Intra-oral device to prevent snoring

This may be useful for patients with mild OSA, but this patient has presented with daytime sleepiness and studies consistent with moderate–severe disease.

7174

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Obstructive Sleep Apnoea

Media

Expanded explanations

Flashcards

A 77-year-old man who smokes 20 cigarettes per day presents to the clinic with a chronic cough and, most recently, increasing haemoptysis. He has lost 4 kg in weight over the past 3 months. On examination he looks thin, with a BMI of 20. His BP is 140/75 mmHg. Auscultation of the chest reveals changes consistent with COPD.

Investigations reveal the following:

Investigation	**Result**	**Normal values**
Haemoglobin (Hb)	119 g/l	135–175 g/l
White cell count (WCC)	8.9 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	231 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	140 μmol/l	50–120 μmol/l
Erythrocyte sedimentation rate (ESR)	67 mm/hour	1–20 mm/hour
Chest X-ray (CXR)	Left-sided mass suspicious of bronchial carcinoma	
Trans-bronchial biopsy	squamous cell carcinoma	

Which of the following would be a contraindication to left pneumonectomy?

Your answer was incorrect

A Hypercalcaemia

B FEV_1 of 0.7 litres post bronchodilators

C MI within the past 6 months

D Tumour extending to the surface of the visceral pleura on CT

E Age 77 years

Explanation

⚙

B FEV$_1$ of 0.7 litres post bronchodilators

Patients who have an FEV$_1$ of less than 2.0 litres post bronchodilators are not recommended for pneumonectomy. VQ scanning may be of value in further assessment of such patients, although where there is a postoperative predicted FEV$_1$ of 0.8, pneumonectomy is effectively ruled out.

A Hypercalcaemia

Hypercalcaemia may be a paraneoplastic phenomenon related to PTHrp, or might actually have no link at all to the underlying bronchial carcinoma. Therefore it isn't a contraindication to surgery.

C MI within the past 6 months

Surgery is not recommended if the patient has suffered an MI within 6 weeks; a history of MI within 6 months necessitates a cardiology opinion and work-up, such as an echocardiogram, to assess preoperative ejection fraction.

D Tumour extending to the surface of the visceral pleura on CT

Tumour abutting the visceral pleura is not necessarily a contraindication to surgery, although more radical resection may be required and assessment of lymph node activity (for example with PET scanning) is likely to be undertaken first.

E Age 77 years

Age may not necessarily contraindicate surgery; functional status is more important in determining the chances of operative success.

21213

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 19-year-old man presents with a 10-day history of headaches and malaise, with the more recent development of a dry cough and breathlessness.

On examination, he was pyrexial, and he had a rash which looked like this:



Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	94 g/l	135 - 175 g/l
White cell count (WCC)	10 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	9.2 × 10^9/l	1.5 - 7.0 × 10^9/l
Platelets (PLT)	200 × 10^9/l	150 - 400 × 10^9/l
Reticulocytes	5.1%	0.5 - 2.5 %
Sodium (Na^+)	129 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.0 mmol/l	3.5 - 5.0 mmol/l
Urea	4.4 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	90 μmol/l	50 - 120 μmol/l
Bilirubin	30 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	44 u/l	10 - 40 u/l
Alanine aminotransferase (ALT)	30 u/l	7 - 55 u/l
Alkaline phosphatase (ALP)	121 u/l	30 - 150 u/l
Albumin	35 g/l	35 - 55 g/l
Lactate dehydrogenase (LDH)	495 u/l	140 - 280 u/l

His chest X-ray is shown below:



What is the most likely diagnosis?

Your answer was correct

A Epstein–Barr virus (EBV)

B *Legionella* pneumonia

C Streptococcal pneumonia

D Tuberculosis

E *Mycoplasma* pneumonia

Explanation

E *Mycoplasma* pneumonia

Mycoplasma usually affects young individuals and occurs in epidemics every 3–4 years. Chest symptoms are usually preceded by non-specific symptoms such as malaise and headaches.

The CXR usually shows only one lobe to be involved; however, about 20% show bilateral pneumonia. There is often a discrepancy between X-ray appearances and the clinical condition of the patient.

Diagnosis is by *Mycoplasma* serology – cold agglutinins occur in 50%. The image shows erythema multiforme, which further supports the diagnosis. The blood results suggest a haemolytic anaemia with a reticulocytosis, hyperbilirubinaemia and elevated LDH; autoimmune haemolytic anaemia (AIHA), when caused by cold agglutinins, is associated with *Mycoplasma* and can be diagnosed with a direct Coomb’s test.

Extra-pulmonary complications of *Mycoplasma* include:

- Cardiovascular – myocarditis and pericarditis

- Dermatological – erythema multiforme (as shown here), erythema nodosum, Stevens–Johnson syndrome and non-specific rashes
- Gastrointestinal – hepatitis, pancreatitis, nausea, vomiting, anorexia and transient abdominal pain
- Neurological – meningoencephalitis, meningitis, ascending paralysis, transient myelitis, cranial nerve palsies, peripheral neuropathy, bullous myringitis
- Haematological – cold autoimmune haemolytic anaemia, thrombocytopenia, disseminated intravascular coagulation
- Renal – glomerulonephritis
- Others – arthralgia, arthritis, bullous myringitis.

Treatment: macrolides.

Causes of erythema multiforme:

- Infections – herpes simplex virus (most common cause), orf, HBV, HIV, EBV, mumps (paramyxovirus), *Mycoplasma*, psittacosis, *Rickettsiae*, *Streptococcus*, typhoid, diphtheria
- Drug reactions – barbiturates, penicillin, sulphonamides, phenytoin
- Connective tissue disease – SLE
- Vasculitis – polyarteritis nodosa, granulomatosis with polyangiitits

Others – underlying malignancy, sarcoidosis, rheumatoid arthritis, ulcerative colitis.

A Epstein–Barr virus (EBV)

EBV causes infectious mononucleosis but is not associated with pneumonia. The disease is characterised by malaise and *upper* respiratory tract symptoms.

B *Legionella* pneumonia

Whilst dry cough, malaise, hyponatraemia and pneumonia are compatible with a diagnosis of *Legionella* pneumonia, the presence of erythema multiforme and headache, with normal LFTs, in a teenage patient is more consistent with *Mycoplasma* pneumonia.

C Streptococcal pneumonia

This patient presents with features of an atypical pneumonia. *Streptococcus pneumoniae* is the commonest cause of community-acquired pneumonia and would be associated with a sudden onset of symptoms, including fever and productive cough.

D Tuberculosis

TB tends to affect the upper lobes and is usually associated with fever, night sweats and weight loss. Additionally, no potential contact source of TB is mentioned, making this diagnosis more unlikely.

70075

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

As the medical SHO, you are admitting a 22-year-old asthma patient with an exacerbation of his asthma with a cough and severe wheeze that has been worsening over the past two days. He is normally on regular, short-acting β-agonist only and his predicted peak flow is 600 l/min. You are concerned that he may need review by the intensive care team.

On examination in the Emergency Department, his blood pressure (BP) is 110/72 mmHg, with pulse 110/min and regular. He has severe bilateral wheeze and his respiratory rate is 28/ min.

Investigations:

Investigation	Result	Normal value
PaO_2	9.4 kPa	10.5–13.5 kPa
$PaCO_2$	4.4 kPa	4.6–6.0 kPa
pH	7.37	7.35–7.45
PEFR	180 l/min (600 predicted)	

Which of the findings is most concerning?

Your answer was incorrect

A Respiratory rate 28/min

B Widespread wheeze

C Heart rate 110/min

D $PaCO_2$ 4.4 kPa

E Peak flow of 180 l/min

Explanation

E Peak flow of 180 l/min

The criteria for life-threatening asthma are well described in the BTS asthma guidelines. From the clinical scenario listed above, it is only his PEFR of 180 l/min which fits with those criteria, being less than 200 (33% predicted level). Given his peak flow, however, it’s obviously important that he is managed aggressively according to the guidelines, including the addition of IV magnesium if early response to nebulisers isn’t seen.

Criteria for life-threatening asthma:

- PEF <33% best or predicted
- SpO_2 <92%
- PaO_2 <8 kPa
- Normal (or elevated) $PaCO_2$ (4.6–6.0 kPa)
- silent chest
- cyanosis
- poor respiratory effort
- arrhythmia
- exhaustion, altered conscious level

C Heart rate 110/min

Tachycardia is an unreliable marker in asthma as it can easily occur due to repeat doses of salbutamol. However, arrhythmia implies a far worse state and should be treated quickly.

A Respiratory rate 28/min

Monitoring respiratory rate is important in asthma even though tachypnoea is not included in the criteria for life-threatening asthma, as a fall in respiratory rate can be associated with reduced respiratory effort and exhaustion.

B Widespread wheeze

Widespread wheeze indicates good respiratory effort in an acute asthma attack. If the chest were to become silent then this would indicate life-threatening asthma.

D $PaCO_2$ 4.4 kPa

Patients with exacerbation of asthma have a high respiratory rate and would be expected to have a low pCO_2 due to the increased CO_2 elimination. When CO_2 starts to rise, this implies that the patient is becoming tired and that means that the asthma can be classified as life-threatening.

32379

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Asthma

Media

Expanded explanations

Flashcards

A 28-year-old man is referred from the Infertility Clinic. He has a past history of recurrent sinusitis and a persistent cough with sputum production. The sweat test reveals a sodium concentration of 50 mmol/l.

His CXR is shown below:



What is the most likely diagnosis?

Your answer was incorrect

A	Bronchiectasis
B	Cystic fibrosis
C	α1-antitrypsin deficiency
D	Situs inversus

E Kartagener syndrome

Explanation

E Kartagener syndrome

This is primary ciliary dyskinesia associated with recurrent sinusitis, bronchiectasis and azoospermia. The CXR shows situs inversus with transposition of the major organs (as is the case in up to 50% of patients with ciliary dyskinesia). It also causes azoospermia, explaining his infertility. The mainstay of treatment for bronchiectasis is appropriate self-physio with postural drainage. Some physicians advocate the use of rotating antibiotic therapy, but this is associated in many cases with multi-drug resistance.

Image source: https://commons.wikimedia.org/wiki/File:Situs_inversus_chest_Nevit.jpg (https://commons.wikimedia.org/wiki/File:Situs_inversus_chest_Nevit.jpg)

D Situs inversus

While this patient has situs inversus as a component of his Kartagener syndrome, it would not adequately account for his sinusitis and infertility.

A Bronchiectasis

While bronchiectasis is a component of Kartagener syndrome, this alone would not adequately explain his presentation; Kartagener syndrome also consists of sinusitis and dextrocardia, and can cause azoospermia.

B Cystic fibrosis

This typically presents in infancy with respiratory symptoms and features of pancreatic insufficiency. Cystic fibrosis is diagnosed when the sweat test reveals a chloride concentration of >60 mmol/l.

C α1-antitrypsin deficiency

This is associated with respiratory symptoms, predominantly cough and dyspnoea, but would not adequately explain his situs inversus or infertility.

6608

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Asthma

Media

Expanded explanations

Flashcards

An 18-year-old male is brought to the Emergency Department by his parents who are concerned about the cough and fever that he has developed. The onset of the cough was about 5 days ago and over this time it has progressively worsened. The patient is also producing purulent sputum and complains of feeling short of breath. He has no past medical history and is not on any medication. On examination, he has a fever of 39°C and is visibly tachypnea. Examination of the chest shows reduced expansion on the right side associated with a reduced percussion note and bronchial breathing on auscultation.

Bloods are taken and his chest x-ray is shown below.



What is the most likely cause of this patients symptoms?

Your answer was incorrect

A Staphylococcal pneumonia

B *Pneumocystis jirovecii* pneumonia

C *Legionella* pneumonia

D *Haemophilus influenzae* pneumonia

E Pneumococcal pneumonia

Explanation

E Pneumococcal pneumonia

Despite appropriate antimicrobial therapy, pneumococcal pneumonia remains a cause of significant mortality owing to the ability of the organism to replicate in host tissues.

Young children and elderly adults are particularly prone to infections with *Streptococcus pneumoniae*. This organism is also the commonest cause of community-acquired pneumonia.

Clinical symptoms of this condition include fever and rigors, malaise, anorexia, cough, shortness of breath, production of purulent sputum and pleuritic chest pain. Clinical signs include tachycardia, hypoxia, fever, tachypnoea and signs of pulmonary consolidation (diminished expansion, dullness to percussion, bronchial breathing and increased tactile fremitus).

CXR classically shows lobar consolidation, this patient has right lower lobe pneumonia.

Image source: https://commons.wikimedia.org/wiki/File:Pneumonia_of_Right_Lung_Middle_Lobe.jpg

D *Haemophilus influenzae* pneumonia

Haemophilus influenzae infections are rare in patients over the age of 6 years, unless immunocompromised. The most common presentation is with meningitis.

A Staphylococcal pneumonia

The CXR in staphylococcal pneumonia typically shows a bilateral cavitating bronchopneumonia. Primary staphylococcal pneumonia is more commonly seen in children and young patients, with secondary disease being seen most commonly secondary to influenza infection.

B *Pneumocystis jirovecii* pneumonia

This is the most common opportunistic infection seen in patients with HIV infection.

C *Legionella* pneumonia

This is associated with a prodromal period of myalgia and headache, initially dry cough, confusion and patchy lower lobe consolidation on CXR. Pleural effusion is seen in as many as 30% of cases. It is associated with poorly maintained air conditioning systems, e.g. hotels, offices, etc.

6804

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 34-year-old woman presents with increasing breathlessness, complaining that she is no longer able to chase after her 4-year-old child. She also complains of chest pain and light-headedness and has recently noticed swelling of the ankles. There is no past medical history of note, particularly with respect to chronic chest disease or venous thromboses. On examination her body mass index (BMI) is 24, the jugular venous pressure (JVP) is elevated, and there is evidence of tricuspid regurgitation and right heart failure. Chest X-ray (CXR) shows enlargement of the pulmonary vessels. An echocardiogram is arranged, which shows markedly elevated right-sided pressures. There is hypoxia and hypocapnia on arterial blood gas estimation, with desaturation on exercise. Ventilation–perfusion scan is unremarkable.

Which of the following diagnoses fits best with this clinical picture?

Your answer was correct

A Chronic venous thromboembolism

B Idiopathic right heart failure

C Primary pulmonary hypertension

D Secondary pulmonary hypertension

E Pulmonary fibrosis

Explanation

C Primary pulmonary hypertension

Mean age at presentation for primary pulmonary hypertension is 36 years; patients characteristically complain of exertional dyspnoea, syncope and symptoms of right heart failure. Diagnosis is confirmed with echocardiogram, right heart catheterisation and computed tomography (CT) pulmonary angiogram or ventilation–perfusion scan to rule out significant thromboembolic disease. Autoantibodies such as anti-SCL-70 and anti-centromere may point to associated connective tissue disease. Options for intervention include PDE-5 inhibitors, endothelin receptor antagonists and prostacyclin receptor agonists.

A Chronic venous thromboembolism

Chronic venous thromboembolism is less likely given the absence of features to suggest thrombophilia, and the fact that this patient’s BMI is in the normal range.

B Idiopathic right heart failure

Isolated right heart failure is associated with the peripheral oedema seen here, although the pulmonary hypertension demonstrated is inconsistent with right heart failure alone.

D Secondary pulmonary hypertension

Secondary pulmonary hypertension occurs as a result of pulmonary fibrosis, either idiopathic or related to connective tissue disease, and as a result of chronic venous thromboembolism.

E Pulmonary fibrosis

There is no past medical history to suggest an underlying cause for pulmonary fibrosis, and no abnormalities are seen on the CXR to suggest extensive fibrotic disease.

32426

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pulmonary Embolism

Media

Expanded explanations

Flashcards

A 68-year-old alcoholic patient with known chronic obstructive pulmonary disease (COPD) was admitted to A&E with a 5-day history of green sputum and fevers. Past history of an inferior myocardial infarction is noted, and regular medication includes high-dose combination salmeterol/fluticasone inhaler, atorvastatin, ramipril and aspirin.

On examination, his BP is 95/65 mmHg, with pulse 100/min and regular. His respiratory rate is 25/min. There are signs of left lower lobe consolidation. He answers questions appropriately and his AMTS is 9/10.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	138 g/l	135–175 g/l
White cell count (WCC)	12.9×10^9/l	$4–11 \times 10^9$/l
Platelets	266×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5–5.0 mmol/l
Creatinine	90 µmol/l	50–120 µmol/l
Urea	8.2 mmol/l	2.5–6.5 mmol/l
Chest X-ray	Extensive left lower lobe consolidation	
pH	7.38	7.35–7.45
pO_2	8.9 kPa	10.5–13.5 kPa
pCO_2	4.6 kPa	4.6–6.0 kPa

What is his CURB-65 score?

Your answer was incorrect

A 1

B 2

C 3

D 4

E 5

Explanation

B | 2

The CURB-65 score comprises:

- age > 65 years
- confusion (defined as AMTS scoring ≤ 8)
- urea > 7 mmol/l
- respiratory rate > 30/min
- blood pressure < 90 systolic or 60 diastolic.

In this case, he scores increased risk for age and urea, so the correct answer is B, CURB-65 score of 2. The risk of mortality in patient scoring a 2 is 9.2%, therefore, inpatient treatment is indicated. A score of over 3 conveys a 22% mortality and patients should be considered for critical care admission, if appropriate. A score of 1 implies a 1.5% risk of mortality, indicating outpatient care can be instigated safely.

C | 3

This patient scores only a 2 on age and urea level. A score of over 3 conveys a 22% mortality and patients should be considered for critical care admission, if appropriate.

A | 1

This would have been correct if the patient only had one point. A CURB-65 score of 0–1 suggests that home treatment is possible as the risk of mortality stands at 1.5%.

D | 4

This patient scores only a 2 on age and urea level. A score of over 3 conveys a 22% mortality and patients should be considered for critical care admission, if appropriate.

E | 5

This patient scores only a 2 on age and urea level. A score of 4 or 5 conveys a 22% mortality and patients should be considered for critical care admission, if appropriate.

32378

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BTS management of CAP
brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-mana...
(https://www.brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/)

Pneumonia

Media

Expanded explanations

Flashcards

A 19-year-old man presents to the Emergency Department with a 3-day history of minor right-sided chest discomfort and lethargy. He smokes 20 cigarettes a day. A chest X-ray is obtained which shows a right-sided pneumothorax that is <1 cm from the chest wall. He is not distressed and he is not in significant pain. His oxygen saturation is 98% on air, his BP is 124/82 mmHg and his pulse is 81/min and regular.

How should he be managed?

Your answer was incorrect

A Pleural aspiration

B Small-bore chest drain insertion

C Oxygen therapy

D Outpatient clinic review

E Large-bore chest drain insertion

Explanation

D Outpatient clinic review

In a patient under the age of 50 without evidence of pre-existing lung disease, this is a primary pneumothorax. As the pneumothorax is <1 cm from the chest wall, it is reasonable to consider discharging the patient for review in the outpatient clinic.

C Oxygen therapy

Admission for high-flow oxygen therapy is indicated in secondary pneumothoracies <1 cm from the chest wall, or where aspiration has reduced the pneumothorax to this size. Care needs to be taken where there is oxygen sensitivity.

A Pleural aspiration

Primary pneumothoracies which are >2 cm from the chest wall, or are associated with breathlessness, should be aspirated. Secondary pneumothoarcies which are 1–2 cm from the chest wall should be aspirated.

B Small-bore chest drain insertion

In primary or secondary pnemothoracies where aspiration was initially indicated but has failed, and in secondary pneumothoracies >2 cm from the chest wall, insertion of an 8–14 F chest drain is indicated.

E Large-bore chest drain insertion

Large-bore chest drains are usually greater than 20 F. Such drains are indicated in haemothoracies and complex empyema.

21545

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Management of spontaneous pneumothorax: British Thoracic Society pleural disease guideline 2010
thorax.bmj.com/content/thoraxjnl/65/Suppl_2/ii18.full.pdf
(http://thorax.bmj.com/content/thoraxjnl/65/Suppl_2/ii18.full.pdf)

Pneumothorax

Media

Expanded explanations

Flashcards

A 56-year-old man is referred to the endocrine clinic with an 18-month history of constant tiredness. He works as a taxi driver and on one occasion fell asleep at the wheel. He finds it difficult to concentrate during the day and has frequent morning headaches. His wife stated that he is restless and snores loudly at night and occasionally has 'choking' episodes. He has also had very poor libido recently. He is a smoker of 30 cigarettes per day and drinks 4 pints of lager every night. His only medication is an inhaler for 'bronchitis', and he has had a chronic cough with sputum over the last few years.

On examination he was a large man with coarse features and was clinically obese, with body mass index (BMI) of 35. His heart sounds were normal, there was some scattered wheezing in the chest and he had early pitting ankle oedema.

What would be the most practical investigation to establish the diagnosis?

Your answer was incorrect

A Thyroid function tests

B Overnight home sleep oximetry

C Formal respiratory function tests

D CT thorax

E Epworth sleepiness score

Explanation

B Overnight home sleep oximetry

The diagnosis is obstructive sleep apnoea, which affects 1–2% of the population and is most commonly found in obese male smokers with a background of chronic obstructive pulmonary disease (COPD). Rarer causes include acromegaly, Cushing's disease, nasal deformities and tonsillar enlargement. Apnoea occurs when the walls of the pharynx collapse during sleep, which leads to hypoxia and increasingly strenuous respiratory efforts until the resistance is overcome. Apnoea can occur hundreds of times per night, resulting in reduction of REM sleep. The diagnosis can usually be made by a good history from the sleeping partner, supported by non-invasive oximetry performed at home. Full polysomnographic studies are rarely required for diagnosis but can be useful for research purposes. Management includes treating any underlying cause, weight reduction and avoidance of sedative drugs and alcohol. Continuous positive airway pressure (CPAP), which keeps the walls of the pharynx open, is the mainstay of treatment and can lead to a dramatic improvement in symptoms.

C Formal respiratory function tests

These may show evidence of obstructive lung disease and/or obesity hypoventilation syndrome, although this wouldn't prove or disprove obstructive sleep apnoea.

A Thyroid function tests

Although this patient is obese, it's most likely this is related to his lifestyle rather than to any underlying hypothyroidism. Hence TFT is unlikely to be of value here.

D CT thorax

This may show early changes consistent with COPD, although this won't prove or disprove obstructive sleep apnoea.

E Epworth sleepiness score

The Epworth sleepiness score is merely a pointer to daytime sleepiness associated with OSA; it isn't a confirmatory test. It can be used as screening of OSA.

9146

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219

Responses Total: 301

Responses - % Correct: 27%

Obstructive Sleep Apnoea

Media

Expanded explanations

Flashcards

A 27-year-old pregnant lady is referred to the Emergency Admission for sudden development of shortness of breath. She was well until 2 days ago when she developed shortness of breath. She has some slight left-sided chest pains, but no cough or haemoptysis. She is 28 weeks pregnant and has had no problems with the pregnancy. There is no history of long flights or swelling of her legs.

On examination, she appears anxious with a blood pressure of 110/60 mmHg, pulse rate of 92/min and a respiratory rate of 26/min. Her oxygen saturation on air is 98%. Her abdomen is distended and fetal movements are seen. Her chest is clinically clear. Her cardiovascular and lower limb examinations are unremarkable.

Investigations reveal the following:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	113 g/l	115–155 g/l
White cell count (WCC)	10.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	278 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	3.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	112 µmol/l	50–120 µmol/l
Mean corpuscular volume (MCV)	89 fl	76–98 fl
Urea	6.1 mmol/l	2.5–6.5 mmol/l
C-reactive protein (CRP)	6.0 mg/l	< 10 mg/l
D-dimers	0.62 mg/l	< 0.25 mg/l
pH	7.48	7.35–7.45
pa(O_2) on air	13.7 kPa	10.5–13.5 kPa
pa(CO_2) on air	3.9 kPa	4.6–6.0 kPa
Chest X-Ray (CXR)	No focal abnormality	
V/Q scan	Low probability of pulmonary embolism (PE)	

What would be the most appropriate step in her immediate management?

Your answer was correct

A Commence low-molecular weight heparin

B Commence warfarin

C Arrange for her discharge

D Commence low-dose aspirin

E Thrombolysis

Explanation

C Arrange for her discharge

A low-probability V/Q scan has a low clinical suspicion of pulmonary emboli (PE) and should prompt a search for another cause for the patient's symptoms. The slightly elevated pH, with O_2 towards the upper end of the normal range, accompanied by low CO_2 are more suggestive of mild hyperventilation and anxiety. Taking these facts into account, it is safe to facilitate discharge.

A Commence low-molecular weight heparin

There is no indication here for anticoagulation given the V/Q scan result.

B Commence warfarin

There is no indication for anticoagulation taking into account the V/Q result, and warfarin is not advised in early or in very late pregnancy.

D Commence low-dose aspirin

There is no indication for anti-platelet therapy, and even if there were evidence of thromboembolic disease, aspirin is not an effective intervention.

E Thrombolysis

Both thoracotomy with embolectomy and thrombolysis can be considered for pregnant women with massive pulmonary embolism, although that isn't the diagnosis here.

7511

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Thromboembolic Disease in Pregnancy and the Puerperium: Acute Management
rcog.org.uk/globalassets/documents/guidelines/gtg-37b.pdf
(https://www.rcog.org.uk/globalassets/documents/guidelines/gtg-37b.pdf)

A 55-year-old asthmatic patient presented in December with increasing cough and wheeze. She described a cough productive of brown sputum. She had no other past medical history. She was taking a steroid inhaler regularly and increasing doses of beta-agonist with no alleviation of her symptoms. She worked in a bakery for many years. She is an ex-smoker and stopped smoking 20 years ago. She has a dog at home.

On examination, she had a temperature of 38 °C. She was breathless at rest. She had reduced expansion anteriorly on the left side, with corresponding reduced breath sounds. She had a mild expiratory wheeze.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	141 g/l	135 - 175 g/l
White cell count (WCC)	12.7 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	7.37 × 10^9/l	1.5 - 7.0 × 10^9/l
Lymphocytes	2.55 × 10^9/l	1.5 - 4.0 × 10^9/l
Monocytes	0.6 × 10^9/l	0.0 - 0.8 × 10^9/l
Basophils	0.05 × 10^9/l	< 0.1 × 10^9/l
Eosinophils	1.9 × 10^9/l	< 0.45 × 10^9/l
Mean corpuscular volume (MCV)	87.3 fl	80 - 100 fl
Platelets (PLT)	314 × 10^9/l	150 - 400 × 10^9/l
Erythrocyte sedimentation rate (ESR)	28 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	16 mg/l	< 10 mg/l
Sodium (Na^+)	138 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5 - 5.0 mmol/l
Urea	4.8 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	95 µmol/l	50 - 120 µmol/l
Bilirubin	12 µmol/l	1 - 22 µmol/l
Alanine aminotransferase (ALT)	49 u/l	7 - 55 u/l
Alkaline phosphatase (ALP)	61 u/l	30 - 150 u/l
Albumin	36 g/l	35 - 55 g/l

Chest X-ray on presentation is shown below:



Chest X-ray taken one month earlier:



What is the diagnosis?

Your answer was incorrect

A Occupational asthma

B Exacerbation of asthma

C Allergic bronchopulmonary aspergillosis

D Churg-Strauss syndrome

E Loeffler syndrome

Explanation

C Allergic bronchopulmonary aspergillosis

The chest X-ray shows left upper lobe collapse and also the ‘veil sign’. This patient has allergic bronchopulmonary aspergillosis (ABPA). Pulmonary infiltrates, blood eosinophilia and asthma are usually secondary to Churg–Strauss or allergic bronchopulmonary aspergillosis.

ABPA is suspected in any patient with asthma who has an abnormal chest X-ray and high peripheral blood eosinophilia.The chest X-ray may show diffuse pulmonary infiltrates, and pulmonary, lobar or segmental collapse occur as transient features. The most common cause is sensitivity to *Aspergillus fumigatus* spores. The lack of other features of vasculitis such as mononeuritis, or markers of elevation such as raised CRP counts against vasculitis as an alternative diagnosis here.

The diagnostic criteria include:

- Asthma (in most cases)
- Peripheral blood and sputum eosinophilia
- Abnormal chest X-ray (infiltrates, segmental or lobar collapse)
- Positive skin tests/RAST to an extract of *A. fumigatus*
- *A. fumigatus* IgG serum-precipitating antibodies
- Raised total IgE >1000 ng/ml
- Fungal hyphae of *A. fumigatus* on microscopy of sputum.

The lung and eosinophils:

- Loeffler syndrome (eosinophils about 10% of blood WCC), also known as acute eosinophilic pneumonia/simple eosinophilic pneumonia. Mild self-limiting illness with transient migratory pulmonary shadows. Associated with parasitic infections, drug allergies and exposure to inorganic chemicals
- Tropical pulmonary eosinophilia (eosinophils >20%) – in tropical countries usually due to migrating larvae of the filarial worms *Wucheria bancrofti* and *Brugia malayi*
- Chronic/prolonged pulmonary eosinophilia (eosinophils >20%) – eosinophilic pneumonia persisting for more than one month. Chronic debilitating illness characterised by malaise, weight loss, fever and

dyspnoea

- Allergic bronchopulmonary aspergillosis (eosinophils 5–20%) as in this case
- Eosinophilic granulomatosis with polyangiitis (eosinophils 5–20%) associated with asthma
- Hyper-eosinophilic syndrome (eosinophils >20%: eosinophilic infiltrations of various organs – e.g. lungs, heart, bone marrow. Can be associated with an eosinophilic arteritis
- External agents – drugs, toxins, parasitic infection.

D Churg-Strauss syndrome

This patient does not have the diagnostic criteria for Churg–Strauss syndrome. Although the eosinophil count is elevated there is no other evidence of vasculitis.

A Occupational asthma

The patient working as a baker is irrelevant – occupational asthma is unlikely to cause a productive cough or these chest X-ray features. It can be screened for using serial peak flow measurements and comparing those at home to those at work.

B Exacerbation of asthma

The chest radiograph features and clinical examination findings are not in keeping with an exacerbation of asthma.

E Loeffler syndrome

Loeffler syndrome is associated with fleeting pulmonary infiltrates but not with intermittent lobar collapse. It is associated with parasitic worm migration to the lungs. We're given no indication that this is a potential cause here.

70094

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 62-year-old man with chronic lymphocytic lymphoma (CLL) presents to the General Practitioner, complaining of increased shortness of breath and wheezing. He has a history of asthma, for which he takes Seretide® (fluticasone propionate/salmeterol xinafoate), and he takes fludarabine for CLL.

On examination, he is pyrexial with a temperature of 38.1 °C, and there is a wheeze and coarse bilateral crackles on auscultation of the chest. He is diagnosed with *Pneumocystis jirovecii pneumonia*.

Which is the most appropriate prophylaxis against future *Pneumocystis* infection?

Your answer was incorrect

A Clindamycin 300 mg orally once a day (OD) and pyrimethamine 15 mg OD

B Co-trimoxazole 80 mg/400 mg orally OD

C Co-trimoxazole 80 mg/400 mg orally OD with pentamidine 300 mg by nebuliser once a month

D Pentamidine 300 mg by nebuliser once a month

E Co-trimoxazole 160 mg/800 mg orally three times weekly

Explanation

E Co-trimoxazole 160 mg/800 mg orally three times weekly

Patients who take fludarabine are at increased risk of infection with a range of pathogens, including *Pneumocystis, Listeria, Neisseria* and *Legionella*. Primary prophylaxis of Pneumocystis in these patients is with co-trimoxazole at a dose of 960 mg (total 16/800 split) three times per week. Indications for Pneumocystis prophylaxis in immunocompromised patients who are human immunodeficiency virus (HIV)-negative and not haematopoietic or solid organ transplant recipients include patients receiving chemotherapy regimens (associated with a 3.5% risk of Pneumocystis infection) (eg > 29 mg per day of prednisolone of > 1 month or those taking purine analogues), patients with inflammatory disorders receiving at least 16 mg per day of prednisolone for > 1 month and patients with certain primary immunodeficiencies.

C Co-trimoxazole 80 mg/400 mg orally OD with pentamidine 300 mg by nebuliser once a month

Co-trimoxazole alone is usually efficient prophylaxis against Pneumocystis and therefore, combination therapy is not recommended.

A Clindamycin 300 mg orally once a day (OD) and pyrimethamine 15 mg OD

This treatment combination may be considered as prophylaxis against *Pneumocystis* infection in immunocompromised patients who are solid organ transplant recipients. However, it is a third-line treatment option in this situation and would not be considered for this patient, as he has not received a solid organ transplant.

B Co-trimoxazole 80 mg/400 mg orally OD

This is the recommended dose of co-trimoxazole (trimethoprim/sulfamethoxazole) as prophylaxis against *Pneumocystis* infection in patients who are human immunodeficiency virus (HIV)-positive and have a CD4 count < 200 cells/µl.

D Pentamidine 300 mg by nebuliser once a month

Pentamidine is a suitable alternative prophylactic treatment if this patient is unable to tolerate co-trimoxazole. However, co-trimoxazole is the preferred primary agent, and there is nothing in the scenario to suggest that this would not be tolerated.

21473

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 70-year-old farmer is admitted with a 4-day history of feeling increasingly short of breath. He tells you he was kicked by a cow and was laid up with a swollen leg before the symptoms started. On examination there is extensive bruising and haematoma around the left knee, and the leg is more swollen than the right. He has bilateral basal crackles and occasional wheezes on inspiration, blood pressure (BP) is 110/80 mmHg and heart rate is 105 and regular; oxygen saturations 89–91% on pulse oximetry. The chest X-ray is shown below.



Which of the following represents the best treatment choice for this man?

Your answer was correct

A	Oral prednisolone
B	Anticoagulation
C	Intravenous furosemide
D	Nebulised salbutamol

E Penicillin IV

Explanation

B Anticoagulation

We are presented here with a patient who has had a blow to the knee, a period of inactivity and now has hypoxia with circulatory compromise. There are no signs of severe asthma on examination and the chest x-ray is normal. Given the differential leg size, pulmonary embolus is the most likely diagnosis and anticoagulation the treatment of choice. The injury to the knee is some days old, and therefore anticoagulation is unlikely to result in increased size of the knee haematoma. He may well also have other lung disease related to his farming occupation that accounts for the wheezes and crackles heard on auscultation.

Image source: https://search.creativecommons.org/photos/7785d80a-8ada-4c96-b21e-77ac49fdad7b (https://search.creativecommons.org/photos/7785d80a-8ada-4c96-b21e-77ac49fdad7b)

A Oral prednisolone

This would be an appropriate treatment for an exacerbation of asthma or COPD. The lack of significant wheeze counts against asthma as the cause of his symptoms here.

C Intravenous furosemide

This would be an appropriate treatment for cardiac failure/pulmonary oedema. The normal CXR counts against heart failure as the likely cause of his symptoms here.

D Nebulised salbutamol

This would be an appropriate treatment for an exacerbation of asthma or COPD. The lack of significant wheeze, and relatively acute onset of symptoms, counts against asthma or COPD as the underlying cause of this patient’s symptoms.

E Penicillin IV

This would be an appropriate treatment for pneumonia. The negative chest X-ray and absence of features such as pyrexia and raised inflammatory markers makes pneumonia a less likely diagnosis in this case.

6605

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pulmonary Embolism

Media

Expanded explanations

Flashcards

A 27-year-old woman presents with right-sided pleuritic chest pain. She says the pain came on suddenly while lifting her son into his push-chair some 30 h earlier. Her only medication of note is the oral contraceptive pill, and she is usually fit and well. The symptoms have largely resolved and she has not taken paracetamol for some 6 h.

On examination, BP is 120/70 mmHg, pulse 80/min and regular, breath sounds seem normal and O_2 saturation on air is 97%.

Investigations:

Investigation	Result	Normal value
Haemoglobin	129 g/l	115–155 g/l
White cell count (WCC)	8.2 × 10^9/l	4–11 × 10^9/l
Platelets	192 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine	110 μmol/l	50–120 μmol/l
Chest X-ray	Small right pneumothorax – rim less than 1cm	

Which of the following is the most appropriate next course of action?

Your answer was incorrect

A Send home with a plan for early review

B Admit for observation and oxygen therapy

C Admit for air aspiration

D Admit for formal chest drain insertion

E Admit for thoracoscopy

Explanation

A Send home with a plan for early review

This woman has a pneumothorax which is more than 1 day old, and the chest X-ray indicates that the size is small. Given that her symptoms are improving, it would be most appropriate to discharge her with a plan to review in the respiratory clinic, or to return if her pain significantly worsens. Air aspiration might be considered for a larger pneumothorax, whereas formal chest drain insertion would be indicated if the pneumothorax was larger and there was a previous history of chest disease.

C Admit for air aspiration

In the case of a primary spontaneous pneumothorax, air aspiration is only indicated if the pneumothorax is >2 cm and/or the patient is breathless. If this fails to resolve the symptoms or the pneumothorax does not decrease in size, then formal chest drain insertion would be needed.

B Admit for observation and oxygen therapy

The patient’s oxygen saturations are 97% on air, so there is no indication for supplemental oxygen. As this is a spontaneous pneumothorax in an otherwise well patient

D Admit for formal chest drain insertion

Formal chest drain insertion is only indicated in symptomatic/>2 cm primary spontaneous pneumothorax where aspiration has failed. It is first line if there is underlying lung disease and the patient is breathless or the pneumothorax is >2 cm in size.

E Admit for thoracoscopy

Thoracoscopy is usually performed to investigate pleural disease or in preparation for pleurodesis. This patient has had a spontaneous pneumothorax and there is no current suspicion of underlying lung disease. The annual incidence of primary spontaneous pneumothorax is around 5/100,000 in women and around 25/100,000 in men. The greatest risk factor is smoking.

27768

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Management of spontaneous pneumothorax: British Thoracic Society pleural disease guideline 2010
thorax.bmj.com/content/65/Suppl_2/ii18
(http://thorax.bmj.com/content/65/Suppl_2/ii18)

BTS Pleural Disease Guideline 2010
brit-thoracic.org.uk/document-library/clinical-information/pleural-disease/pleural-disease-guidelines-2010/appendix-3-spontaneous-pneum...
(https://www.brit-thoracic.org.uk/document-library/clinical-information/pleural-disease/pleural-disease-guidelines-2010/appendix-3-spontaneous-pneumothorax-poster-pleural-disease-guideline/)

Pneumothorax

Media

Expanded explanations

Flashcards

A 27-year-old known asthmatic patient is brought into the Emergency Department by the ambulance crew after an acute severe attack of asthma. On arrival, he is cyanosed, his respiratory rate is 9 breaths/min and his heart rate is 48 bpm. He is started on high-flow oxygen (non-rebreather mask). While assessing him, the patient suddenly develops respiratory arrest. The patient still has radial pulse of 50 bpm and intubation is attempted. While trying to intubate him, he starts to actively vomit a little.

What would be the immediate next appropriate step?

Your answer was incorrect

A Apply firm cricoid pressure

B Release the cricoid and perform suctioning

C Go ahead and intubate quickly

D Call the anaesthetist

E Turn the patient over

Explanation

B Release the cricoid and perform suctioning

During emergency intubation, to minimise the risk of gastric aspiration, the Sellick manoeuvre (firm pressure over the cricoid cartilage) may be initiated as soon as positive pressure ventilation is started and should normally be continued until inflation of the tracheal cuff of the endotracheal tube in the trachea. However active vomiting, which creates oesophageal pressures greater than 60 cmH_2O, is a contraindication to using this technique. In this case, the appropriate action is therefore to release the cricoid and suction.

C Go ahead and intubate quickly

The priority in this case is to minimise the risk of gastric aspiration, followed by return to pre-oxygenation with 100% oxygen.

A Apply firm cricoid pressure

Active vomiting, which creates oesophageal pressures greater than 60 cm H_2O is a contraindication to using the Sellick manoeuvre. Applying firm cricoid pressure in this case can result in the rupture of the oesophagus.

D Call the anaesthetist

Though senior specialist support should be sought, initial measures must be undertaken to stabilise, and prevent further deterioration of the patient.

E Turn the patient over

In the context of a respiratory arrest, the risk of gastric aspiration is best minimised by suctioning aspirates rather than repositioning the patient.

7503

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Respiratory Failure

Expanded
explanations

A 65-year-old man with known COPD comes to see you at the Respiratory Clinic. He has been a smoker for 40 years and previously worked in the docks at Southampton. He has experienced no increase in his sputum volume or any change in its colour. On examination, he is pursed-lip breathing but managing complete sentences. His BP is 140/70 mmHg, pulse 80 bpm and regular, $Sp(O_2)$ 93% o/a, respiratory rate 18 breaths/min and temperature 37°C. Examination of his chest revealed a widespread wheeze with coarse crepitations heard in the left mid-zone. FEV_1:FVC ratio in the clinic today was 69%.

What is the most useful investigation in establishing the diagnosis?

Your answer was incorrect

A Repeat spirometry

B Chest X-ray

C V/Q scan

D High-resolution CT

E Full blood count

Explanation

D High-resolution CT

Patients with longstanding chronic obstructive pulmonary disease (COPD) often develop localised areas of bronchiectasis because of progressive damage to the lung tissue. It is important to distinguish between this and fibrosis (secondary to asbestos exposure) in the absence of infective processes so that adequate treatment can be employed. The most useful investigation is therefore a CT scan.

C V/Q scan

VQ scanning is most useful in identifying evidence of an acute pulmonary embolus. That would not fit with the picture of wheeze and coarse crepitations on auscultation here.

A Repeat spirometry

Spirometry is not useful and it is likely that it would confirm an obstructive picture, with a low level of sensitivity to identify the onset of any restrictive lung disease.

B Chest X-ray

CXR has only a low pick-up rate for bronchiectasis or pulmonary fibrosis versus CT and would therefore not be considered useful here.

E Full blood count

FBC may show an increase in Hb because of chronic hypoxia, although this would be non-specific with respect to the cause of any underlying hypoxia (i.e. worsening COPD versus asbestos-related pulmonary fibrosis). 2360

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Asbestos-related Disease

Media

Expanded explanations

Flashcards

A 62-year-old patient presents to the Respiratory Clinic. He has been complaining of progressively worsening shortness of breath and dry cough. He denies any haemoptysis or chest pain. He smoked when he was younger and has roughly a ten pack-year history of smoking cigarettes. He used to work as a welder. He has a pet dog and lives in a house with his wife. He does not recall any asbestos exposure.

His past medical history includes gastro-oesophageal reflux disease (GORD), cholecystectomy for gallstones and hypertension. He normally takes amlodipine, omeprazole and ramipril.

Auscultation demonstrates crepitations in both lung bases.

His hands are as shown below.

What is the most likely diagnosis?



Your answer was incorrect

A	Bronchiectasis
B	Pulmonary fibrosis
C	Congestive cardiac failure
D	Pulmonary arterial hypertension
E	Pulmonary embolism

Explanation

B Pulmonary fibrosis

The presence of a slow-onset shortness of breath, dry cough, bibasal crepitations and clubbing makes pulmonary fibrosis the most likely diagnosis. The patient's demographics fit well with this diagnosis as most patients with pulmonary fibrosis present in their 60s. Pulmonary fibrosis is accumulation of excessive fibrous tissues in a process termed fibrosis. This causes reduced oxygenation. Many cases are idiopathic, but environmental pollutants can be involved, such as in asbestosis and silicosis. Some cases are associated with underlying connective tissue diseases, and others are medication induced, eg amiodarone, methotrexate, nitrofurantoin and bleomycin.

The first-line investigations would be spirometry, for which a restrictive pattern would be found, and a chest X-ray, which would demonstrate a reticulonodular shadowing. Furthermore, a high-resolution computed tomography (HRCT) of the chest would be done and, if the diagnosis remains in doubt, a biopsy may be needed.

C Congestive cardiac failure

Heart failure can cause progressive shortness of breath, especially in patients who have a smoking history and hypertension. It can also cause cough, but this is generally productive with white or pink sputum. However, the differentiating factor is clubbing. Clubbing is not caused by heart failure.

A Bronchiectasis

Bronchiectasis is a suppurative chronic lung disease characterised by scarred and thickened airways with chronic mucous secretion. It can be idiopathic, but many cases are due to viral infection, cystic fibrosis, autoimmune disease and tuberculosis. While bronchiectasis would explain worsening shortness of breath and cough, as well as the clubbing, bronchiectasis is clinically characterised by high mucous production. The absence of mucous makes this very unlikely. One would also expect crepitations which change in character on coughing.

D Pulmonary arterial hypertension

Pulmonary arterial hypertension is a rare condition characterised by sustained elevation in pulmonary arterial pressure and pulmonary vascular resistance without any other clear cause. Presentation is with progressive dyspnoea progressing to oedema and ascites, and late stage may be complicated by angina and arrhythmia. Clinically, one can expect a right ventricular heave, loud second heart sound and raised jugular venous pressure (JVP) on examination. The absence of these findings and the presence of clubbing make this unlikely.

E Pulmonary embolism

Pulmonary embolism is an obstruction of the pulmonary artery due to thrombosis, fat, amniotic fluid or air. Presentation is with sudden dyspnoea, pleuritic chest pain, cough and haemoptysis. It is not a cause of clubbing and would be unlikely to cause the slow progression of symptoms or the presence of crepitations bilaterally. It could be considered if there was further evidence of malignancy and the symptoms were more sudden in their nature, as the clubbing could then be associated with a malignancy.

72629

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 24-year-old man presents to the Emergency Department having felt unwell for the last couple of days. He tells you that he has had flu for the last four days. He complains of a dry cough, headache, fever, malaise and muscle pains.

On examination, he is unwell with a blood pressure of 110/60 mmHg, pulse of 92 beats per minute and a respiratory rate of 22 breaths per minute. His oxygen saturation is 95% on air. On auscultation, he has scattered crackles in both lung fields. His cardiovascular and abdominal examinations are unremarkable.

Investigations reveal the following:

Investigation	Result	Normal Values
Haemoglobin (Hb)	98 g/l	135 - 175 g/l
White cell count (WCC)	12.1 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	278 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	130 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.5 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	112 μmol/l	50 - 120 μmol/l
Mean corpuscular volume (MCV)	89 fl	80 - 100 fl
C-reactive protein (CRP)	86 mg/l	< 10 mg/l
Blood film	Fragmented red cells, reticulocytosis	
Coombs' test	+	

How would you treat this patient?

Your answer was correct

A Ciprofloxacin

B Clarithromycin

C Flucloxacillin

D Gentamicin

E Rifampicin

Explanation

B Clarithromycin

This is the typical presentation of *Mycoplasma pneumoniae* with a few days' history of dry cough and flu-like symptoms. Cold-agglutinins associated with the infection may also drive an autoimmune haemolytic anaemia with positive Coombs' test. Infection is seen in epidemics, particularly where young adults live in close proximity, for example in an army barracks.

A Ciprofloxacin

Ciprofloxacin could be an option in combination with vancomycin for the treatment of community-acquired pneumonia where patients are penicillin allergic but is not a first line choice.

C Flucloxacillin

Flucloxacillin is an option for post-influenza pneumonia, which is considered as a possible diagnosis when patients present with cavitating pneumonia following flu infection.

D Gentamicin

Gentamicin was considered additive to the effects of penicillins in severe infection – for example in endocarditis, although is now mainly reserved for the treatment of severe Gram-negative infections.

E Rifampicin

Rifampicin is used as dual therapy in the treatment of severe legionella infection.

71080

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 21-year-old comes to the Emergency Room with symptoms of an upper respiratory tract infection and sharp stabbing chest pain over his left upper chest when he breathes in. He is usually well and is doing a degree in physical education. He is a non-smoker who only drinks small amounts of alcohol.

On examination, his blood pressure (BP) is 135/70 mmHg, his temperature is 37.9 °C, his pulse is 84/min and regular. Auscultation of the chest is clear, although there are some transmitted sounds from his upper respiratory tract. He has some limited anterior chest wall tenderness.

Investigations reveal the following:

Investigation	**Result**	**Normal values**
Haemoglobin (Hb)	141 g/l	135–175 g/l
White cell count (WCC)	8.2×10^9/l	$4.0–11.0 \times 10^9$/l
Platelets (PLT)	202×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	100 µmol/l	50–120 µmol/l
Electrocardiogram (ECG)	Sinus rhythm with a rate of 84; no acute changes	
$_PO_2$	13 kPa	10–14 kPa
$_PCO_2$	4.8 kPa	4.7–6.0 kPa

His chest x-ray is shown below.



Which of the following is the most likely diagnosis?

Your answer was incorrect

A Viral pericarditis

B Viral myocarditis

C Viral costochondritis

D Hyperventilation syndrome

E Pleurisy

Explanation

C Viral costochondritis

In the context of a young, previously well individual with a recent upper respiratory tract infection, normal ECG, normal CXR and normal ABG the history points towards viral costochondritis. Adequate analgesia is the most important component of successful treatment, with simple analgesia such as paracetamol combined with an appropriate non-steroidal. Without analgesia, the risk of atelectasis and consequent bacterial superinfection is increased.

Image source: https://commons.wikimedia.org/wiki/File:Chest_x-ray.jpg (https://commons.wikimedia.org/wiki/File:Chest_x-ray.jpg)

B Viral myocarditis

Signs and symptoms of myocarditis include chest pain, general malaise, history consistent with a recent viral illness, congestive heart failure, arrhythmias, fever and sudden death. ECG changes include diffuse T-wave inversion and saddle-shaped ST-elevation. There are many causes of myocarditis, but viral infection is the most common. The gold standard diagnostic test is myocardial biopsy.

A Viral pericarditis

Pericarditis classically presents with sharp, pleuritic chest pain which is relieved by sitting up and leaning forwards. Additionally, there may be a pericardial friction rub on auscultation. Classical ECG findings are widespread saddle-shaped ST-elevation. Patients may also develop a pericardial effusion.

The common viral causes of pericarditis include enteroviruses (e.g. Coxsackieviruses, echo- viruses); herpesviruses (e.g. Epstein–Barr virus, cytomegalovirus, human herpesvirus 6); adenoviruses, parvovirus B19. Non-infectious causes of pericarditis include auto-immune, neoplastic, metabolic, traumatic and drug related. NSAIDs are used in the treatment of pericarditis.

Reference: Imazio M, Gaita F, LeWinter M. Evaluation and treatment of pericarditis: A systematic review. *JAMA*. 2015;314(14):1498–1506. doi:10.1001/jama.2015.12763

D Hyperventilation syndrome

In hyperventilation syndrome, the patient breathes rapidly causing P_aCO_2 to fall. This results in a respiratory alkalosis. Patients often describe chest pain, and tingling around the mouth and fingers. These features may accompany a panic attack.

E Pleurisy

Pleurisy describes inflammation of the pleura which causes sharp, stabbing pain when breathing. On auscultation of the chest it may be possible to hear a pleural friction rub. Patients often complain of lateral chest pain rather than the anterior upper chest pain seen here and don't complain of pain on chest wall palpitation as is seen here.

21231

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Viral Infections and Antivirals

Expanded explanations

A woman with known rheumatoid arthritis presents with breathlessness, which has increased over the past 10 days, accompanied by night sweats and a dry cough. Currently, she takes prednisolone 3 mg and methotrexate 10 mg once per week, ranitidine and folic acid.

Her chest X-ray is shown below.



By -- Samir धर्म 06:38, 14 January 2007 (UTC) [GFDL (http://www.gnu.org/copyleft/fdl.html) or CC-BY-SA-3.0 (http://creativecommons.org/licenses/by-sa/3.0/)], via Wikimedia Commons

Oxygen saturations are 97% on air but fall to 90% on exercise. Other bloods are unremarkable, apart from her lymphocyte count, which is slightly below the lower limit of the normal range.

What is the best investigation to elicit the likely diagnosis?

Your answer was incorrect

A Bronchoscopy and lavage

B Lung function tests

C Arterial blood gases

D Ventilation/perfusion (V/Q) scan

E Echocardiogram

Explanation

A Bronchoscopy and lavage

The presence of exercise-induced oxygen desaturation in an immunocompromised or immunosuppressed person should always raise the possibility of *Pneumocystis jiroveci* pneumonia (PCP). In this case, however, the possibility of methotrexate-induced pneumonitis should also be considered. Both conditions can be associated with dyspnoea, exercise-induced desaturation and fever. Radiographic features vary considerably, with diffuse, bilateral interstitial or alveolar infiltrates also possible in both conditions. In this case, the X-ray shows diffuse bilateral interstitial changes.

Laboratory findings in PCP are generally unhelpful. Diagnosis can be made by staining spontaneous or induced sputum samples. In patients with human immunodeficiency virus (HIV), bronchoalveolar lavage diagnostic yield approaches 95% and this can be raised to nearer 100% with the use of transbronchial biopsy.

C Arterial blood gases

Arterial blood gases may reveal hypoxia but will not rule significant infection in or out. As such, they are not the preferred investigation here.

B Lung function tests

This patient may have underlying methotrexate-related pneumonitis where a restrictive picture is seen on lung function tests, but it is important to exclude PCP first.

D Ventilation/perfusion (V/Q) scan

A V/Q scan may reveal ventilatory defects consistent with infection but will not confirm the nature of the pathogen.

E Echocardiogram

Although many immune modulators can affect myocardial function, methotrexate leads to pulmonary and hepatic fibrosis.

9837

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

As the medical registrar, you are asked on the phone by your FY1 to review a 26-year-old male patient who she has just examined. The patient has presented with progressively worsening dyspnoea and chest discomfort over the previous 24 h. He had been given a salbutamol inhaler by his general practitioner (GP) some three months earlier for exercise-induced wheeze and shortness of breath.

On examination, he is tachycardic at 120 bpm and tachypnoeic at 30 bpm, with bilateral wheeze on auscultation. On examination of the supraclavicular areas and neck, there is palpable cutaneous emphysema.

What is the most likely diagnosis?

Your answer was correct

A Oesophageal rupture

B Asthma

C Gas gangrene

D Pulmonary embolism

E Lung malignancy

Explanation

B Asthma

Surgical emphysema, also known as subcutaneous emphysema, is a pathological accumulation of air or gas in the subcutaneous tissues of the body. It is a rare condition, but it can occur from air tracking from the lungs (pneumothorax, COPD, asthma), from trauma – particularly barotrauma (pneumo-mediastinum, mechanical ventilation) or, rarely, a rupture of the oesophagus. However, it can also occur from the production of gas within a tissue by an infection such as in gas gangrene. It causes a crackling sensation felt on palpation of the skin of the neck and chest but it may also cause a diffuse swelling of the chest wall and neck.

A Oesophageal rupture

This is most commonly associated with procedures such as bronchoscopy, with spontaneous rupture occurring most often in association with vomiting. There may be dysphagia, tachycardia, dyspnoea and fever.

C Gas gangrene

While this can cause cutaneous emphysema, patients are usually significantly unwell, with evidence of sepsis or septic shock and visible necrotic tissue on examination.

D Pulmonary embolism

This is not commonly associated with cutaneous emphysema, and we are not given any classical risk factors for DVT/PE.

E Lung malignancy

While lung malignancy can lead to cutaneous emphysema, it is a relatively rare complication; the patient's young age and absence of risk factors for lung malignancy also make this a less likely option.

6805

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

You review a 26-year-old woman who has been admitted to the neurological ward for treatment. She complained of ascending symmetrical weakness and loss of sensation affecting both lower limbs. This began after a diarrhoeal illness. A diagnosis of Guillain–Barré syndrome was made. She is complaining to the nurses of increasing shortness of breath. You are aware that a deterioration in respiratory function can predict mortality. She is 50 kg in weight. FVC on admission was noted to be 1.5 l. You arrange for it to be checked again.

What level of FVC would indicate the threshold for respiratory support?

Your answer was correct

A 1.5 l

B 1.4 l

C 1.3 l

D 1.0 l

E 0.6 l

Explanation

D 1.0 l

Deterioration in FVC is closely predictive of deterioration with opportunistic respiratory tract infection, and associated increased morbidity and mortality. Patients with Guillain–Barré syndrome should be monitored with regular spirometry. The threshold for increased respiratory intervention is said to be around 20 ml/kg FVC, which would put this at around 1 l in this patient. Minimal sedation and aggressive physiotherapy while on the Intensive Care Unit can serve to reduce the possibility of infection. Overall poor prognosis in Guillain–Barré is associated with rapid progression of symptoms, advanced age, prolonged ventilation (>1 month) and severe action potential reduction on neuromuscular testing.

A 1.5 l

1.5 l corresponds to around 30 ml/kg FVC, which is above the recommended threshold for respiratory support.

B 1.4 l

1.4 l is above the level where respiratory support is likely to be required, meaning that patients may be inappropriately admitted to the High-dependency Unit, putting pressure on bed space for other patients.

C 1.3 l

1.3 l is around 30% above the level where respiratory support is likely to be required.

E 0.6 l

At 0.6 l FVC, type 2 respiratory failure and pneumonia are possible sequelae. Respiratory support should happen at the higher threshold of 1.0 l for FVC.

71074

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Guillain-Barré Syndrome

Media

Expanded explanations

Flashcards

A 32-year-old woman presents to her doctor complaining of increasing shortness of breath. Over the past year or so she has also noticed that her hands go white and are very painful if she ventures out into cold weather. On examination there is evidence of subcutaneous calcification in the fingers. Fine bi-basal crepitations and occasional wheezes are heard on auscultation of the chest.

Investigations reveal the following:

Investigation	Result
Anti-nuclear antibody	Positive
Anti scl-70 antibody	Positive
FVC	3.2 litres
FEV_1	2.6 litres

Which of the following represents the most appropriate treatment in this case?

Your answer was correct

A Commence inhaled corticosteroids

B Commence inhaled short-acting beta-2-agonist therapy

C Commence immuno-suppressant therapy

D Commence combined inhaled corticosteroid and beta-2 agonist therapy

E Commence inhaled long-acting anti-cholinergic therapy

Explanation

C Commence immuno-suppressant therapy

This woman has evidence of systemic sclerosis, with clinical features and the scl-70 positivity pointing towards associated lung fibrosis. Her lung function tests also suggest the possibility of a restrictive lung defect, further supporting this diagnosis. Around 15% of patients have disease which is associated with more rapidly progressive lung fibrosis. Lung fibrosis in systemic sclerosis may also be associated with an increased risk of lung cancer, and is associated with the development of pulmonary hypertension.

Management is usually with immunosuppressant. Mycophenolate (https://www.uptodate.com/contents/mycophenolate-mofetil-cellcept-and-mycophenolate-sodium-myfortic-drug-information?source=see_link) mofetil has a good evidence base and is often used first line, with cyclophosphamide being an alternative. Low-dose corticosteroids can be used as an adjunct in some cases, but high-dose steroids should be avoided if possible due to risk of scleroderma crisis.

A Commence inhaled corticosteroids

Unfortunately inhaled corticosteroids are ineffective in the treatment of pulmonary fibrosis and have no role in the management of this patient.

B Commence inhaled short-acting beta-2-agonist therapy

Inhaled short-acting beta-2-agonists are of course the initial therapy of choice for asthma, but have no value in relieving symptoms associated with pulmonary fibrosis.

D Commence combined inhaled corticosteroid and beta-2 agonist therapy

Combined inhaled corticosteroid and LABA therapy is utilised for patients with asthma who fail to gain control of symptoms on inhaled corticosteroids alone.

E Commence inhaled long-acting anti-cholinergic therapy

Long-acting anti-cholinergics are of value in the treatment of COPD, not pulmonary fibrosis.

18555

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 64-year-old man was referred to the clinic with insomnia. He complained of falling asleep during the day and having an early morning headache. He had a past history of a cholecystectomy and gout. He took no other regular medication. Examination was unremarkable. His weight and height were 120 kg and 175 cm, respectively

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	161 g/l	135 - 175 g/l
White cell count (WCC)	6.4×10^9/l	$4.0 - 11.0 \times 10^9$/l
Platelets (PLT)	94×10^9/l	$150 - 400 \times 10^9$/l
Mean corpuscular volume (MCV)	101 fl	80 - 100 fl
Thyroid-stimulating hormone (TSH)	1.2 mu/l	0.4 - 5.0 mu/l

Which of the following investigations would most aid diagnosis?

Your answer was incorrect

A ABG

B Sleep studies

C CXR

D CT thorax

E Body plethysmography

Explanation

B Sleep studies

This patient has obstructive sleep apnoea (OSA), suggested by insomnia, daytime somnolence, morning headache and obesity. Other symptoms include poor concentration during the day, and partners may describe snoring followed by apnoeic episodes.

Diagnosis is made using the Epworth score, which is a measure of daytime somnolence, and sleep studies, which would show apnoeic/hypopnoeic episodes associated with desaturations, increase in heart rate and arousal from sleep. Treatment will depend on severity of symptoms and desaturations. In most cases weight loss is advisable.

In mild OSA syndrome a mandibular advancement device may control symptoms. In moderate to severe OSA syndrome, nocturnal non-invasive ventilation (NIV) using continuous positive airway pressure (CPAP) can be considered. Early-morning headaches can be caused by a transient hypercapnia, which corrects to normal when the patient is awake. If a patient has hypercapnia in waking hours (and not just on an early morning blood gas) then overlap syndrome with obesity hypoventilation syndrome should be considered, although severe OSA may cause persistent hypercapnia in some cases. Patients with persistent hypercapnia should be considered for bi-level positive airway pressure (BiPAP).

It is important to exclude underlying causes; this patient has a history of gout which, together with the raised MCV and low platelets, suggests alcohol consumption, which can precipitate the problem. Hypothyroidism, acromegaly and sedating drugs also need to be excluded. Retrognathia can cause OSA, and large tonsils may obstruct the airway.

C CXR

Radiology is usually normal in obstructive sleep apnoea.There are no respiratory symptoms to suggest an intrathoracic pathology that would require a CXR.

A ABG

Arterial blood gases would be helpful in assessing for hypercapnoea, but are not diagnostic of OSA.

D CT thorax

Radiology is usually normal in obstructive sleep apnoea.There are no respiratory symptoms to suggest an intrathoracic pathology that would require a CT.

E Body plethysmography

Body plethysmography may reveal restrictive lung function with a normal gas transfer but this is not a diagnostic test for OSA.

70139

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy



Peer Responses %

Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Obstructive Sleep Apnoea

Media

Expanded explanations

Flashcards

A 27-year-old woman is referred to the Emergency Department for sudden development of shortness of breath, with left-sided chest pains but no cough or haemoptysis. She is 28 weeks into an uneventful pregnancy.

On examination, she appears anxious; her BP is 110/60 mmHg, heart rate 92 bpm and respiratory rate 26 breaths per minute. Her oxygen saturation on air is 98%. Her chest is clinically clear. Her cardiovascular and lower limb examinations are unremarkable.
Investigations:

Investigation	Result	Normal Value
Haemoglobin (Hb)	113 g/l	115–155 g/l
White cell count (WCC)	10.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	278 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	3.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	112 μmol/l	50–120μmol/l
Urea	6.1 mmol/l	2.5–6.5 mmol/l
C-reactive protein (CRP)	6 mg/l	< 10 mg/l
D-dimers	0.62 mg/l	< 0.25 mg/l

Which of the following investigations would be the most appropriate next step?

Your answer was incorrect

A Bilateral lower limb Doppler scan

B CXR

C Echocardiography

D CT pulmonary angiography

E Ventilation/perfusion scan

Explanation

B CXR

This patient has symptoms of a potential pulmonary embolus, and Green-top guidance from the Royal College of Obstetricians and Gynaecologists (RCOG) recommends a CXR as an initial step in investigation. A D-dimer level is unreliable when measured in pregnancy and guidance does not support its use.

C Echocardiography

Echocardiography is most useful as a bedside investigation when there is a high suspicion of pulmonary embolism and the patient is unstable. It may reveal evidence of right ventricular strain. Here, the patient is stable, and therefore, echocardiography is less likely to contribute towards the diagnosis.

A Bilateral lower limb Doppler scan

Given that the lower limbs appear normal on examination, a bilateral lower limb Doppler scan in this case, is unlikely to prove useful in ruling in or out a pulmonary embolus.

D CT pulmonary angiography

CT pulmonary angiography (CTPA) is the preferred investigation post-CXR in the event that the chest X-ray is found to be abnormal. The choice of performing either CTPA or ventilation/perfusion scan (V/Q) is a balance between the risk of radiation exposure to the pregnant woman's breast tissue (much higher with CTPA) and that of radiation exposure to the fetus (which is higher with V/Q scanning).

E Ventilation/perfusion scan

According to the RCOG guidelines, ventilation/perfusion V/Q or CTPA can be considered once a CXR has been performed, although where the CXR has proven to be abnormal, CTPA is preferred to V/Q.

73464

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Thromboembolic Disease in Pregnancy and the Puerperium: Acute Management
rcog.org.uk/globalassets/documents/guidelines/gtg-37b.pdf
(https://www.rcog.org.uk/globalassets/documents/guidelines/gtg-37b.pdf)

You are asked to review a 64-year-old man with a history of chronic obstructive pulmonary disease (COPD), prior to planned surgery for a large inguinal hernia. You take a history and examine him. He is worked up with investigations before proceeding to the operation.

Which of the following is most associated with increased risk of peri-operative complications?

Your answer was correct

A Nature of the surgery (hernia repair)

B CO_2 level of 6.2 kPa on air-breathing arterial blood gas sample

C FEV$_1$ (forced expiratory volume in 1 s) of 1.4 l

D History of treated hypertension

E Ability to walk 1.2 miles on the flat

Explanation

B CO_2 level of 6.2 kPa on air-breathing arterial blood gas sample

A raised CO_2 on air would have most impact on increased peri-operative risk in this case. Whilst epidural anaesthesia may offer a reduction in peri-operative risk, atelectasis due to continued abdominal pain is still associated with increased incidence of postoperative lower respiratory tract infection. Postoperative pain relief and aggressive physiotherapy support are therefore essential components of treatment.

A Nature of the surgery (hernia repair)

Although abdominal surgery is associated with increased risk of postoperative respiratory failure, hernia repair is a relatively minor procedure and adequate analgesia should minimise the risk of perioperative complications.

C FEV$_1$ (forced expiratory volume in 1 s) of 1.4 l

FEV$_1$ <0.8 l is particularly associated with operative risk. Obesity, because of splinting of the diaphragm, may also drive postoperative respiratory failure.

D History of treated hypertension

Controlled blood pressure should not be associated with significant operative risk. Recent ischaemic cardiovascular events on the other hand do significantly impact on outcome.

E Ability to walk 1.2 miles on the flat

Ability to walk 1.2 miles implies a functional status associated with low risk of operative complications. 32412

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media Expanded explanations Flashcards

A young alcoholic patient presents with a fever, haemoptysis, green sputum and a left-sided effusion clinically. He has been unwell with swinging fevers for the last week. On further questioning he admits to drinking 8 pints of beer per day and to not eating properly. You are concerned that it may be an empyema.

Which of the following tests would be most useful in supporting your suspicion?

Your answer was incorrect

A Pleural fluid microscopy and culture

B Pleural fluid white cell count and differential

C Urinary pneumococcal antigen

D Serum pneumococcal antigen

E Pleural fluid pH

Explanation

E Pleural fluid pH

A pleural fluid pH of less than 7.2 is highly suggestive of an empyema, and is the most useful investigation here.

D Serum pneumococcal antigen

As with a urinary pneumococcal antigen test, though a serum pneumococcal antigen result would be of diagnostic help, neither a positive nor a negative result would confirm the presence of empyema. Given the patient's history of alcohol excess, the likelihood of a *Klebsiella* pneumonia, a common cause of empyema, is high, rather than a pneumococcal pneumonia.

A Pleural fluid microscopy and culture

While indispensible if positive in order to tailor antimicrobial therapy, only 60% of pleural fluid cultures are positive.

B Pleural fluid white cell count and differential

Although a raised WCC (> 10) is seen in empyema, and more rarely in other exudative effusions, it may not be diagnostic here. A low pleural fluid pH in contrast is likely to be confirmatory for empyema.

C Urinary pneumococcal antigen

While a urine pneumococcal antigen result would be of diagnostic help, neither a positive nor a negative result would confirm the presence of empyema.

7178

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Empyema

Expanded
explanations

54-year-old stonemason was referred to the clinic with a 3-year history of progressive breathlessness. He is an ex-smoker (40 pack-years), having stopped 14 years ago.

On examination he looked well, was not clubbed and there was no lymphadenopathy. His chest was hyper-expanded and he had basal inspiratory crackles.

His spirometry revealed:

FEV_1	58% predicted (1.71 litres)
FVC	70% predicted (2.4 litres)
KCO	55% predicted

His CXR is shown below:



What is the most underlying diagnosis?

Your answer was incorrect

A Silicosis

B Asbestosis

C Byssinosis

D Berylliosis

E Simple pneumoconiosis

Explanation

A Silicosis

Silicosis is a fibrotic lung disease associated with inhalation of silica (silicon dioxide), which is highly fibrogenic. It is usually seen in quarrying and mining occupations and in sandblasters, ceramic workers, pottery workers and in stonemasons if the dust generated contains quartz.

Acute silicosis may present within 12 months if exposure has been heavy, and is characterised by dry cough and breathlessness. Patients deteriorate rapidly over 1–2 years and there is no effective treatment to slow the progression.

Chronic silicosis runs a more chronic course, as in this patient, and is associated with a gradual worsening of breathlessness, with restrictive lung function and a reduced transfer factor.

Chronic silicosis is associated with silicotic nodule formation (3–5 mm in diameter), predominantly affecting the upper lobes. These nodules may coalesce and cause progressive massive fibrosis. Diagnosis is made on history of exposure and XR changes. The CXR may show hilar eggshell calcification (pathognomonic of silicosis), as in the CXR shown, with or without upper zone pulmonary nodules and fibrosis.

B Asbestosis

There is no mention of potential exposure to asbestos, and the chest radiograph shows no pleural or diaphragmatic calcification to suggest previous asbestos exposure.

C Byssinosis

Byssinosis is associated with exposure to cotton dust, flax and hemp.

D Berylliosis

Berylliosis is associated with inhalation of fumes from the molten metal beryllium.

E Simple pneumoconiosis

Simple pneumoconiosis, also known as simple coal worker’s pneumoconiosis, is associated with exposure to coal dust and is classically seen in miners. It is usually relatively asymptomatic unless progressive massive fibrosis develops.

70077

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Silicosis

Media

Expanded explanations

Flashcards

A 17-year-old woman with stable cystic fibrosis presents to the Emergency Department with a cough productive of purulent sputum tinged with blood, and significantly worse shortness of breath over the past 48hrs. Her CF is currently well managed and she is Pseudomonas negative. On examination she is pyrexial 38.1°C, BP is 95/55 mmHg; heart rate is 85/min and regular, respiratory rate is 32. There are coarse crackles at the right base, and you also notice a couple of cold sores at the corner of her mouth. O_2 saturation on air is 92%. Her BMI is 23.

Investigations;

Investigation	Result	Normal value
Haemoglobin	121 g/l	115–155 g/l
White cell count (WCC)	13.4 × 10^9/l (neutrophilia)	4–11 × 10^9/l
Platelets	189 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Urea	8 mmol/l	2.5–6.5 mmol/l
Creatinine	90 µmol/l	50–120 µmol/l
Glucose	5.1 mmol/l	3.5–5.5 mmol/l
Alanine aminotransferase (ALT)	34 U/l	5–30 IU/l
Alkaline phosphatase (ALP)	95 U/l	30–130 IU/l
Albumin	37 g/l	35–55 g/l
Chest X-ray	Right lower lobe consolidation	

Which of the following is the most appropriate antibiotic option?

Your answer was incorrect

A IV Ciprofloxacin

B IV Gentamicin

C Nebulised Tobramycin

D Oral Azithromycin

E Oral Clarithromycin and IV Co-amoxiclav

Explanation

E Oral Clarithromycin and IV Co-amoxiclav

The answer is Oral Clarithromycin and IV Co-amoxiclav -

Patients with cystic fibrosis are more susceptible to community-acquired pneumonia. The short history, coupled with lobar consolidation and evidence of herpes simplex, raises the possibility of *Streptococcus pneumoniae* infection. Oral Clarithromycin and IV Co-amoxiclav is therefore the most appropriate intervention with respect to antibiotics. Past medical history, examination and investigations including both albumin and glucose, suggest this patient is maintaining relatively good nutritional status with respect to her cystic fibrosis.

B IV Gentamicin

IV Gentamicin is incorrect, it is an alternative option for Pseudomonas infection, and is not first line for community-acquired pneumonia.

A IV Ciprofloxacin

IV Ciprofloxacin is incorrect, it is an option for Pseudomonas infection, but that seems less likely given the clinical history here.

C Nebulised Tobramycin

Nebulised Tobramycin is incorrect, it is a nebulised option for chronic Pseudomonas colonisation. When *P. aeruginosa* is first detected, prompt treatment using inhaled tobramycin alone (300 mg in 5 ml, administered twice daily) for 28 days, rather than a regimen including other antibiotics is recommneded. The therapy is repeated only if surveillance cultures show recurrence of *P. aeruginosa*.

D Oral Azithromycin

Oral Azithromycin is incorrect, it is given for 6 months in patients colonised with Pseudomonas because of data suggesting it improves outcomes.

40423

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 32-year-old woman with a history of intravenous drug abuse attends for review. She has a history of gradually increasing shortness of breath over the past 18 months to 2 years. There has been associated intermittent fevers and weight loss of some 5 kg. She has tested negative for hepatitis B some 6 months earlier. On examination there are lower zone inspiratory crackles.

Investigations reveal haemoglobin (Hb) of 92 g/l and a diffuse increase in immunoglobulins, renal function testing is normal. Chest X-ray is shown below. Lung biopsy reveals diffuse interstitial lymphoid infiltrates and pulmonary function tests suggest a restrictive lung defect.



Which of the following diagnoses fits best with this clinical picture?

Your answer was incorrect

A Lymphoid interstitial pneumonia

B Idiopathic pulmonary fibrosis

C Viral pneumonia

D Hypersensitivity pneumonitis

E Respiratory bronchiolitis-associated interstitial lung disease

Explanation

A Lymphoid interstitial pneumonia

Clinical features of lymphoid interstitial pneumonia include gradual-onset breathlessness over a considerable time, intermittent fevers and weight loss, with inspiratory crackles on examination. There is mild anaemia with a diffuse increase in immunoglobulins. The chest x-ray showing a thin walled cystic lesion in the right mid zone and histology revealing lymphoid interstitial infiltrates are in keeping with the diagnosis. Steroids may improve symptoms, although around one-third of patients progress to extensive fibrosis. Associations include HIV, connective tissue disease, hepatitis B infection, chronic active hepatitis and Hashimoto's thyroiditis.

Image source: https://radiopaedia.org/cases/lymphocytic-interstitial-pneumonitis-with-infective-exacerbation-of-bronchiectasis (https://radiopaedia.org/cases/lymphocytic-interstitial-pneumonitis-with-infective-exacerbation-of-bronchiectasis)

B Idiopathic pulmonary fibrosis

Idiopathic pulmonary fibrosis implies there is no identified underlying cause; against a background of IV drug abuse, secondary interstitial lung disease should be excluded before describing the changes seen here as idiopathic.

C Viral pneumonia

Given the chronic nature of this patient's symptoms, viral pneumonia is very unlikely to be responsible.

D Hypersensitivity pneumonitis

Although hypersensitivity pneumonitis may be seen, particularly against a background of cocaine smoking, it is stated that this patient's mode of drug abuse is primarily intravenous.

E Respiratory bronchiolitis-associated interstitial lung disease

RBALD is primarily associated with heavy smoking and presents with pigment-loaded macrophages in the pulmonary infiltrates rather than the lymphocytes seen here.

32439

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 52-year-old immigrant from the Balkans presents for review. Over the past few months he has suffered a number of fevers and night sweats and has lost a significant amount of weight. His chest x-ray shows:



Tuberculin skin testing is positive. Tapping of the fluid reveals a lymphocytosis, although staining for acid and alkali-fast bacilli (AAFB) is negative, and protein count is 41 g/l.

What diagnosis fits best with this clinical picture?

Your answer was incorrect

A Lymphomatous pleural effusion

B Tuberculous pleural effusion

C Para-pneumonic effusion

D Carcinomatous pleural effusion

E Transudate effusion secondary to heart failure

Explanation

B Tuberculous pleural effusion

TB effusions present with this pattern of symptoms, accompanied by a small to moderate effusion rich in lymphocytes. Acid and alkali-fast bacilli (AAFB) staining is only positive in 5–10%, although cultures are positive in 25% of cases over 2–6 weeks. Induced sputum or polymerase chain reaction (PCR) for TB done at a specialist centre may improve the diagnostic yield, as may pleural biopsy. Fluid volumes may increase during anti-tuberculous therapy; corticosteroids are associated with reduced fluid volume but have no effect on outcome; and pleural thickening and calcification unfortunately occur commonly. Tuberculin skin testing and the fact that he has come from the Balkans are pointers to underlying TB.

Image source: https://radiopaedia.org/cases/pleural-effusion-7?lang=gb (https://radiopaedia.org/cases/pleural-effusion-7?lang=gb)

A Lymphomatous pleural effusion

This diagnosis could explain the fever, night sweats, weight loss and effusion, but is very rare. The presence of a tuberculin skin test being positive points to a much more common diagnosis.

C Para-pneumonic effusion

In the presence of pneumonia, a para-pneumonic effusion can develop. However, as the patient has no evidence of chest infection, this is an unlikely diagnosis. Also, such an effusion is likely to be smaller in volume.

D Carcinomatous pleural effusion

Carcinomatous pleural effusion is the second most common reason for an exudative pleural effusion. It is most associated with cancer of the lung, ovary and breast, as well as lymphoma. In malignancy, there can be increased pleural membrane permeability and increased pressure as the lymphatic system becomes obstructed. The presence of night sweats, fevers and weight loss makes this a strong differential diagnosis, but the positive tuberculin test and immigration from an endemic area makes TB of higher likelihood.

E Transudate effusion secondary to heart failure

In heart failure, patients can develop pleural effusions. However, these effusions are typically transudate rather than exudate in nature, and would be unlikely to have lymphocytes. In heart failure, the effusion would more likely to be bilateral as well, and this patient does describes no features of heart failure, nor any risk factors.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total: 301

Responses - % Correct: 27%

Pleural Effusion

Media

Expanded explanations

Flashcards

A 50-year-old housewife is being followed up in the chest clinic for her asthma. Over the past 2 years she has had intermittent low-grade fevers, weight loss, cough and shortness of breath. On examination she has end-expiratory wheeze with occasional crackles that do not resolve on coughing. She does not have any clubbing. She is a non-smoker. Her past medical history includes asthma and atrial fibrillation. Her drug history includes salbutamol inhaler, high-dose steroid inhaler and amiodarone. Her chest X-ray shows peripheral infiltrates on the outer two-thirds of the peripheries. Her sputum microbiology is negative but she has a raised number of eosinophils. Her blood tests show a raised serum eosinophil count, IgG and ESR.

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Amiodarone-induced fibrosis

B Chronic eosinophilic pneumonia (CEP)

C Allergic bronchopulmonary aspergillosis (ABPA)

D Treatment-resistant asthma

E Acute eosinophilic pneumonia (AEP)

Explanation

B Chronic eosinophilic pneumonia (CEP)

Eosinophilic pneumonia is a rare respiratory pathology characterised by pulmonary infiltrates and pulmonary eosinophilia. It can be idiopathic or secondary to parasites, drugs or systemic diseases such as vasculitis. The peripheral eosinophil count is not always raised. The serum IgG is raised in two-thirds and the ESR is elevated. Sputum and bronchoalveolar lavage often show an eosinophilia.

This presents over months to years. CEP usually occurs in those aged in their 50s, with a female:male ratio of 2:1. It can be irreversible. About 60% of patients with CEP have asthma and 90% are non-smokers. Patients present with constitutional symptoms, cough, wheeze and fever. Chest X-ray in CEP shows dense, bilateral peripheral infiltrates often ‘opposite to pulmonary oedema’ with a distribution in the outer two-thirds of the lung. Treatment is with steroids.

C Allergic bronchopulmonary aspergillosis (ABPA)

There would be an elevated total serum IgE rather than IgG and IgG precipitins to *Aspergillus fumigatus*.

A Amiodarone-induced fibrosis

In this condition, changes are usually seen in the bases. The treatment is to stop the drug, and steroids may help.

D Treatment-resistant asthma

Asthma does not present with intermittent low-grade fevers, and CXR would be normal rather than show pulmonary infiltrates.

E Acute eosinophilic pneumonia (AEP)

This presents over a few days. Patients with AEP are acutely unwell with temperature, tachycardia and tachypnoea. AEP chest X-ray may show bilateral diffuse infiltrates associated with an effusion.

32445

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 31-year-old woman with asthma presents to the Emergency Department with a 1-day history of a productive cough and shortness of breath. Before this, her symptoms have been stable for the last six months and she stopped taking her inhalers.

On examination, she is unable to speak in full sentences. Her respiratory rate is 30 breaths per minute, heart rate 115 bpm and BP 100/60 mmHg. Auscultation of her chest reveals an expiratory wheeze throughout.

Investigations:

Investigation	Result	Normal value
Peak expiratory flow rate (PEFR)	150 litres/min	400–700 litres/min
pH	7.47	7.35–7.45
Partial pressure of carbon dioxide ($p(CO_2)$)	4.06 kPa	4.7–6.0 kPa
pO_2	12 kPa	10.5–13.5 kPa
Bicarbonate	21.5 mmol/l	22–32 mmol/l

Which of the following would be the best initial management for this patient?

Your answer was incorrect

A High-flow oxygen, PO/IV steroid, ipratropium nebulisers and salbutamol nebulisers

B 60% oxygen, PO/IV steroid, salbutamol nebulisers, ipratropium bromide nebulisers and leukotriene receptor antagonist

C 60% oxygen, PO/IV steroid, salbutamol and IV magnesium

D High-flow oxygen, salbutamol nebulisers, ipratropium bromide nebulisers and IV magnesium

E High-flow oxygen, IV/PO steroids, ipratropium bromide nebuliser and IV aminophylline

Explanation

A High-flow oxygen, PO/IV steroid, ipratropium nebulisers and salbutamol nebulisers

After stopping this patient's inhalers, she is no longer on any medication to control her asthma. She presents with symptoms and signs of an acute severe asthma exacerbation – inability to complete a sentence in one breath, a respiratory rate greater than 25 breaths per minute, tachycardia greater than 110 bpm and peak expiratory flow rate (PEFR) less than 50%. The correct management includes:

- the administration of high-flow oxygen, as patients with pure asthma do not retain carbon dioxide
- salbutamol driven by oxygen; and steroid therapy – prednisolone 40 mg PO if the patient is able to take PO medication or 100 mg IV hydrocortisone
- Ipratropium may also be used
- IV magnesium should be given in life-threatening asthma or if there is no response to initial therapy
- IV aminophylline can also be used.

C 60% oxygen, PO/IV steroid, salbutamol and IV magnesium

IV magnesium is reserved for patients with a PEFR less than 50% predicted and who have not had a good response to inhaled bronchodilator therapies.

B 60% oxygen, PO/IV steroid, salbutamol nebulisers, ipratropium bromide nebulisers and leukotriene receptor antagonist

Leukotriene receptor antagonists are used in the long-term management of asthma but are not appropriate as an acute treatment for an exacerbation.

D High-flow oxygen, salbutamol nebulisers, ipratropium bromide nebulisers and IV magnesium

Steroids are a key aspect of acute treatment for asthma. Additionally, IV magnesium is reserved for patients with a PEFR less than 50% predicted and who have not had a good response to inhaled bronchodilator therapies.

E High-flow oxygen, IV/PO steroids, ipratropium bromide nebuliser and IV aminophylline

Salbutamol nebulisers are highly effective in acute asthma exacerbations and are missing here. Additionally, while aminophylline can be used in such patients, they are used only in treatment-resistant patients.

70124

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British guideline on the management of asthma
brit-thoracic.org.uk/quality-improvement/guidelines/asthma/
(https://www.brit-thoracic.org.uk/quality-improvement/guidelines/asthma/)

A 35-year-old man is admitted with night sweats, cough and general malaise. He is tachycardic at 121 bpm and dyspnoeic, and chest examination reveals scattered wheeze.

On the second day of his admission, he experiences haemoptysis and haematuria is noted. He has a background of anxiety and a previous deep vein thrombosis (DVT). His regular medications include cetirizine, warfarin and diazepam.

Investigations:

Haemoglobin (Hb)	145 g/l	(135–155 g/l)
White cell count (WCC)	15.1 × 10^9/l	(4–11 × 10^9/l)
Creatinine (Cr)	176 µmol/l	(50–120 µmol/l)
MPO antibodies	Negative	
PR3 antibodies	Positive	
24-hour urine 5-hydroxyindoleacetic acid (5-HIAA)	Positive	
First sputum sample	Negative	

His chest X-ray is shown below.



What is the most likely cause of this patient's presentation?

Your answer was correct

A Anti-glomerular basement membrane (GBM) antibody disease

B Carcinoid tumour of the lung

C Eosinophilic granulomatosis with polyangiitis

D Granulomatosis with polyangiitis

E *Mycobacterium tuberculosis*

Explanation

D Granulomatosis with polyangiitis

This case features widespread symptoms, suggesting a multi-system cause. The combination of renal impairment, haemoptysis, haematuria, pyrexia and malaise may raise suspicion of a vasculitis. The positive PR3 points to granulomatosis with polyangiitis as a likely cause.

A Anti-glomerular basement membrane (GBM) antibody disease

This patient has an unusual widespread history, which indicates pulmonary involvement and possible vasculitis. While anti-GBM is associated with haemoptysis, the blood results point to an **ANCA-positive** vasculitis, rather than anti-GBM in this case.

B Carcinoid tumour of the lung

The non-specific symptoms and pulmonary involvement, alongside the raised urine 5-HIAA levels, raise suspicion of carcinoid here. However, this picture is complicated by renal impairment and PR3 antibody positivity. It is important to remember that urine 5-HIAA levels can be artificially raised by certain medications, eg diazepam and warfarin. This lowers the likelihood of carcinoid in this case and points to an alternate cause.

C Eosinophilic granulomatosis with polyangiitis

Eosinophilic granulomatosis with polyangiitis is associated with positive MPO antibodies, rather than PR3.

E *Mycobacterium tuberculosis*

Tuberculosis may seem to be a possible diagnosis here with a history of dyspnoea, haemoptysis and night sweats. The renal impairment could possibly be renal tuberculosis. However, the blood results point to an ANCA-positive vasculitis, in this case granulomatosis with polyangiitis.

71776

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Granulomatosis with Polyangiitis

Media

Expanded explanations

Flashcards

A 74-year-old male smoker with a 30-pack-year smoking history has an FEV1 of 40% predicted. He has successfully reduced his smoking habit to only 3 cigarettes per day with help from smoking cessation services. However, he can only walk 100 m at a time before he becomes out of breath and feels that he needs to stop. His medication includes an inhaled long-acting beta-agonist and tiotropium. On examination, his BP is 155/85 mmHg and he has coarse crackles and wheeze throughout both lung fields.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	121 g/l	135–175 g/l
White cell count (WCC)	5.9×10^9/l	$4–11 \times 10^9$/l
Platelets	210×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5–5.0 mmol/l
Creatinine	139 µmol/l	50–120 µmol/l
Bicarbonate	30 mmol/l	24–30 mmol/l
pH	7.4	7.35–7.45
P_aO_2	10.1 kPa	10.5–13.5 kPa
P_aCO_2	6.4 kPa	4.6–6.0 kPa

Which would best improve his exercise tolerance?

Your answer was correct

A Inhaled beclometasone

B Inhaled salbutamol

C Portable oxygen

D Pulmonary rehabilitation

E Smoking cessation

Explanation

D Pulmonary rehabilitation

Given his limited smoking habit, it is likely that a pulmonary rehabilitation programme will have more impact.

A Inhaled beclometasone

Low-dose inhaled corticosteroids have no impact on disease progression in COPD, although it appears that high-dose inhaled corticosteroids combined with a long-acting beta-2 agonist does reduce exacerbations, at the expense of increased pneumonia.

B Inhaled salbutamol

Salbutamol is not likely to impact significantly as most patients have very poor reversibility.

C Portable oxygen

His P_aO_2 is too high for portable oxygen to be of benefit.

E Smoking cessation

Smoking cessation will have some impact on exercise tolerance, but given his limited habit now, it is likely that a pulmonary rehabilitation programme will have more impact.

21484

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Pulmonary rehabilitation in chronic obstructive pulmonary disease
sciencedirect.com/science/article/pii/S1745045407000329
(http://www.sciencedirect.com/science/article/pii/S1745045407000329)

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 71-year-old woman with a previous history of pulmonary tuberculosis presents to the Respiratory Clinic complaining of lethargy, weight loss, night sweats and intermittent severe haemoptysis. She has a history of hypertension managed with ramipril and indapamide, but is otherwise well.

On examination, her blood pressure is 142/87 mmHg, she is apyrexial and her body mass index (BMI) is 21 kg/m^2. Her FEV$_1$ is 1.8 l.

Investigations reveal the following:

Investigation	Result	Normal values
Haemoglobin (Hb)	131 g/l	115–155 g/l
White cell count (WCC)	10.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	233 × 10^9/l	150–400 × 10^9/l
Sodium (Na$^+$)	139 mmol/l	135–145 mmol/l
Potassium (K$^+$)	4.9 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	130 µmol/l	50–120 µmol/l
Chest X-ray (CXR)	Left upper lobe cavity with rim of air surrounding a solid lesion	
Aspergillus precipitant test	+	
FEV$_1$	1.8 l	

A computed tomography (CT) chest scan confirms the presence of a density with the classical appearance of an aspergillioma.

Which of the following is the most appropriate treatment for her?

Your answer was incorrect

A IV amphotericin

B Surgical resection

C Bronchial artery embolisation

D Voriconazole

E Itraconazole

Explanation

B Surgical resection

Pulmonary aspergilloma is defined as a conglomeration of intertwined *Aspergillus* hyphae, fibrin, mucus and cellular debris within a pulmonary cavity or an ectatic bronchus. Patients usually have an underlying chronic lung condition. Surgical resection is curative in patients with aspergilloma who have an adequate reserve of lung function to enable them to undergo surgery; pre-existing lung disease may preclude thoracotomy. A preoperative FEV1 of >1.5 l for lobectomy and >2.0 l for pneumonectomy generally indicate suitability for resection procedures on the lungs. Postoperative complications include haemorrhage, bronchopleural fistulae and *Aspergillus* infection of the pleural space.

C Bronchial artery embolisation

Where there is life-threatening haemoptysis, arterial embolisation may be another potential option.

A IV amphotericin

Amphotericin is not first-line treatment for aspergilloma. Amphotericin is associated with febrile reactions, kidney disease and electrolyte disturbances.

D Voriconazole

Voriconazole is a triazole antifungal medication that can be used to treat invasive aspergillosis, which may occur in immunocompromised patients. Visual disturbance is a reported side effect of voriconazole, and thus patients taking voriconazole should be advised against driving at night or other potentially hazardous tasks.

E Itraconazole

Itraconazole is a triazole antifungal medication. In patients whose lung function is too poor to tolerate surgical resection, long-term itraconazole therapy may be effective in reducing symptoms of haemoptysis and actually leading to resolution of the aspergilloma lesion in up to 60% of patients.

For Further Reference:

Walsh et al. Treatment of aspergillosis: Clinical Practice Guidelines of the Infectious Diseases Society of America. *Clin Infect Dis* 2008; 46(3):327–360. doi: 10.1086/525258

Gould and Pearce. Assessment of suitability for lung resection. *Contin Educ Anaesth Crit Care Pain* 2006; 6(3):97–100. doi: 10.1093/bjaceaccp/mkl016

For Further Reference: Walsh et al. Treatment of aspergillosis: Clinical Practice Guidelines of the Infectious Diseases Society of America. Clin Infect Dis 2008; 46(3):327–360. doi: 10.1086/525258 Gould and Pearce. Assessment of suitability for lung resection. Contin Educ Anaesth Crit Care Pain 2006; 6(3):97–100. doi: 10.1093/bjaceaccp/mkl016

21247

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Scott Weingart. EMCrit Podcast 199 – Management of Massive Hemoptysis with Oren Friedman. EMCrit Blog. Published ...
emcrit.org/podcasts/massive-hemoptysis/.
(https://emcrit.org/podcasts/massive-hemoptysis/.)

A 55-year-old dairy farmer attends the Respiratory Clinic with shortness of breath, chronic cough, lethargy and night sweats. He has a history of type II diabetes, for which he takes metformin. Apparently, the farm has had financial troubles over the past few months because of the need to cull approximately 50% of the herd due to tuberculosis (TB).

On examination, his BP is 135/75 mmHg and his heart rate is 75 bpm and regular. His BMI is 35 kg/m^2. He has wheeze on auscultation of the chest, as well as a few scattered coarse crackles.

Pulmonary function testing (PFT) reveals a mild restrictive defect.

His CXR is shown below:



What is the most likely diagnosis?

Your answer was correct

A Allergic bronchopulmonary aspergillosis

B Asthma

C Idiopathic pulmonary fibrosis

D Miliary TB

E Obesity hypoventilation syndrome

Explanation

D Miliary TB

This patient has symptoms of chronic lung disease, and the CXR shows bilateral interstitial infiltrates. Given the fact that the patient has been exposed to TB, this is most likely to indicate miliary TB. The diagnosis should be confirmed with at least three sputum samples, and given the extent of the infection, contact tracing and testing are essential.

Image source: https://commons.wikimedia.org/wiki/File:Chest_radiograph_of_miliary_tuberculosis_1.jpg

A Allergic bronchopulmonary aspergillosis

This leads to an obstructive picture on PFT and symptoms consistent with poorly controlled asthma. Patchy shadowing is usually apparent on the CXR rather than the nodular appearance seen in this patient.

B Asthma

Asthma is more likely to be associated with a normal appearance on the CXR and would not fit with other constitutional symptoms of infection, including lethargy and night sweats seen here.

C Idiopathic pulmonary fibrosis

This would be associated with a restrictive defect on PFT, although coarse crepitations and nodular appearance on the CXR would not be expected. Patchy shadowing on the chest X-ray and inspiratory crepitations would be expected.

E Obesity hypoventilation syndrome

This is associated with a mild restrictive defect on PFT but not with nodular shadowing on the CXR, nor with the auscultation findings seen in this patient.

39656

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 72-year-old man comes to the Emergency Department some 48 hours after starting a course of amoxicillin from his General Practitioner for right lower lobe pneumonia. His condition has deteriorated with persistent fever, a cough productive of purulent, blood stained sputum and diarrhoea. Apparently, he has recently returned from a holiday to Portugal and became ill over the last few days on holiday.

On examination, his blood pressure is 123/85 mmHg; pulse is 85 bpm and regular, his temperature is 38.9 °C. There are signs of right lower lobe consolidation on auscultation of the chest. His abdomen is soft although there is mild tenderness on palpation. He is mildly confused.

Investigations:

Investigation	Result	Normal value
Haemoglobin (Hb)	129 g/l	135–175 g/l
White cell count (WCC)	11.7×10^9/l	$4–11 \times 10^9$/l
Platelets (PLT)	181×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	131 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	103 μmol/l	50–120 μmol/l
C-reactive protein (CRP)	189 mg/l	0–10 mg/l
Alanine aminotransferase (ALT)	95 IU/l	5–30 IU/l
Alkaline phosphatase (ALP)	105 IU/l	30–130 IU/l
Bilirubin	12 μmol/l	2–17 μmol/l
Chest X-ray	Right lower lobe consolidation	

Which of the following antibiotics is appropriate for treating his underlying condition?

Your answer was incorrect

A Ceftriaxone

B Co-amoxiclav

C Meropenem

D Flucloxacillin

E Levofloxacin

Explanation

E Levofloxacin

The suspicion here is that this patient has an atypical pneumonia, potentially Legionnaire’s disease as characterised by confusion, diarrhoea, hyponatraemia and abnormal liver function tests, coupled with the lack of response to amoxicillin. The treatment of choice for Legionnaire’s disease includes modern macrolides including clarithromycin and azithromycin and quinolones such as levofloxacin. Given the side-effects of levofloxacin, including tendon rupture, clarithromycin would seem a sensible first line choice. Should treatment fail, levofloxacin or a broader spectrum agent such as meropenem would be next in line.

D Flucloxacillin

Flucloxacillin may be an appropriate choice for suspected staphylococci pneumonia. This typically occurs post influenzae or measles and causes cavitation on the chest X-ray.

A Ceftriaxone

Ceftriaxone is rarely used in patients with pneumonia given its antimicrobial spectrum. This patient is most likely suffering from an atypical pneumonia; therefore, ceftriaxone is not clinically indicated here.

B Co-amoxiclav

Although co-amoxiclav covers a wider spectrum of organisms than amoxicillin, the clinical scenario indicates a possible atypical pneumonia and therefore requires macrolide cover.

C Meropenem

While carbapenems such as meropenem certainly have efficacy against legionella pneumonia, sensible antibiotic stewardship should be practised here, and the patient should be given a chance to respond to an appropriate antibiotic with a narrower spectrum of activity. Should clarithromycin fail, meropenem would be a reasonable alternative.

39663

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 50-year-old woman presents to the Gastroenterology Clinic with a 6-month history of dysphagia, dyspepsia and occasional regurgitation of food. She reports that sometimes food is difficult to swallow, and she feels the food is stuck in her throat until she vomits it back up. Her past medical history includes asthma and allergic rhinitis.

She is referred for an oesophagogastroduodenoscopy (OGD) that shows generalised oesophageal oedema with widespread white exudates and oesophageal rings.

Given the likely diagnosis, what is the recommended first-line management for this patient?

Your answer was incorrect

A Budesonide

B Hydrocortisone

C Nystatin

D Oesophageal dilation

E Omeprazole

Explanation

A Budesonide

This patient has a likely diagnosis of eosinophilic oesophagitis given the history of repeated dysphagia, dyspepsia and regurgitation of food products. Eosinophilic oesophagitis is an allergen-mediated condition characterised by oesophageal dysfunction, including dyspepsia, dysphagia, vomiting and abdominal pain along with eosinophilic infiltration in the oesophageal epithelium on biopsy. The history of allergic rhinitis and asthma makes eosinophilic-driven conditions such as eosinophilic oesophagitis more likely. Biopsy and OGD are necessary for diagnosis and the findings include oesophageal oedema, oesophageal rings and strictures, with white plaques or exudates representing eosinophilic micro-abscesses. The first-line treatment includes topical corticosteroids such as orodispersible budesonide. Current guidelines recommend continuing treatment with topical corticosteroids for eight weeks, followed by a repeat OGD to assess response.

C Nystatin

Nystatin is used in the treatment of oropharyngeal thrush. It is not absorbed by the gastrointestinal tract and is, therefore, given as an oral suspension and topical delivery. Although this patient has white plaques on endoscopy, this likely represents oesophageal micro-abscesses due to eosinophilic infiltration, and these will not respond to nystatin. Rather, topical corticosteroids are the first-line treatment.

B Hydrocortisone

Systemic corticosteroid use is not recommended in the management of eosinophilic oesophagitis unless rapid relief of severe dysphagia that is causing an impairment of nutrition or hydration is present (eg weight loss, dehydration). It can also be considered when other treatments have failed. This does not seem to be the case here.

D Oesophageal dilation

Oesophageal dilatation is required for patients with severely impaired oesophageal stenosis. There is no mention in this patient's OGD report that there is significant stricture formation or stenosis making oesophageal dilatation an unlikely first-line step in this patient's management.

E Omeprazole

Proton-pump inhibitors (PPIs) such as omeprazole may provide some symptomatic relief with eosinophilic oesophagitis and, in fact, sometimes both conditions may be present. However, PPIs are not the mainstay of treatment for eosinophilic oesophagitis and are not recommended as the first-line for this patient.

73485

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219

Responses Total:	301
Responses - % Correct:	27%

External Links

NICE CKS- Budesonide orodispersible
nice.org.uk/guidance/ta708
(https://www.nice.org.uk/guidance/ta708)

A 32-year-old man is referred to the Gastroenterology Clinic by his General Practitioner with an 8-month history of dyspepsia and dysphagia that has failed to respond to oral proton-pump-inhibitor therapy. In the last week he presented to the Emergency Department with an acute food bolus while eating a lamb biryani but reported that this cleared after drinking fizzy cola, and he was discharged home. He is otherwise well in himself and denies weight loss. His past medical history includes eczema and childhood asthma.

What is the most appropriate investigation to confirm the likely diagnosis?

Your answer was correct

A Barium swallow

B Laryngoscopy

C Oesophagogastroduodenoscopy and biopsy

D Oesophageal manometry

E Urea breath test

Explanation

C Oesophagogastroduodenoscopy and biopsy

This patient has a likely diagnosis of eosinophilic oesophagitis, given the history of prolonged dysphagia and dyspepsia that has failed to respond to proton-pump-inhibitor therapy. The recent presentation with an acute food bolus is also suggestive of the condition. Eosinophilic oesophagitis is an allergen-mediated condition characterised by oesophageal dysfunction including dyspepsia, dysphagia, vomiting and abdominal pain along with eosinophilic infiltration in the oesophageal epithelium on biopsy.

The history of eczema and asthma makes eosinophilic-driven conditions such as eosinophilic oesophagitis more likely. Oesophagogastroduodenoscopy (OGD) and biopsy are necessary for diagnosis, and findings include oesophageal oedema, oesophageal rings and strictures, with white plaques or exudates representing eosinophilic micro-abscesses. Multiple biopsies are taken with a peak eosinophil count and/or 15 eosinophils per high-power microscopy field being diagnostic for the condition.

A Barium swallow

Barium swallow is not used in the diagnosis of eosinophilic oesophagitis but is used in the diagnosis of pharyngeal pouch and achalasia. Achalasia is characterised by motor dysfunction of the oesophagus that causes symptoms of dysphagia and regurgitation due to loss of oesophageal peristalsis. Meanwhile, a pharyngeal pouch causes symptoms of dysphagia, regurgitation, chronic cough, aspiration and halitosis and is most commonly seen in men aged above 70 years. These diagnoses do not completely fit with the presentation of this patient.

B Laryngoscopy

Laryngoscopy is used in the diagnosis of laryngeal disease (eg laryngeal carcinoma). While this can present with dysphagia, the symptoms of dyspepsia and food bolus obstruction are less common. A laryngoscopy would not be a suitable investigation for this patient, as it would not provide adequate visualisation of the oesophagus, to ascertain the cause of this presentation.

D Oesophageal manometry

Oesophageal manometry is used in the investigation of achalasia which is characterised by motor dysfunction of the oesophagus, causing dysphagia and regurgitation due to loss of oesophageal peristalsis. High-resolution manometry is now the gold standard for the diagnosis and findings of incomplete relaxation of the lower oesophageal sphincter and aperistalsis being key criteria for diagnosis. However, the history of dyspepsia and food bolus obstruction in conjunction with a background of eczema and asthma make eosinophilic oesophagitis the more likely diagnosis for which oesophageal manometry is not helpful.

E Urea breath test

A urea breath test is used in the diagnosis of Helicobacter pylori, a Gram-negative organism capable of causing peptic ulcer disease and dyspepsia. However, dysphagia and acute food bolus obstruction are more suggestive of other diagnoses.

73486

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 77-year-old woman was referred to the Respiratory Outpatient Clinic with increasing shortness of breath. Her chest X-ray (CXR) is shown below.



What operation has she had?

Your answer was correct

A Left upper lobectomy

B Left pneumonectomy

C Thoracoplasty

D Phrenic crush

E Plombage

Explanation

C Thoracoplasty

This question is a 'spot diagnosis'. Clues that this is a thoracoplasty and not any of the alternative options are that there is evidence of rib resection in conjunction with loss of lung volume. The chest X-ray shows a thoracoplasty. This lady had tuberculosis in the 1940s, prior to the use of anti-tuberculosis chemotherapy. She had a thoracoplasty, which was one of the main surgical treatments for tuberculosis between 1930 and 1955. Several ribs were resected, which reduced the thoracic volume and collapsed the underlying lung. The aim was to close the tuberculous cavity and 'rest the lung'. Control of pulmonary tuberculosis and survival was good. However, patients were left with severe chest deformity and the associated respiratory compromise, a restrictive defect.

Image courtesy of Dr Ian Bickle.

A Left upper lobectomy

Upper lobectomy would only be associated with loss of upper lung volume.

B Left pneumonectomy

This would not be associated with the bony changes seen here.

D Phrenic crush

Phrenic crush is associated with elevated hemidiaphragm rather than the extensive bony resection seen here.

E Plombage

The CXR would show material inserted into the chest cavity to promote lung collapse.

70126

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Tuberculosis

Media

Expanded explanations

Flashcards

A 31-year-old woman presented to the Respiratory Outpatients Department with a 3-year history of increasing shortness of breath and cough. She had required numerous courses of antibiotics in the last few years. She was a smoker of 5/day for 5 years but gave up 4 years ago. Respiratory examination revealed a hyper-inflated chest and a mild wheeze throughout.

Lung function tests:

FEV_1	50% predicted
FVC	80% predicted
TLCO	68% predicted
KCO	71% predicted
RV:TLC	Increased

What is the most likely diagnosis?

Your answer was incorrect

A COPD related to smoking

B Asthma

C Cystic fibrosis

D Bronchiectasis

E Alpha-1-antitrypsin deficiency

Explanation

E Alpha-1-antitrypsin deficiency

Alpha-1-antitrypsin deficiency is a rare autosomal recessive disease associated with the development of premature panlobular emphysema, most severe in the basal areas of the lungs.Emphysema is thought to result from an imbalance in the lung between neutrophil elastase (which destroys elastin) and the elastase inhibitor α1-antitrypsin (which protects against the proteolytic degradation by elastin). The decline in lung function is accelerated in smokers.

Patients usually present with progressive increasing shortness of breath and weight loss. Corpulmonale and polycythaemia occur later in the course of the disease. A chest X-ray usually shows bilateral bibasal emphysema. Lung function tests show an obstructive picture and the RV:TLC ratio is raised, indicating gas

trapping. Transfer factor will be reduced. There are a number of different phenotypes depending on the serum level of α1-antitrypsin:
Pi – protease inhibitor
M – normal allele
Z, X and null are disease alleles

Phenotype	α1-Antitrypsin level	Disease
PiMM	Normal	
PiMS	80%	
PiMZ	60%	
PiSS	60%	
PiSZ	40%	Liver cirrhosis/premature emphysema
Pi Null Null	Nil	Premature emphysema

Treatment is similar to that of asthma. Alpha1-antitrypsin replacement can be given weekly or monthly and is recommended for those with very low serum levels and abnormal lung function. Lung transplantation should be considered in advanced disease. Patients should be encouraged to stop smoking.

C Cystic fibrosis

Respiratory problems in CF would usually be present since infancy. She has no other features of CF such as pancreatic insufficiency, sinus disease, liver disease or diabetes.

A COPD related to smoking

Her disease severity coupled with her minimal smoking history makes α1-antitrypsin deficiency more likely.

B Asthma

The lung function pattern and history suggesting lack of variability in symptoms is more in keeping with emphysema.

D Bronchiectasis

The gas trapping and reduced gas transfer are more in keeping with emphysema in this history.

70138

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

An acutely unwell 62-year-old man is brought to the Emergency Department by his wife. They had returned from holiday to Spain three days ago. Apart from some mild flu-like symptoms, he had been otherwise well on returning to the United Kingdom. Over the last 24 hours, he had developed a severe headache, violent shivers and muscle pains. He also had some abdominal pain, with several episodes of diarrhoea. He had no significant past medical history and was on no medication. He had consumed some seafood on holiday, but his wife had eaten the same dish and had no symptoms.

On examination, he was confused and agitated, with a fever of 40 °C. His respiratory rate was 32 bpm, pulse 110 bpm and blood pressure 100/62 mmHg. His heart sounds were normal, and he had left basal crepitations on chest auscultation. His abdomen was soft and non-specifically tender, with scanty bowel sounds. There was no focal neurology.

Investigations reveal the following:

Investigation	**Result**	**Normal value**
Haemoglobin (Hb)	136 g/l	135–175 g/l
White cell count (WCC)	12.0 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	380 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	122 mmol/l	135–145 mmol/l
Potassium (K^+)	3.2 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	170 µmol/l	50–120 µmol/l
Neutrophils	10.2 × 10^9/l	2.5–7.58 × 10^9/l
Lymphocytes	0.5 × 10^9/l	1.5–3.5 × 10^9/l
Monocytes	0.8 × 10^9/l	0.0–0.8 × 10^9/l
Eosinophils	0.4 × 10^9/l	0.04–0.4 × 10^9/l
Basophils	0.1 × 10^9/l	< 0.1 × 10^9/l
Gamma-glutamyltransferase (GGT)	65 IU/l	5–30 IU/l
Total calcium (Ca^{2+})	2.06 mmol/l	2.20–2.60 mmol/l
Bilirubin	16 µmol/l	2–17 µmol/l
Alanine aminotransferase (ALT)	92 IU/l	5–30 IU/l
Albumin	30 g/l	35–55 g/l
Alkaline phosphatase (ALP)	98 IU/l	30–130 IU/l
Urea	7.6 mmol/l	2.5–6.5 mmol/l
Electrocardiogram (ECG)	Sinus tachycardia	

Urinalysis	Blood+++, protein+	
Chest X-ray (CXR)	Shown below	



By Antonella GrottolaComments to Author , Fabio Forghieri, Marisa Meacci, Anna Fabio, Lorena Pozzi, Patrizia Marchegiano, Mauro Codeluppi, Monica Morselli, Leonardo Potenza, Ambra Paolini, Valeria Coluccio, Mario Luppi, Fabio Rumpianesi, and Monica Pecorari [Public domain], via Wikimedia Commons

What is the most likely causative organism?

Your answer was incorrect

A *Legionella pneumophila*

B *Mycoplasma pneumoniae*

C *Coxiella burnetii* (Q fever)

D *Salmonella typhi*

E *Chlamydia pneumoniae*

Explanation

A *Legionella pneumophila*

This man has the typical features of legionnaires' disease. *Legionella* infection is said to be the cause of between 1 and 5% of adult pneumonias. The organism is a small coccobacillus, which is not seen on Gram staining and can only be cultured on special media. Infection is spread by the aerosol route and is usually acquired from contaminated air-conditioning or shower systems in institutions such as hotels and hospitals. Investigations may show hyponatraemia, normal or slightly elevated WCC with lymphopenia, hypoalbuminaemia and raised liver enzymes. Chest X-ray may show either unilobar or multilobar involvement and/or a pleural effusion. Haematuria and kidney disease can occur. Diagnosis is made by either a fourfold increase in antibody titre or a highly specific urine antigen test. Macrolides or quinolones with rifampicin are used in combination for patients with severe infection. Mortality is 15–30%, particularly in the elderly.

C *Coxiella burnetii* (Q fever)

Coxiella burnetii is a cause of Q fever, with abrupt onset of fever, malaise, profuse sweating, severe headaches, muscle and joint pains, loss of appetite, dry cough, pleuritic chest pain and confusion.

B *Mycoplasma pneumoniae*

Mycoplasma is associated with a dry cough and pleuritic chest pain, and in some patients, it may be associated with haemolysis. It is less likely than *Legionella* to be associated with gastrointestinal disturbance and is treated with macrolides as the first-line intervention.

D *Salmonella typhi*

Typhoid fever presents with marked pyrexia and abdominal pain, with bradycardia, rather than tachycardia, as seen here.

E *Chlamydia pneumoniae*

Chlamydia presents with nasal stuffiness, a low-grade fever, laryngitis and a slowly worsening cough that can last for a period of weeks.

9144

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 51-year-old man, while an inpatient for a routine hernia operation, was noted to have a pneumothorax on his pre-op chest X-ray. He has a history of chronic bronchitis, and a 50 pack-year smoking history. His chest X-ray shows a unilateral pneumothorax of 2.2 cm. He is not breathless, but has chest pain.

Which of the following would be the ideal management for the clinical scenario described?

Your answer was incorrect

A You decide to aspirate the pneumothorax with a 16G cannula

B You decide to insert a 14 French gauge chest drain

C You decide to give the patient 10 litres of oxygen by mask

D You decide to admit the patient for observation only

E You decide to discharge the patient to outpatient ambulatory clinic in 2 weeks

Explanation

B You decide to insert a 14 French gauge chest drain

This patient has a secondary pneumothorax as he is over the age of 50 and has a significant smoking history. The British Thoracic Guidelines suggest you should insert a chest drain 8-14 Fr if the pneumothorax is greater than 2 cm in size or the patient is breathless. The intrapleural distance should be measured at the level of the hilum.

D You decide to admit the patient for observation only

Admission without intervention can be considered for patients with secondary pneumothoraces where the pneumothorax less than 1 cm in size, and the patient asymptomatic (not breathless).

A You decide to aspirate the pneumothorax with a 16G cannula

An attempt at aspirating a secondary pneumothorax should only be made where the pneumothorax is between 1 and 2 cm, and the patient is not breathless.

C You decide to give the patient 10 litres of oxygen by mask

High-flow oxygen may be used as a treatment option for secondary pneumothoraces less than 1 cm in size, provided you do not suspect the patient is at risk of oxygen sensitivity.

E You decide to discharge the patient to outpatient ambulatory clinic in 2 weeks

Admission without intervention can be considered for patients with secondary pneumothoraces where the pneumothorax less than 1 cm in size, and the patient asymptomatic (not breathless).

7181

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total: 301

Responses - % Correct: 27%

Pneumothorax

Media

Expanded explanations

Flashcards

A 62-year-old man attends for a chest X-ray as part of a medical prior to emigrating to Australia. Unfortunately the chest X-ray reveals a number of rounded opacities predominantly in the left lung field, all less than 3 cm in diameter. He has had 5 kg of unintentional weight loss over the past year but no other history of note.

Which of the following features would be the strongest pointer to malignancy when the chest X-ray is reviewed?

Your answer was incorrect

A Spiculated margin to one of the nodules, with distortion of neighbouring vessels

B Smaller lesion size

C A rounded appearance to the nodules

D No enhancement with contrast on future CT thorax

E Age 62 at presentation

Explanation

A Spiculated margin to one of the nodules, with distortion of neighbouring vessels

A spiculated margin to one of the nodules, particularly when there is distortion of neighbouring vessels, raises the possibility of malignancy. Associated ground-glass shadowing, or the presence of cavitation or pseudo-cavitation within the lesion, increases the chances of malignant disease. CT thorax to further characterise the lesions is indicated.

C A rounded appearance to the nodules

Rounded appearance to the nodules, implying organised growth, is less likely to be associated with an underlying malignancy.

B Smaller lesion size

Lesions larger than 3 cm are more likely to be associated with underlying malignancy.

D No enhancement with contrast on future CT thorax

Enhancement of nodules with contrast on CT thorax increases the likelihood that they are malignant in origin.

E Age 62 at presentation

Older age at presentation and history of smoking are associated with increased likelihood of malignancy. 32434

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 25-year-old man presents with left-sided pleuritic chest pain. It had come on suddenly three days previously. Initially, he thought it was secondary to his workout in the gym. However, since it did not improve and he noticed that he was slightly more breathless than usual, he went to the Emergency Department. He has no significant past medical history. He is a smoker.

On examination, he looked well. He was a tall, thin man and was not short of breath at rest. Examination was unremarkable except for a clicking sound which was synchronous with the heart sounds. CXR was unremarkable.

How would you best manage this patient?

Your answer was incorrect

A Analgesia and discharge with follow-up in the clinic

B Aspiration of chest

C Chest drain insertion

D Non-steroidal anti-inflammatory drugs and admit patient

E Refer to Cardiology Department

Explanation

A Analgesia and discharge with follow-up in the clinic

A synchronous click with the heart sounds is a recognised sign of a small left apical pneumothorax. He is a tall thin man and a smoker, which are both risk factors for developing a pneumothorax. There is a strong association between pneumothoraces and smoking. The most recent British Thoracic Society (BTS) guidelines[1], suggest that if the postero-anterior (PA) chest X-ray is normal and a small pneumothorax is suspected, a lateral decubitus chest X-ray provides added information in up to 14% of cases. Expiratory films add little and are not recommended. The treatment of a primary pneumothorax in a non smoker with no evidence of underlying lung disease <50yrs old with a rim of air of <2 cm and no breathlessness is to discharge and follow up as an outpatient. Although this patient is a smoker there is nothing to suggest underlying lung disease and it is likely that his apical pneumothorax is too small to safely aspirate. Patients should be given analgesia if required and clear instructions to return if their symptoms get any worse.

[1] Management of spontaneous pneumothorax: British Thoracic Society pleural disease guideline 2010 | Thorax (bmj.com)

(https://thorax.bmj.com/content/65/Suppl_2/ii18)

C Chest drain insertion

There is no indication to aspirate a small pneumothorax in a young patient with no lung disease who is not breathless when reviewed.

B Aspiration of chest

There is no clinical indication to admit this patient, although he should be followed up as an outpatient.

D Non-steroidal anti-inflammatory drugs and admit patient

There is no indication insert a chest drain in a small pneumothorax in a young patient with no lung disease who is not breathless when reviewed.

E Refer to Cardiology Department

This would not be helpful in diagnosing pneumothorax.

70120

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BTS Pleural Disease Guideline 2010
brit-thoracic.org.uk/document-library/clinical-information/pleural-disease/pleural-disease-guidelines-2010/appendix-3-spontaneous-pneum...
(https://www.brit-thoracic.org.uk/document-library/clinical-information/pleural-disease/pleural-disease-guidelines-2010/appendix-3-spontaneous-pneumothorax-poster-pleural-disease-guideline/)

A 72-year-old man is admitted to the Intensive Care Unit with post-influenza pneumonia. Apparently, he has been ill for the past week with flu-like symptoms and developed a severe cough productive of purulent sputum over the past 48 hours. He has a past history of ischaemic heart disease, hypertension and type 2 diabetes and is known to be penicillin-allergic. He was also treated with broad-spectrum antibiotics in past. Examination reveals a blood pressure of 105/60 mmHg, a pulse of 95 bpm and regular and a temperature of 38.2 °C. He is intubated and ventilated. There is extensive right-sided consolidation on auscultation of his chest.

Investigations;

Investigations	**Results**	**Normal values**
Haemoglobin (Hb)	129 g/l	135–175 g/l
White cell count (WCC)	15.8×10^9/l	$4–11 \times 10^9$/l
Platelets (PLT)	205×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	135 mmol/l	135–145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine	152 µmol/l	50–120 µmol/l
C-reactive protein (CRP)	240 mg/l	0–10 mg/l

Chest X-ray (CXR): Extensive right sided consolidation, some left sided basal consolidation.

Which of the following is the most appropriate intervention?

Your answer was incorrect

A Clarithromycin

B Doxycycline

C Flucloxacillin

D Linezolid

E Vancomycin

Explanation

D Linezolid

This patient has severe staphylococcal pneumonia which can occur after infection with influenza. Studies have shown that linezolid is effective in treating both non-meticillin-resistant and meticillin-resistant staphylococcal pneumonia. One study of linezolid versus vancomycin showed that linezolid was associated with more rapid clearance of pneumonia, although no significant difference was seen in 60-day mortality. Moreover, this patient also has renal impairment and hence it's better to avoid vancomycin here.

B Doxycycline

Doxycycline is less effective than linezolid in the treatment of staphylococcal pneumonia.

A Clarithromycin

Clarithromycin is less effective than linezolid in the treatment of staphylococcal pneumonia.

C Flucloxacillin

This patient has a penicillin allergy. Therefore, he would be at risk for developing anaphylaxis if flucloxacillin were to be administered. Flucloxacillin would be the first-line agent for staphylococcal pneumonia in a patient with no allergies.

E Vancomycin

As discussed above, one study of linezolid versus vancomycin showed that linezolid was associated with more rapid clearance of pneumonia, although no significant difference was seen in 60-day mortality. It is therefore reasonable to consider linezolid as first line, as opposed to vancomycin.

38214

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 46-year-old man is admitted to the Emergency Department complaining of chest pain, most marked on the left. He smokes ten cigarettes a day, drinks 3–4 pints of beer each week and has a history of recent travel.

On examination, following a 1-litre fluid bolus, his BP is 80/60 mmHg and his heart rate is 105 bpm and regular. His jugular venous pressure (JVP) is markedly elevated. Saturation is 92% on 40% oxygen by mask. His chest is clear. The ECG shows significant tachycardia and deep T wave inversion in leads V2–V4. His troponin level is 88 ng/ml (normal range under 14 ng/ml). Unfractioned heparin has already been commenced.

Which of the following is the most appropriate next step?

Your answer was incorrect

A Aspirin 300mg

B Apixaban

C Pericardiocentesis

D Synchronised DC cardioversion

E Thrombolysis

Explanation

E Thrombolysis

The patient most likely has a massive pulmonary embolism (PE) and is at risk of an imminent cardiac arrest. As such, thrombolysis is an appropriate next step. The patient is significantly hypoxic, and his BP has not improved in response to a fluid bolus. According to guidelines, indications for thrombolysis in massive PE include:

- a systolic pressure of less than 90 mmHg or a drop of over 40 mmHg for longer than 15 minutes if not caused by a rhythm disturbance
- new-onset right heart failure, as evidenced by raised JVP, loud P2 and ECG abnormalities where there is evidence of thrombus formation on echocardiography.

C Pericardiocentesis

Pericardiocentesis is the treatment for a cardiac tamponade. Similarly to this patient’s presentation, cardiac tamponade can present with raised JVP and hypotension. However, in the context of a recent flight, a PE is more likely.

A Aspirin 300mg

Antiplatelet agents do not improve the prognosis in massive pulmonary embolism (PE). Furthermore, they do not even have any useful impact as prophylaxis for PE. Instead, a loading dose of aspirin is used as an initial treatment for life-threatening myocardial or cerebral ischaemia. The patient's ECG and troponin changes likely demonstrate right heart strain rather than myocardial ischaemia.

B Apixaban

Apixaban monotherapy is appropriate for patients presenting with a PE, where there is haemodynamic stability. Here, with haemodynamic instability, thrombolysis is indicated.

D Synchronised DC cardioversion

Synchronised DC cardioversion is used as an emergency treatment option for tachyarrhythmia with adverse features (eg ischaemia, syncope, heart failure, hypovolaemia). Although this patient is tachycardic and hypotensive, it is more likely that this is secondary to a PE, given the history of recent travel, chest pain and hypoxia. Therefore, urgent thrombolysis is the preferred treatment of choice.

39759

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect: 219

Responses Total:	301
Responses - % Correct:	27%

External Links

NICE Guidelines Pulmonary Embolism
cks.nice.org.uk/pulmonary-embolism#!scenario:1
(https://cks.nice.org.uk/pulmonary-embolism#!scenario:1)

A 62-year-old man is admitted to hospital with a 3-day history of dyspnoea, pleuritic chest pain and several episodes of haemoptysis.

Before this, he described several months of constant rhinorrhoea, with some nasal crusting, several large epistaxes and constant pain below his eyes. More recently, he had noticed some double vision and swelling of his right eye.

On examination, he had low-grade pyrexia and was normotensive. There was some nasal mucosal ulceration and right-sided proptosis. Cardiovascular examination was unremarkable. He had localised areas of crepitations throughout both lung fields. His abdomen was soft and non-tender, with no masses. There were no focal neurological signs or skin lesions.

Investigation results are below:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	100 g/l	135 - 175 g/l
White cell count (WCC)	12.0×10^9/l	$4.0 – 11.0 \times 10^9$/l
Platelets (PLT)	390×10^9/l	$150 - 400 \times 10^9$/l
Mean corpuscular volume (MCV)	90 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	80 mm/h	1 - 20 mm/h
Sodium (Na^+)	142 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.7 mmol/l	3.5 - 5.0 mmol/l
Urea	22.0 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	620 µmol/l	50 - 120 µmol/l
C-reactive protein	238 mg/L	< 10 mg/l
Bilirubin	15 µmol/l	1.7 - 20.5 µmol/l
Alkaline phosphatase (ALP)	88 u/l	20 - 140 u/l
Aspartate aminotransferase (AST)	32 u/l	10 - 40 u/l
Calcium (Ca^{2+})	2.42 mmol/l	2.2 - 2.7 mmol/l
Gamma-glutamyl transferase (GGT)	48 u/l	9 - 48 u/l
Urinalysis	Blood +++ protein ++	
Chest X-ray	Multiple large cavitating nodules throughout both lung fields	
Electrocardiogram (ECG)	Normal sinus rhythm, no acute changes	

What would be the most useful investigation in pointing to the diagnosis if positive?

Your answer was correct

A High-resolution computed tomography (CT) scan of the thorax

B Serum anti-glomerular basement membrane (anti-GBM) antibodies

C Serum antineutrophil cytoplasmic antibodies

D Serum antinuclear antibodies

E Urine microscopy

Explanation

C Serum antineutrophil cytoplasmic antibodies

This patient has the typical features of Granulomatosis with polyangiitis (GPA), which is a necrotising granulomatous arteritis of the upper and lower respiratory tract and kidney. It is rare, with an incidence of 5–10 per million. It is a multisystem disease and may affect the eyes, skin, joints, heart and nervous system. Pulmonary involvement is seen in 95% of cases, and renal involvement in 85% of cases. A positive serum antineutrophil cytoplasmic antibody (ANCA) is present in > 90% of cases of GPA and strongly supports the diagnosis. Renal biopsy is sometimes required.

Treatment is with steroids and cyclophosphamide. Untreated, 1-year mortality is 80%, but with appropriate therapy, remission can be achieved in up to 90% of patients.

A High-resolution computed tomography (CT) scan of the thorax

This may show the cavitating lung lesions in more detail but will not aid in the diagnosis.

B Serum anti-glomerular basement membrane (anti-GBM) antibodies

This scenario is not anti-GBM disease; therefore, these antibodies will be normal. Anti-GBM disease typically causes macroscopic haematuria & proteinuria with mild renal impairment along with alveolar haemorrhage.

D Serum antinuclear antibodies

Serum antinuclear antibodies will not aid in the diagnosis.

E Urine microscopy

Urine microscopy may show blood and protein; however, it is non-specific for the diagnosis.

71306

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 35-year-old woman with a 15-year smoking history and a diagnosis of myasthenia gravis is referred by the neurologists for review. She has complained of progressive dyspnoea over the past few months. On examination her BP is 123/82, with pulse 67 and regular. You note she seems extremely fatigued, with obvious partial ptosis and proximal muscle weakness. The weakness increases with repetitive testing. Chest X-ray reveals a mediastinal mass and detailed pulmonary function testing reveals evidence of extrinsic obstruction.

Which of the following diagnoses is most likely given this clinical history?

Your answer was incorrect

A Multinodular goitre

B Thymoma

C Schwannoma

D Neurofibroma

E Bronchial carcinoma

Explanation

B Thymoma

Given the clinical history of myasthenia gravis and a mediastinal mass, thymoma represents the most likely diagnosis in this case (30–40% of thymoma cases have co-existent myasthenia gravis). Thymoma has equal sex distribution and rarely presents below the age of 20 years. Removal of the thymoma can significantly improve symptoms of myasthenia and is the next logical step once the diagnosis is confirmed in this patient.

C Schwannoma

Myasthenia gravis can be associated with multiple schwannomas, although these are not associated with extrinsic airways obstruction.

A Multinodular goitre

Multinodular goitre is associated with extrinsic airways obstruction, although it isn’t associated with myasthenia gravis.

D Neurofibroma

Superficial plexiform neurofibromas have been reported to occur in patients with myasthenia gravis, and this may occur because of inflammation around the neuromuscular junction.

E Bronchial carcinoma

Bronchial carcinoma is associated with Lambert-Eaton myasthenic syndrome, which is associated with potentiation on repeated muscle testing rather than exacerbated weakness as seen here.

32437

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Thyroid Cancer

Media

Expanded explanations

A 63-year-old man is reviewed in the Respiratory Clinic some six weeks after his latest admission with an exacerbation of COPD. He is currently managed by his GP with salmeterol inhaled BD and PRN Salbutamol, and has finished a reducing course of oral corticosteroids in the past few weeks. He is still significantly short of breath and is unable to walk more than 100 metres without stopping despite having given up smoking. He had a diagnosis of asthma for which he took inhalers until he was in his mid thirties.

On examination, his BP is 155/85 mmHg, and heart rate is 70 bpm, (atrial fibrillation), auscultation of the chest reveals bilateral poor air entry and fine wheeze consistent with emphysema. O$_2$ saturation is 93% on air.

Investigations;

Hb	131 g/l
WCC	7.4 x10^9/l
PLT	181 x10^9/l
Na$^+$	137 mmol/l
K$^+$	4.5 mmol/l
Creatinine	122 micromol/l
Glucose	5.6 mmol/l
ECG	inferior Q waves, pulse 70 atrial fibrillation
CXR	Evidence of emphysema
FEV$_1$/FVC	0.65
FEV$_1$	35% of predicted

Which of the following is the most appropriate add-on therapy, according to NICE guidelines?

Your answer was incorrect

A Add ipratropium bromide PRN

B Add theophylline tablets

C Add beclomethasone

D Add tiotropium

E Start home oxygen therapy

Explanation

C Add beclomethasone

The NICE guidelines recommend for patients with COPD and ‘asthmatic features’ (previous diagnosis of asthma, high eosinophil count, more than 400ml variation in FEV1 over time or at least 20% diurnal variation in peak expiratory flow) then treatment with a long-acting β2-agonists (LABA) and inhaled corticosteroid (ICS) should be considered. There is an increased risk of pneumonia in patients who take ICS. The next step would be to add a long-acting muscarinic antagonist (LAMA).

D Add tiotropium

Tiotropium, a LAMA, would be indicated along with a LABA in patients with no features of steroid responsiveness after a short-acting bronchodilator or short-acting muscarinic antagonist.

A Add ipratropium bromide PRN

Short-acting bronchodilators or muscarinic antagonists are recommended in the NICE guidelines as a first-line therapy, although they are not both required. Additionally, they do not impact prognosis.

B Add theophylline tablets

Add Theophylline tablets, is incorrect because whilst they can impact on symptoms, they do not impact on prognosis and may increase the risk of fast AF, particularly in a patient with rate controlled AF, as here.

E Start home oxygen therapy

Home oxygen therapy is not currently required; the COPD therapy is not currently maximised, given his measured O2 saturation, it is unlikely he is hypoxic on air, and there are no signs of right heart failure.

40432

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

NICE - COPD management
nice.org.uk/guidance/ng115/resources/visual-summary-treatment-algorithm-pdf-6604261741
(https://www.nice.org.uk/guidance/ng115/resources/visual-summary-treatment-algorithm-pdf-6604261741)

NICE - Chronic obstructive pulmonary disease in over 16s: diagnosis and management
nice.org.uk/guidance/ng115/chapter/Recommendations#managing-stable-copd
(https://www.nice.org.uk/guidance/ng115/chapter/Recommendations#managing-stable-copd)

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 55-year-old man presents to his GP, with his wife complaining of 3–4 months of increased shortness of breath, irritability and daytime somnolence. The patient is a known smoker of one pack per day. He reports no other medical problems and is not on any medication. Clinical examination shows the patient to have a BMI of 22, with mild peripheral oedema and cyanosis. The patient is tachypnoeic. Examination of the chest shows hyperinflation, with the use of accessory muscles of respiration and quiet breath sounds. Pulse oximetry shows a pa(O_2) of 94%. Appropriate investigations are initiated for the patient, including lung function tests which show non-reversible obstruction and air trapping. Additionally, the patient is referred to the Sleep Disorders Unit of the nearest specialised Ear, Nose and Throat hospital, where arterial blood gases show a chronic respiratory acidosis.

The results of the sleep study are below.



His chest x-ray is shown below.



What is the most likely diagnosis for this patient’s symptoms?

Your answer was correct

A Obstructive sleep apnoea

B Pickwickian syndrome

C Chronic obstructive pulmonary disease (COPD) and nocturnal hypoventilation

D Chronic obstructive pulmonary disease (COPD) and simple snoring

E Narcolepsy

Explanation

C Chronic obstructive pulmonary disease (COPD) and nocturnal hypoventilation

The combination of symptoms, obstructive lung function tests and chest x-ray showing COPD is reached as a diagnosis here hyperexpansion and increased lung markings. COPD patients commonly suffer from nocturnal oxygen desaturation that is not usually caused by sleep apnoea. Sleep-related hypoxaemia can result in the patient reporting morning headaches and daytime somnolence. The diagnosis is made via overnight home monitoring with a pulse oximeter, which shows dramatic falls in saturation levels that occur during REM sleep.

Image source: https://radiopaedia.org/cases/chronic-obstructive-pulmonary-disease-marked-hyperinflation

A Obstructive sleep apnoea

Obstructive sleep apnoea is a disorder characterised by obstructive apnoeas, daytime symptoms secondary to reduced sleep and signs of disturbed sleep such as snoring. It typically occurs in obese patients with craniofacial or upper airway soft tissue abnormalities. Treatment is with CPAP.

B Pickwickian syndrome

Pickwickian syndrome is the combination of severe obesity, a plethoric face and obstructive sleep apnoea causing hypoxia and hypercapnia, resulting in marked daytime somnolence. It is named after the ‘fat and red-faced boy in a state of somnolency’ as described in the novel *The Pickwick Papers* by Charles Dickens.

D Chronic obstructive pulmonary disease (COPD) and simple snoring

This patient has evidence of disturbed sleep, and the study demonstrates hypoxia during sleep, meaning that this is more than simple snoring.

E Narcolepsy

Narcolepsy is a neurological disorder characterised by a sudden and uncontrollable, though often brief, compulsion to sleep, sometimes accompanied by cataplexy and sleep paralysis.

6807

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 73-year-old man is sent to the Emergency Department by ambulance after a GP home visit. The GP found him to be extremely short of breath with a cough productive of rusty-coloured sputum. He has herpetic lesions at the corner of his mouth. On examination he has right-sided crackles and bronchial breathing consistent with pneumonia.

You use the British Thoracic Society Community Acquired Pneumonia Severity Score (CURB-65) criteria to assess the severity of infection with respect to the need for HDU/ITU.

Which one of the following features would be consistent with severe pneumonia by CURB-65 criteria?

Your answer was incorrect

A Urea 9.2 mmol/l

B Systolic BP 105 mmHg

C Diastolic BP 65 mmHg

D Presence of X-ray changes

E Raised ESR

Explanation

A Urea 9.2 mmol/l

The British Thoracic Society Community Acquired Pneumonia Severity Score (CURB-65) uses five variables where a point is scored for each one. It is a mortality score.

The variables are:

- confusion (AMT of 8 or less);
- urea >7 mmol/l;
- respiratory rate 30/min or greater;
- systolic blood pressure 90 mmHg or less or diastolic BP 60 mmHg or less;
- age >65

A score of 3 is associated with a risk of mortality of 17% from community-acquired pneumonia; for 5 this rises to a risk of mortality of 57%. These data are extrapolated to give a guide as to whether and where the patient should be admitted: those with a score of 0 or 1 may safely be managed at home, 2 should receive hospital care, while HDU/ITU should be considered for a score of 3 or more. This is a guide and should not replace clinical judgement.

C Diastolic BP 65 mmHg

Again, though this does not fulfil the criteria of CURB-65, it may be of clinical importance. CURB-65 is a guide to likely mortality and is used for this reason as a pointer towards the requirement for ITU/HDU admission. It is not, however, a management guide.

B Systolic BP 105 mmHg

This blood pressure does not fit the criteria, although may be of concern depending on his other clinical observations.

D Presence of X-ray changes

X-ray changes are not involved in these prognostic criteria, though may be of diagnostic and prognostic value.

E Raised ESR

ESR is not routinely measured in community-acquired pneumonia.

32332

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

NICE guidance
nice.org.uk/guidance/cg191/chapter/1-recommendations#community-acquired-pneumonia-2
(https://www.nice.org.uk/guidance/cg191/chapter/1-recommendations#community-acquired-pneumonia-2)

Pneumonia

Media

Expanded explanations

Flashcards

A previously well 23-year-old student presents to the Emergency Department with an acute onset of pleuritic chest pain and dyspnoea. She has just returned from a gap year to Australia and had been well until the morning of admission.

She has no significant past medical history or family history, and her only medication included the contraceptive pill. She is a non-smoker.

On examination, she is acutely distressed, with a respiratory rate of 32 breaths/min, a pulse of 128 bpm and a blood pressure of 98/60 mmHg. She is peripherally cool. Her chest is clear to auscultation, and her heart sounds are normal. The jugular venous pressure (JVP) is elevated. There is no obvious leg swelling.

Investigation results are below:

Investigation	Result	Normal Values
Haemoglobin (Hb)	152 g/l	115 - 155 g/l
White cell count (WCC)	14 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	310 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	136 mmol/l	135–145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5–5.0 mmol/l
Urea	3.2 mmol/l	2.5–6.5 mmol/l
Creatinine (Cr)	90 µmol/l	50–120 µmol/l
Chest X-ray	Normal	

Electrocardiogram is shown below



Arterial blood gases (on 15 l/min oxygen) were:

pH	7.48

pa(O_2)	8.2 kPa
pa(CO_2)	3.6 kPa

Intravenous (IV) heparin is commenced.

Which of the following is the most appropriate initial management?

Your answer was incorrect

A Limited IV crystalloid fluid

B Thrombolysis with streptokinase

C IV dobutamine

D IV doxapram

E IV dopamine

Explanation

A Limited IV crystalloid fluid

The diagnosis is an acute major pulmonary embolism. The ECG shows sinus tachycardia, the commonest ECG finding in such a presentation. Fluids should be given in the first instance to increase the right ventricular filling pressure and maintain cardiac output. Heparinisation should prevent the occurrence of further emboli.

Image source: https://commons.wikimedia.org/wiki/File:Sinustachycardia_(CardioNetworks_ECGpedia).jpg (https://commons.wikimedia.org/wiki/File:Sinustachycardia_(CardioNetworks_ECGpedia).jpg)

C IV dobutamine

If hypotension is severe and does not respond to plasma expanders, inotropic support with agents such as dobutamine may be necessary.

B Thrombolysis with streptokinase

Thrombolysis should be considered if the patient is not improving after 15–30 min with the above measures. The British Thoracic Society guidelines do not deliver any strong guidance with respect to thrombolysis, except that it should be considered where there is circulatory collapse. Pulmonary embolectomy should only be undertaken when thrombolysis is contraindicated. The diagnosis of pulmonary embolism is made by computed tomography (CT) or ventilation–perfusion (V/Q) scan.

D IV doxapram

The use of doxapram is not indicated in this scenario. Doxapram is a respiratory stimulant, which can lead to an increase in tidal volume and respiratory rate in patients with respiratory failure. It may be useful for treating respiratory depression in patients who have taken excessive doses of opiates and failed to respond adequately to treatment with naloxone.

E IV dopamine

Dopamine is not indicated in the management of this patient. There is, however, some evidence that it may be beneficial in patients with pulmonary embolism, a low cardiac index and a normal blood pressure.

71457

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

2014 ESC Guidelines on the diagnosis and management of acute pulmonary embolism: The Task Force for the Diagnosis ...
academic.oup.com/eurheartj/article/35/43/3033/503581
(https://academic.oup.com/eurheartj/article/35/43/3033/503581)

A 24-year-old asthmatic is on a salbutamol inhaler as needed (PRN) and a budesonide 200 µg inhaler twice daily (BD). He continues to wake up at night due to coughing and uses his salbutamol inhaler at least ten times during the week. His General Practitioner believes his asthma is not well controlled and needs his treatment stepping up.

Which of the following would be recommended according to the current (2017) National Institute for Health and Care Excellence (NICE) guidelines?

Your answer was incorrect

A Add oral montelukast to current therapy

B Add tiotropium inhaler to current therapy

C Increase budesonide to 800 µg BD

D Stop budesonide inhaler and start montelukast inhaler

E Stop budesonide inhaler and start oral montelukast

Explanation

A Add oral montelukast to current therapy

Montelukast is a leukotriene receptor antagonist. According to the 2017 National Institute for Health and Care Excellence (NICE) guidelines; if asthma is uncontrolled in adults (aged 17 and over) on a low dose of inhaled corticosteroids as maintenance therapy, offer a leukotriene receptor antagonist in addition to the inhaled corticosteroids and review the response to treatment in 4–8 weeks.

D Stop budesonide inhaler and start montelukast inhaler

Montelukast is given in the form of an oral tablet. In recent years, there have been studies to test the possibility of using montelukast inhalers. However, this product has not yet been brought to the market.

B Add tiotropium inhaler to current therapy

Tiotropium is a long-acting antimuscarinic licensed for maintenance treatment for chronic obstructive pulmonary disease and for treatment of asthma in patients not controlled on moderate-dose inhaled corticosteroids and long-acting beta-agonist (with or without montelukast). It is therefore not warranted at this stage of the therapeutic ladder.

C Increase budesonide to 800 µg BD

The standard dose of budesonide is 100–400 µg, and the high dose is 0.8–2 mg. Before using a high-dose steroid, a leukotriene antagonist should be considered and a long-acting beta-agonist should be tried.

E Stop budesonide inhaler and start oral montelukast

The NICE guidelines indicate that montelukast should be added to the current therapeutic regime, rather than given in place of steroid therapy. Asthma is a condition of hyperresponsive airways which is invariably steroid-responsive, so it is useful to remember that steroid therapy tends not to be stopped as part of the therapeutic escalation. Chronic obstructive pulmonary disease, on the other hand, is often not steroid-responsive and therefore, if steroid therapy is tried and found not to be helpful, it might be stopped in place of alternative therapy.

72952

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Asthma: diagnosis, monitoring and chronic asthma management
nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment
(https://www.nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment)

Asthma

Media

Expanded explanations

Flashcards

A 39-year-old woman from Zimbabwe presents to the Emergency Department with a 5-day history of increasing shortness of breath and dry cough. She has a 1-day history of sudden onset of sharp chest pain which is worse on coughing, movement and deep inspiration. She describes having had night sweats and a poor appetite for 2 months. Her weight has decreased by about 5 kg. She has a past medical history of genital herpes and depression. She is married and has two children who live in Zimbabwe. She moved to England 2 years ago and has not travelled since. She is a smoker of 20/day and drinks little alcohol.

On examination, she looked unwell. Observations: temperature 38.3 °C, BP 108/72 mmHg, pulse 120/min, regular, respiratory rate 30/min. Auscultation of her chest revealed inspiratory crackles bi-basally.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	102 g/l	135 - 175 g/l
White cell count (WCC)	6.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Lymphocytes	1.0 × 10^9/l	1.5 - 4.0 × 10^9/l
Platelets (PLT)	439 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	75 μmol/l	50 - 120 μmol/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
C-reactive protein (CRP)	73 mg/l	< 10 mg/l
Urea	5.0 mmol/l	2.5 - 6.5 mmol/l
Glucose	5.7 mmol/l	3.9 - 7.1 mmol/l

Blood gases on 80% oxygen:

pH	7.46
PCO_2	3.44 kPa
PO_2	10.77 kPa
Bicarbonate	17.6 mmol/l
Base excess	–3.5

Her chest X-ray is shown below.



What is the most likely underlying cause?

Your answer was correct

A Lymphoma

B HIV

C Bronchiectasis

D Sarcoid

E Common variable immune deficiency

Explanation

B HIV

For the purpose of the exam, people who have lived abroad (especially Africa and South America), businessmen who work abroad, men who have sex with men (MSM) and intravenous drug users, are more likely to have HIV.

This lady has HIV – the history of living in Zimbabwe and the low lymphocyte count are the clues. *Pneumocystis jirovecii* is the most common opportunistic infection to cause pneumonia in AIDS – especially when the CD4 count is <200/mm^3. It accounts for about 50% of cases of pneumonia in AIDS and 40% of all AIDS-defining illnesses.

Patients usually present with a fever, dry cough and breathlessness. They are usually hypoxic and desaturate on exercise.

In *P. jirovecii* pneumonia (PJP) the chest X-ray usually shows bilateral interstitial shadowing and cysts in the mid and lower zones. However, the CXR may be normal. Pneumothorax (because the cysts rupture) may be present in up to 10%.

Diagnosis is now usually made by PCR of sputum or BAL fluid. Other diagnostic methods include staining induced sputum or BAL with indirect immunofluorescence with monoclonal antibodies.

Treatment is with high-flow oxygen and high-dose co-trimoxazole. Prednisolone should be added in severe cases (PO_2 <9.5 kPa).

A Lymphoma

Given the patient is from sub-Saharan Africa and is presenting with a history suggestive of disseminated lung infection HIV is more a likely cause than lymphoma for the night sweats and weight loss.

C Bronchiectasis

The patient has a non-productive cough and gives an acute history of pulmonary infection, making bronchiectasis unlikely.

D Sarcoid

This patient presents with a history of pulmonary infection with fever, sweats and raised CRP. Sarcoidosis would be expected to present with chronic respiratory symptoms including a dry cough, and other features such as erythema nodosum.

E Common variable immune deficiency

This condition is present from birth and there would be a history of previous recurrent infections since childhood.

70100

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

You are asked to urgently review a 24-year-old woman who has been admitted to the Emergency Department. She has poorly controlled asthma for which she takes combination salmeterol/fluticasone two puffs bd and montelukast 10 mg daily. History of note includes two previous admissions to the Intensive Care Unit. On examination she looks tired, her BP is 132/72 mmHg and pulse is 100/min and regular. There is poor air entry and bilateral severe wheeze on auscultation of the chest. Peak flow on admission was 150 l/min, and this has only improved to 210 after three salbutamol nebulisers; her predicted is 580. Her PaO_2 is 7.8 kPa on arterial blood gas measurement.

Her X-ray is shown below:



Which one of the following features would indicate life-threatening asthma?

Your answer was incorrect

A PaO_2 of 7.8 kPa on air

B $PaCO_2$ of 4.0 kPa on air

C Distress and agitation

D Changes suggesting infection on CXR

E pH of 7.46

Explanation

A PaO_2 of 7.8 kPa on air

The fact that she is becoming severely hypoxic is a clear indication that this is life-threatening asthma. It is likely the underlying cause is pneumonia, as evidenced by the right middle lobe changes on the chest x-ray.

Criteria for the diagnosis of life-threatening asthma are clearly defined:

- PEF <33% best or predicted
- SpO_2 <92%
- PaO_2 <8 kPa
- normal $PaCO_2$ (4.6–6.0 kPa)
- silent chest
- cyanosis
- poor respiratory effort
- arrhythmia
- exhaustion, altered conscious level.

C Distress and agitation

Patients with asthma attacks are generally distressed and agitated due to the severity of the condition. If they start to tire then this can become very worrying and signals impending respiratory arrest.

B $PaCO_2$ of 4.0 kPa on air

A Pa CO_2 of 4.0 implies that the patient has low CO_2. In asthma exacerbation this occurs as the patient has a high respiratory rate and is exhaling CO_2 rapidly. In patients where the CO_2 is normalising, this can be a worrying sign as it implies tiring.

D Changes suggesting infection on CXR

Exacerbations of asthma may be due to underlying chest infection. Whilst chest x-ray here demonstrates right lobar pneumonia and IV antibiotics should be given in addition to the asthma treatment, the changes do not indicate life threatening asthma.

E pH of 7.46

The patient is likely to have exhaled large amounts of CO_2 through a high respiratory rate, and this is not alarming.

32373

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British guideline on the management of asthma 2018
sign.ac.uk/assets/asthma-consultation-draft.pdf
(https://www.sign.ac.uk/assets/asthma-consultation-draft.pdf)

A 19-year-old student is brought to the Emergency Department by his housemates. He has suffered a cough for the past week and using increasing amounts of his blue inhaler. Ordinarily he is maintained on fluticasone/salmeterol combination inhaler, 250 μg/50 mg BD, and montelukast, 10 mg daily. His housemates don't think he has registered with the student health service and are worried that he may not have filled his prescription.

Clinical features on admission to Emergency are respiratory rate 30/min, PEFR 240 l/min and predicted 590 l/min, and O_2 saturation 90%.

You commence oxygen therapy, give a bolus of IV hydrocortisone and give him a nebuliser containing salbutamol 5 mg and ipratropium 500 μg. The nurses repeat the salbutamol nebuliser three times.

Clinical features later in the evening on oxygen reveal the following:

Investigation	Result	Normal value
Respiratory rate	15 breaths per minute	
PEFR	230 l/min	
pH	7.35	7.35–7.45
pO_2	7.7 kPa	10.5–13.5 kPa
pCO^2	5.8 kPa	4.6–6.0 kPa

What is the most appropriate next step for this gentleman?

Your answer was incorrect

A IV salbutamol therapy

B IV aminophylline therapy

C IV magnesium therapy

D Review by the ITU outreach team

E Continuous nebulised salbutamol

Explanation

D Review by the ITU outreach team

The most appropriate next step in this case would be review by an ITU physician. Despite therapy his peak flow has deteriorated, he is making less respiratory effort and there is evidence of hypoxia with hypercapnia on arterial blood gas testing. He is at risk of respiratory arrest and should be assessed for ventilatory support.

B IV aminophylline therapy

IV aminophylline is no longer considered a first-line therapy for acute asthma because of the limited additional beta agonist efficacy versus continuous nebulisers, against a risk of significant arrhythmia.

A IV salbutamol therapy

IV salbutamol gives only limited advantage versus continuous nebuliser therapy, and increases the risk of atrial arrhythmia.

C IV magnesium therapy

IV magnesium is recommended in the management of acute severe asthma, although it may well not reverse the current deterioration in this patient’s symptoms. Hence planning for ITU admission is an essential next step.

E Continuous nebulised salbutamol

Although continuous nebulised salbutamol is appropriate here, review by the ITU outreach team is crucial with respect to outcomes.

18552

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Asthma

Media

Expanded explanations

Flashcards

A 54-year-old man presented with increasing shortness of breath and orthopnoea. He had noticed these symptoms over the last few years, but only went to his general practitioner (GP) after he had had a severe episode of breathlessness when he went wading in the sea on holiday. On further enquiry, he had also become aware of morning headaches and an increasing tendency to fall asleep during the day.

On examination, he looked well. Auscultation of his chest revealed reduced breath sounds and reduced percussion note bi-basally.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	177 g/l	135 - 175 g/l
White cell count (WCC)	12.3 × 10^9/l	4.0 - 11.0 × 10^9/l
Packed cell volume (PCV)	0.51	0.4 - 0.53 (male)
Platelets (PLT)	289 × 10^9/l	150 - 400 × 10^9/l

Investigation	Erect (% predicted)	Supine (% predicted)	Normal value
Residual volume	105%		1200ml
Total lung capacity	80%		80% to 120%
Vital capacity	75%	50%	FEV1 80% to 120% FVC 80% to 120%
TLCO	70%		> 60% to < 120%
KCO	95%		

Which of the following is the most helpful investigation in achieving the correct diagnosis?

Your answer was correct

A Chest X-ray

B Arterial blood gases

C Ultrasound screening of the diaphragm

D Echocardiogram

E Gas transfer and transfer coefficient

Explanation

C Ultrasound screening of the diaphragm

In bilateral diaphragmatic weakness, patients commonly present with breathlessness on exertion and when lying flat. It can be associated with sleep apnoea, resulting in daytime somnolence and headaches. The symptoms are due to the paroxysmal movement of the diaphragm during inspiration. In the supine position, expansion of the ribs results in movement of the abdominal contents into the chest, aided by gravity. Symptoms can be worse when standing up to the waist in water, as this counteracts the effects of gravity and prevents outward movement of the abdomen during inspiration and therefore produces a similar situation to being in the supine position.

A chest X-ray may show relatively small lung fields and basal linear shadowing due to subsegmental collapse. A SNIF test (diaphragmatic screening on ultrasound) will show paradoxical movement of the diaphragm, particularly on sniffing. Blood gases may show a type II respiratory failure, particularly at night. Lung function tests classically show a low vital capacity, which falls further in the supine position. All lung volumes except the residual volume are reduced. The gas transfer tends to be mildly impaired with a normal KCO. Treatment is with non-invasive ventilation.

A Chest X-ray

This may show relatively small lung fields but would not confirm diaphragmatic weakness.

B Arterial blood gases

These may show respiratory failure (either type 1 or chronic type 2) but cannot be used to diagnose diaphragmatic weakness.

D Echocardiogram

This would most likely be normal in this case, as the history is more in keeping with diaphragmatic weakness than cardiac disease.

E Gas transfer and transfer coefficient

This would likely be normal, or only mildly impaired, in this case, and would not aid in confirming the diagnosis.

70149

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 32-year-old man presents to the Respiratory Clinic. He describes being a chronic sufferer of allergic rhinitis, eczema and asthma for decades. These problems have worsened, and he has developed a chronic nocturnal cough despite regular steroid inhaler use. He has also started developing weight loss and malaise. He also complains that he has needed to see his GP several times for sinusitis and inner ear problems. He says that he takes a regular inhaler twice per day and a blue inhaler when he needs it, and uses emollients daily for his skin. He also says that he cannot take aspirin as this worsens his asthma.

Which investigation would be most likely to be diagnostic?

Your answer was incorrect

A Neutrophil count

B HRCT

C FEV_1 and FVC measurements

D IgG

E ANCA

Explanation

E ANCA

The presence of antineutrophil cytoplasmic antibody (ANCA) helps to rule in favour of eosinophilic granulomatosis with polyangiitis (found in 70% of cases), which is similar to asthma and an important differential diagnosis since the treatment is different. Eosinophilic granulomatosis with polyangitis is a vasculitis affecting small- to medium-vessel vasculitis with eosinophilia. It commonly affects many organs, including the lungs, cardiovascular system and renal system, and is heavily associated with atopy, otitis media and asthma, as in this patient. ANCA testing is the most appropriate way to differentiate from severe allergy and other causes of vasculitis.

C FEV_1 and FVC measurements

Although the forced expiratory volume in 1 s (FEV_1) and forced vital capacity (FVC) are non-specific in terms of diagnosis, they are useful in disease monitoring. In asthma attacks, $FEV_1$1 is lower when compared to the patient's established FEV_1. A FEV_1 at 35–50% of expected would be considered a severe asthma attack, whilst FEV_1 <33% of expected would be life-threatening.

A Neutrophil count

An eosinophilia count, rather than neutrophil count and IgE, can suggest allergic bronchopulmonary aspergillosis (ABPA). To diagnose ABPA, the sputum should be examined for *Aspergillus*, an aspergillus skin test can be done or an Aspergillus IgE RAST test can be done.

B HRCT

Computed tomography (CT) in expiration phase shows air trapping and can be seen especially in severe asthmatics. High-resolution computed tomography (HRCT) is only useful when looking for bronchiectasis in patients with ABPA.

D IgG

IgE, rather than IgG, levels of *Aspergillus* can be useful in differentiating ABPA from simple asthma.

2325

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Eosinophilic Granulomatosis with Polyangiitis

Expanded
explanations

A 24-year-old woman presents to the clinic with a nine-month history of nocturnal cough and shortness of breath, particularly first thing in the morning. She works as a PA in a solicitor's office. She notices wheezing on a number of days during the week. She is a non-smoker. Her weight is stable but she reports symptoms of reflux. A salbutamol inhaler used PRN has had limited impact on her symptoms. On examination her BP is 138/72 mmHg, with pulse 70/min and regular. Her chest is clear. Her BMI is 28. Chest X-ray is unremarkable and her peak flow is 400 (predicted is 480).

Which of the following is the most appropriate intervention for her asthma?

Your answer was incorrect

A Omeprazole

B Propping up the head of the bed

C Regular Salbutamol

D Regular Beclomethasone

E Regular salmeterol and fluticasone

Explanation

D Regular Beclomethasone

This patient has symptoms of asthma, particularly with nocturnal cough, which are not responsive to PRN salbutamol. As such, the next step is regular inhaled beclomethasone.

C Regular Salbutamol

Regular salbutamol is relatively short acting and is therefore a poor treatment option for nocturnal symptoms of asthma.

A Omeprazole

Although it was previously thought that proton pump inhibitors are effective at treating asthma in this situation, a meta-analysis (see weblink) has challenged that view.

B Propping up the head of the bed

Although this is an effective option for managing symptoms of reflux, it’s unlikely to impact on her worsening nocturnal cough.

E Regular salmeterol and fluticasone

Regular salmeterol and fluticasone is considered a potential escalation step, in the event that symptoms are not controlled on regular inhaled beclomethasone alone.

36468

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

The efficacy of proton pump inhibitors for the treatment of asthma in adults: a meta-analysis
ncbi.nlm.nih.gov/pubmed/21482834
(http://www.ncbi.nlm.nih.gov/pubmed/21482834)

Asthma

Media

Expanded explanations

Flashcards

A 26-year-old man presents with episodic wheeze and cough. He feels that his symptoms are worse in the morning. The symptoms have been present intermittently over a number of years. While the cough is usually non-productive, in the last two weeks, he has been coughing green sputum.

He has a background of anxiety and previous mumps as a child and works in a shipyard. He smokes approximately ten cigarettes a day.

Chest examination was unremarkable. Chest X-ray and spirometry was undertaken, the results of which are shown below.





Which of the following is the most likely diagnosis?

Your answer was incorrect

A Asbestosis

B Asthma

C Bronchiectasis

D Chronic obstructive pulmonary disease (COPD)

E Idiopathic pulmonary fibrosis

Explanation

B Asthma

This is a history of shortness of breath and cough with diurnal variation and a normal chest x-ray, which raises suspicion of asthma. While classically asthma may have an obstructive picture on spirometry, as shown here, it may also be normal. In this pattern, the maximal expiratory flow rate is reduced, with the concave shape of the curve indicating an obstructive pattern. In investigation of asthma, spirometry is undertaken before and following bronchodilator administration, to assess for reversibility.

Image source: https://commons.wikimedia.org/w/index.php?curid=7036322

D Chronic obstructive pulmonary disease (COPD)

Spirometry is essential in diagnosing COPD, with the picture being obstructive. However, spirometry is a poor predictor of the extent of disease and disability in COPD. Assessing the clinical context, the patient is much too young for COPD despite being a smoker.

A Asbestosis

This is a history of a young person with classic symptoms of asthma. While working on a shipyard may make him more at risk of asbestosis, this usually occurs in older patients. This is due to a latency period, which may be five years at a minimum post-exposure, though this period may extend to 40 or more years. While asbestosis generally has little impact on spirometry, asthma may also present with normal spirometry. The chest x-ray in asbestosis can be normal or may show fine reticular changes with evidence of asbestos exposure in the form of pleural plaques.

C Bronchiectasis

While previous mumps may make an individual at risk of bronchiectasis, the history of productive sputum is not particularly long enough to be convincing of bronchiectasis. This is more likely to be an infection on a background of asthma. Spirometry in bronchiectasis is usually obstructive and the chest x-ray can show an increase in bronchovascular markings, tram-track opacities and cystic changes with air-fluid.

E Idiopathic pulmonary fibrosis

Idiopathic pulmonary fibrosis usually presents with a restrictive pattern on spirometry, rather than the obstructive picture demonstrated here, and fibrosis on the chest x-ray.

71787

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 24-year-old patient comes to the emergency department with acute breathlessness. She is a known asthmatic and has a previous history of requiring admission for her asthma. She usually takes beclomethasone 500 µg bd, but over the past day has needed repeated doses of her salbutamol inhaler. Over the past hour she has becoming increasingly breathless, despite taking regular salbutamol. On examination the patient looks tired and has increased work of breathing. Her respiratory rate is 40 breaths per minute, heart rate is 110 bpm and blood pressure is 105/75 mmHg. Her temperature is 37.2°C. Examination of the chest reveals generalised wheeze and poor air entry. Oxygen saturations are 86% on room air.

Arterial blood gases reveal the following:

Investigation	Result	Normal Value
pH	7.35	7.35–7.45
pO_2	7.1 kPa	10.5–13.5 kPa
pCO_2	5.1 kPa	4.6–6.0 kPa

Which of the following would be the most appropriate next management step?

Your answer was correct

A Back-to-back salbutamol nebulisers and admission to the medical ward

B Back-to-back salbutamol nebulisers, IV corticosteroids and admission to the medical high-dependency unit (HDU)

C IV hydrocortisone

D Nebuliser therapy and later discharge from the Emergency Department

E Addition of long-acting β-agonists to her therapy

Explanation

B Back-to-back salbutamol nebulisers, IV corticosteroids and admission to the medical high-dependency unit (HDU)

Respiratory muscle fatigue and acidosis are known to worsen established hypoxia and lead to a vicious circle of further muscle weakness. As such, early admission to the medical HDU should be considered. It is likely this patient will rapidly need ventilatory support ± intubation. Long-term she is likely to require the addition of long-acting β-agonists to her therapy. A combination inhaler such as salmeterol/fluticasone (sere tide) may be the most effective way to deliver the two drugs together.

A Back-to-back salbutamol nebulisers and admission to the medical ward

Her O_2 is low and CO_2 raised, indicating a severe asthma attack. Salbutamol nebulisation represents a significant component of her management, but the medical ward represents an inadequate place of care; she should be managed in a critical care area.

C IV hydrocortisone

This represents an appropriate therapy but should not represent the sole management of her condition.

D Nebuliser therapy and later discharge from the Emergency Department

This patient is having a severe asthma attack – nebuliser monotherapy and discharge without further investigation/assessment would be inappropriate.

E Addition of long-acting β-agonists to her therapy

This will likely be required in due course, but does not represent the most appropriate next step in her management.

6606

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Asthma

Media

Expanded explanations

Flashcards

A 54-year-old woman has a left adenocarcinoma of the bronchus confirmed on transbronchial biopsy. She tells you that she has lost weight over the past few months and has a nagging cough, particularly in the mornings, but actually has still been able to hold down her job, riding her bike the short distance to work and caring for her teenage children.

Investigations reveal the following:

Investigation	Result	Normal values
Haemoglobin (Hb)	135 g/l	115–155 g/l
White cell count (WCC)	5.6 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	190 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	120 μmol/l	50–120 μmol/l

Which of the following is likely to be most informative with respect to determining the indication for surgery?

Your answer was incorrect

A Pulmonary function testing

B CT thorax

C PET/CT combined scan

D Serum calcium

E CT Head

Explanation

C PET/CT combined scan

PET/CT combined scanning offers the best imaging modality to determine lymph node involvement in bronchial carcinoma. In this case, with a good functional status at the point of diagnosis, it’s lymph node involvement which is likely to be the biggest determinant of the success or failure of surgical intervention.

D Serum calcium

A number of reasons for hypercalcaemia quite unconnected with bronchial carcinoma may be apparent – hence raised calcium *per se* would not represent a contraindication to surgery. PTHrp is also much less likely to be elevated in patients with an underlying adenocarcinoma versus other bronchial carcinomas.

A Pulmonary function testing

Of course, FEV_1 and FVC do have an impact on suitability for surgery, but given that we are told she cycles to work and aren't given information on her smoking status, it is likely that her lung function is adequate to undergo pneumonectomy.

B CT thorax

CT thorax is informative with respect to tumour staging, but won't deliver adequate information with respect to lymph node involvement.

E CT Head

We are given no indication of cerebral metastases, meaning that CT head is of limited value here.

21214

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 25-year-old man presents with left-sided pleuritic chest pain that come on suddenly 3 days previously. Initially he thought it was secondary to his workout in the gym. However, since it did not improve and he noticed that he was slightly more breathless than usual, he went to the Emergency Department. He has no significant past medical history. He is a smoker of 20/day. On examination he looked well. He was a tall, thin man and was not short of breath at rest. Examination was unremarkable except for a clicking sound which was synchronous with the heart sounds.

Chest X-ray is shown below.



What is the diagnosis?

Your answer was correct

A Mitral valve prolapse

B Pericarditis

C Primary pneumothorax

D Pleurisy

E Secondary pneumothorax

Explanation

C Primary pneumothorax

A click synchronous with the heart sounds is a recognised sign of a small left apical pneumothorax. He is a tall, thin man and a smoker, which are both risk factors for developing a pneumothorax. There is a strong association between pneumothoraces and smoking. The most recent British Thoracic Society (BTS) guidelines suggest that if the postero-anterior (PA) chest X-ray is normal, as is the case here, and a small pneumothorax is suspected, a lateral decubitus chest X-ray provides added information in up to 14% of cases (see weblink below). Expiratory films add little and are not recommended.

The treatment of a primary pneumothorax in a non-smoker, <50 years old, with no evidence of underlying lung disease, with a rim of air of <2 cm and no breathlessness is to discharge and follow up as an outpatient. Although this patient is a smoker there is nothing to suggest underlying lung disease, and it is likely that his apical pneumothorax is too small to safely aspirate. Patients should be given analgesia if required and clear instructions to return if their symptoms get any worse.

A Mitral valve prolapse

This is associated with a mid- to late systolic click. It occurs when there is an abnormally thickened mitral valve leaflet that displaces into the left atrium during systole.

B Pericarditis

The patient has pleuritic, not pericarditic, chest pain. Pericarditis typically causes symptoms that include sudden onset of sharp chest pain. The pain may be felt in the shoulders, neck or back. It is typically relieved by sitting up and worse when lying down or deep inspiration. Other symptoms include fever, weakness, palpitations or dyspnoea.

D Pleurisy

Onset was sudden and not associated with infective symptoms. Pleurisy results in a sharp chest pain on breathing. Other symptoms include breathlessness, cough, fever and weight loss.

E Secondary pneumothorax

There is no history of underlying lung disease. Secondary pneumothorax is defined as pneumothorax that occurs in the presence of underlying lung disease. This is in contrast to primary pneumothorax, where there is no lung disease.

70118

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

BTS. Management of pleural infection in adults
thorax.bmj.com/content/65/Suppl_2/ii41
(https://thorax.bmj.com/content/65/Suppl_2/ii41)

Pneumothorax

Media

Expanded explanations

Flashcards

A 25-year-old teacher presents with a 3-year history of recurrent chest infections that never really completely resolved after completing a course of antibiotics. She has a cough most days and produces large volumes of green sputum on a daily basis. She has had mild haemoptysis in the past. She has noticed weight loss over the past year with increased lethargy and shortness of breath on exertion. She has no previous past medical history and is not on any regular medications. On examination she has clubbing and coarse inspiratory crackles in both her bases, which are worse on the right side. She is 135 cm in height, apparently significantly shorter than her brothers and sisters.

Which of the following tests is most likely to contribute to determining her underlying diagnosis?

Your answer was incorrect

A cANCA (anti-neutrophil cytoplasmic antigen)

B Delta F508 mutation

C Hepatitis screen

D Alpha-1-antitrypsin testing

E E ESR (erythrocyte sedimentation rate)

Explanation

B Delta F508 mutation

The patient clinically has bronchiectasis with coarse inspiratory crackles and clubbing and a history of copious purulent sputum and cough. A number of causes exist with respect to bronchiectasis in this age group, but heterozygosity for the cystic fibrosis CFTR mutation is increasingly recognised as being associated with the disease.

D Alpha-1-antitrypsin testing

Alpha-1 antitrypsin deficiency is associated with the development of emphysema, rather than bronchiectasis.

A cANCA (anti-neutrophil cytoplasmic antigen)

ANCA-positive vasculitis may be associated with pulmonary disease, although pulmonary haemorrhage is the usual presentation.

C Hepatitis screen

Viral hepatitis is not usually associated with bronchiectasis, although evidence of cirrhosis may co-exist with emphysema in patients with alpha-1-antitrypsin deficiency.

E E ESR (erythrocyte sedimentation rate)

ESR is a non-specific marker of pulmonary inflammation or infection, and is therefore not useful in identifying the underlying cause of her symptoms.

32431

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Cystic Fibrosis

Media

Expanded explanations

Flashcards

A 67-year-old man is admitted to the Acute Medical Unit with a 5-day history of fever and a cough productive of purulent sputum. He was started on intravenous (IV) antibiotics for community-acquired pneumonia (CAP). His past medical history includes recently diagnosed prostatic cancer, hypertension and stable angina, for which he takes aspirin, bisoprolol, amlodipine, atorvastatin, lansoprazole, tamsulosin and glyceryl trinitrate as required.

Three days into his admission, he complains of left-sided chest pain and shortness of breath. On examination, his temperature is 38.2 °C. He has a heart rate of 110 bpm and a blood pressure of 189/105 mmHg. His respiratory rate is 29 breaths/minute, with saturations of 89% on room air. On auscultation, there are coarse crackles at the left base and heart sounds are normal. He has a prolonged capillary refill time of three seconds and his calves are soft, with mild pitting oedema to the ankles.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin (Hb)	132 g/l	115–155 g/l
White cell count (WCC)	12.3 × 10^9/l	4.0–11.0 × 10^9/l
Neutrophils	11.1 × 10^9/l	2.5–7.5 × 10^9/l
Lymphocytes	2.9 × 10^9/l	0.8–5.0 × 10^9/l
Platelets (PLT)	394 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	136 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Urea	7.3 mmol/l	1.8–7.1 mmol/l
Creatinine (Cr)	125 µmol/l	50–120 µmol/l
Estimated glomerular filtration rate (eGFR)	57 ml/min/1.73 m^2	60+ ml/min/1.73 m^2
International normalised ratio (INR)	1.3	0.9–1.2
Fibrinogen	4.1 g/l	2–4 g/l
D-dimer	800 ng/ml	< 400 ng/ml
Troponin	32 ng/ml	0–14 ng/ml
C-reactive protein (CRP)	115 mg/l	< 10 mg/l

Electrocardiogram (ECG) is shown below



What is the most likely diagnosis?

Your answer was incorrect

A Acute coronary syndrome

B Aortic dissection

C Pericarditis

D Pleurisy

E Pulmonary embolism

Explanation

E Pulmonary embolism

This patient has a pulmonary embolism (PE). The ECG shows a sinus tachycardia, one of the commonest findings in PE. An ongoing significant infection, combined with a reduction in mobility during hospital admission, recent malignancy increases the likelihood of developing a venous thromboembolism (VTE). This accounts for the patient's acute-onset shortness of breath and chest pain. Although a raised D-dimer can be seen in a number of conditions, including infection, malignancy and VTE, a positive value, in conjunction with chest pain and shortness of breath, should point towards a pulmonary embolism. A pulmonary embolism can increase pulmonary arterial pressures and cause right heart strain. This may present as a raised troponin and ischaemic changes in the right precordial leads, such as T-wave inversion in leads V1-V4.

B Aortic dissection

Aortic dissection typically causes chest pain which radiates through to the back. It is more common in patients with hypertension. A discrepancy in blood pressure in both arms can be seen, along with significantly raised D-dimer levels, mediastinal widening on a chest X-ray and ST-segment elevation in the inferior leads in the case of right coronary artery dissection. Given this patient's symptoms and clinical findings, aortic dissection seems less likely.

A Acute coronary syndrome

Acute coronary syndrome (ACS) is an important differential in this setting, particularly with a background of hypertension and ischaemic heart disease. Raised cardiac enzymes, such as troponin, are seen in ACS, along with ECG changes, in keeping with ischaemia. T-wave inversion in leads V1–V4 and a troponin level of 32 ng/ml could represent an ACS. However, the addition of a raised D-dimer suggests an alternative diagnosis.

C Pericarditis

Pericarditis is a cause of pleuritic chest pain. However, the ECG and laboratory findings suggest an alternative diagnosis. Furthermore, given the clinical history and hospital admission, this patient is more likely to have a VTE.

D Pleurisy

Pleurisy is the inflammation of the pleura which commonly arises secondary to an underlying infection, effusion or pulmonary embolism. However, in the context of chest pain, shortness of breath and the laboratory and ECG findings, it seems less likely that this patient's symptoms are from pleurisy alone.

72583

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pulmonary Embolism

Media

Expanded explanations

Flashcards

A 60-year-old South Asian patient presents with shortness of breath and weight loss with night sweats. On examination, his respiratory rate is 24 breaths/min, he has lymphadenopathy and his left lower zone is stony dull to percussion. His chest X-ray demonstrates evidence of a moderate pleural effusion. You suspect tuberculosis as he confirms that his wife was recently diagnosed with the disease.

Investigations:

Investigation	Result	Normal value
Haemoglobin	125 g/l	135–175 g/l
White cell count (WCC)	11.0 × 10^9/l	4–11 × 10^9/l
Platelets	342 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Potassium (K^+)	3.8 mmol/l	3.5–5.0 mmol/l
Creatinine	54 μmol/l	50–120 μmol/l
Mean corpuscular volume (MCV)	89.3 fl	76–98 fl
Urea	4.0 mmol/l	2.5–6.5 mmol/l
C-Reactive protein (CRP)	85 mg/l	0–10 mg/l
Erythrocyte sedimentation rate (ESR)	43 mm/h	0–10 mm in the 1st hour

Which of the following investigations is most useful in confirming the cause of the pleural effusion?

Your answer was incorrect

A CT pleural biopsy and TB-PCR

B High-resolution CT

C Pleural fluid pH

D Quantiferon gamma

E Pleural fluid LDH

Explanation

A CT pleural biopsy and TB-PCR

Pleural biopsy has a high yield in confirming active tuberculosis. The key to diagnosing tuberculosis is obtaining a tissue sample, particularly so that PCR for tuberculous DNA can be carried out to provide rapid diagnosis, to allow early differentiation from non-tuberculosis mycobacterial infection, and to detect multidrug-resistant disease.

D Quantiferon gamma

Quantiferon is an interferon-gamma release assay, specifically acting as a surrogate marker of *Mycobacterium tuberculosis*, and thus it cannot differentiate between latent and active infection. Therefore, a positive quantiferon gamma is of no diagnostic utility in this case, as it could simply indicate previous infection or infection with non-tuberculous mycobacteria.

B High-resolution CT

Imaging such as CT will only determine the location and extent of the effusion, but it won’t give any diagnostic information regarding whether it is caused by tuberculosis. CT will also help determine whether there is any mass underlying the effusion.

C Pleural fluid pH

A pleural fluid pH <7.2 will confirm empyema, but not the infecting organism causing the empyema.

E Pleural fluid LDH

LDH is non-specifically raised in a number of conditions other than tuberculosis, such as a rheumatoid arthritis-related effusion or bacterial infection.

Pleural biopsy has a good detection rate for pleural TB. CT will not help determine the cause of the effusion, merely delineate its size and detect whether there is evidence of an underlying mass. Fluid pH is less than 7.2 in empyema and hence it is useful in pointing towards an infective cause but not necessarily TB. A Heaf test, if grade 4, will tell you that the patient has active TB, but again won’t necessarily indicate that the effusion itself is related to TB infection. LDH is non-specifically raised in many causes of effusion, including infection, rheumatoid and TB.

Pleural biopsy has a good detection rate for pleural TB. CT will not help determine the cause of the effusion, merely delineate its size and detect whether there is evidence of an underlying mass. Fluid pH is less than 7.2 in empyema and hence it is useful in pointing towards an infective cause but not necessarily TB. A Heaf test, if grade 4, will tell you that the patient has active TB, but again won’t necessarily indicate that the effusion itself is related to TB infection. LDH is non-specifically raised in many causes of effusion, including infection, rheumatoid and TB.

32399

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pleural Effusion

Media

Expanded explanations

Flashcards

A 39-year-old lady from Zimbabwe presents to the Emergency Department with a 5-day history of increasing shortness of breath and dry cough. She has a 1-day history of sudden onset of sharp chest pain which is worse on coughing, movement and deep inspiration. She describes having had night sweats and a poor appetite for 2 months. Her weight has decreased by about 5 kg. She has a past medical history of genital herpes and depression. She is married and has two children who live in Zimbabwe. She moved to England 2 years ago and has not travelled since. She is a smoker of 20/day and drinks little alcohol.

On examination she looked unwell. Observations: temperature 38.3°C, BP 108/72 mmHg, pulse 120/min, regular, respiratory rate 30/min. Auscultation of her chest revealed inspiratory crackles bi-basally.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	102 g/l	135 - 175 g/l
White cell count (WCC)	9.4× 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	8.6× 10^9/l	1.5 - 7.0× 10^9/l
Platelets (PLT)	439× 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
C-reactive protein (CRP)	73 mg/l	< 10 mg/l
Sodium (Na^+)	141 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Urea	5.0 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	75 µmol/l	50 - 120 µmol/l
Glucose	5.7 mmol/l	3.9 - 7.1 mmol/l

Blood gases on 80% oxygen	Result
pH	7.46
PCO_2	3.44 kPa
PO_2	10.77 kPa
Bicarbonate	17.6 mmol/l
Base excess	-3.5

Her chest X-ray is shown:



Which investigation would be most helpful in establishing a diagnosis?

Your answer was incorrect

A Bronchoscopy and transbronchial biopsy

B CT pulmonary angiogram

C Lung function tests and transfer factor

D Indirect immunofluorescence of bronchoalveolar lavage fluid

E Blood cultures

Explanation

D Indirect immunofluorescence of bronchoalveolar lavage fluid

Diagnosis is now usually made by PCR of sputum or BAL fluid. Other diagnostic methods include staining induced sputum or BAL with indirect immunofluorescence with monoclonal antibodies.

For the purpose of the exam, people who have lived abroad (especially Africa and South America), businessmen who work abroad, men who have sex with men (MSM) and intravenous drug users, are more likely to have HIV.

This lady has HIV – the history of living in Zimbabwe and the low lymphocyte count are the clues. Incidentally, although the lymphocyte count is not given, it can be inferred from the available data, ie since the white cell total is 9.4 and the neutrophils are 8.6, that leaves 0.8, which means lymphocytes are definitely going to be reduced. *Pneumocystis jirovecii* is the most common opportunistic infection to cause pneumonia in AIDS – especially when the CD4 count is <200/mm^3. It accounts for about 50% of cases of pneumonia in AIDS and 40% of all AIDS-defining illnesses.

Patients usually present with a fever, dry cough and breathlessness. They are usually hypoxic and desaturate on exercise.

In *P. jirovecii* pneumonia (PJP) the chest X-ray usually shows bilateral interstitial shadowing and cysts in the mid and lower zones. However, the CXR may be normal. Pneumothorax (because the cysts rupture) may be present in up to 10%.

Treatment is with high-flow oxygen and high-dose co-trimoxazole. Prednisolone should be added in severe cases (PO_2 <9.5 kPa).

B CT pulmonary angiogram

CTPA is an appropriate investigation for PE. The abnormal chest X-ray with evidence of pulmonary infiltration counts against a PE as the underlying diagnosis here.

A Bronchoscopy and transbronchial biopsy

Transbronchial biopsy would cause undue risk of pneumothorax in a patient already at risk of pneumothorax due to *Pneumocystis* pneumonia. Also, it is likely that this patient will end up invasively ventilated, further adding to risk of pneumothorax.

C Lung function tests and transfer factor

These tests would not be the most likely to give a definitive diagnosis: they would simply show the degree of lung function impairment. In this case, due to pulmonary infiltration, it’s likely that transfer factor will be reduced.

E Blood cultures

Blood cultures are likely to be negative in *Pneumocystis* pneumonia, with local invasion and inflammatory infiltration the drivers for symptoms.

70099

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

An 18-year-old man presents to the Outpatients Department having coughed up a small amount of bright red blood. He is mildly short of breath and feels ‘wheezy’. His mother says he has always been a ‘chesty’ child. He was teased at school as he has always been the smallest in his class. His brother and sister were fit and well. He had a dog at home.

On examination, he looked well and was not breathless at rest. He had clubbing. He had nasal polyps and auscultation of his chest revealed coarse crackles throughout both lung fields. His abdomen was distended but soft.

Urinalysis:

Protein	+
Glucose	++
Nitrites	-
Leucocytes	-

Haematology:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	110 g/l	135 - 175 g/l
White cell count (WCC)	12.1 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	450 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	137 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	62 μmol/l	50 - 120 μmol/l
Neutrophils	91%	
Lymphocytes	8%	
Monocytes	0.4%	
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	35 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	40 mg/l	< 10 mg/l
Urea	6.4 mmol/l	2.5 - 6.5 mmol/l
Bilirubin	31 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	46 u/l	10 - 40 u/l

Alkaline phosphatase (ALP)	221 u/l	30 - 150 u/l
Albumin	38 g/l	35 - 55 g/l
Protein	74 g/l	
Magnesium (Mg^{2+})	0.64 mmol/l	0.6 - 1.1 mmol/l
Corrected calcium (Ca^{2+})	2.1 mmol/l	2.2 - 2.7 mmol/l
Phosphate (PO_4^{3-})	0.75 mmol/l	1.1 - 1.45 mmol/l

Spirometry:

FEV1	60% predicted
FVC	80% predicted
PEF	350 l/min
KCO	70% predicted

His chest X-ray is shown below:



What is the most likely underlying diagnosis?

Your answer was incorrect

A Cystic fibrosis

B Kartagener’s syndrome

C Asthma

D Allergic bronchopulmonary aspergillosis

E Goodpasture’s syndrome

Explanation

A Cystic fibrosis

This patient has cystic fibrosis. He has had recurrent chest infections secondary to bronchiectasis (confirmed by examination findings and the chest X-ray, but also note the long line which has probably been inserted via the left antecubital fossa), he is small for his age and has nasal polyps (which occur in one third of patients with CF) and clubbing. Pulmonary function tests show airflow obstruction. He has mildly deranged liver function tests – secondary to obstruction of biliary ductules in the liver. This can eventually lead to cirrhosis. He has glucose in his urine – pancreatic islet cells are destroyed as the pancreas becomes fibrotic, resulting in CF-related diabetes in 30% of patients by adulthood.

The most likely cause of this patient’s haemoptysis is secondary to infection. Larger-volume hemoptysis can occur in cystic fibrosis due to bronchial artery hypertrophy. Ninety per cent of patients with cystic fibrosis will have steatorrhoea secondary to pancreatic insufficiency if pancreatic enzyme replacement therapy is not taken with food.

E Goodpasture’s syndrome

Goodpasture syndrome, or anti-GBM disease, is a cause of haemoptysis, although it is associated with acute deterioration in renal function with glomerulonephritis and marked haemoptysis.

B Kartagener’s syndrome

Kartagener syndrome causes bronchiectasis and situs inversus. The chest radiograph does not show dextrocardia.

C Asthma

The chest X-ray is inconsistent with a diagnosis of simple asthma. With significant patchy interstitial shadowing and failure to thrive, cystic fibrosis is suspected.

D Allergic bronchopulmonary aspergillosis

ABPA is associated with proximal bronchiectasis, fleeting pulmonary infiltrates, lobar collapse and wheeze. It responds to treatment with systemic corticosteroids and itraconazole.

70085

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 38-year-old woman presents to the Respiratory Clinic for review of her asthma control. She is currently taking 200 µg twice daily (BD) of inhaled beclomethasone and wakes up coughing at night. Peak flows average at 360 litres/min (compared to 490 litres/min predicted). Her BP is 110/80 mmHg, and her heart rate is 75 bpm and regular. There is scattered wheeze on auscultation of the chest.

According to the National Institute for Health and Care Excellence (NICE) guidelines, which of the following is the most appropriate next step?

Your answer was incorrect

A Add low-dose oral prednisolone

B Add montelukast

C Add theophylline

D Increase inhaled steroid

E Add salmeterol

Explanation

B Add montelukast

Montelukast would be the most appropriate here as the NICE guidelines recommend a trial of montelukast in patients who have failed to gain control of their symptoms on low-dose inhaled corticosteroids.

D Increase inhaled steroid

Increasing inhaled steroids is not appropriate according to the NICE 2017 guidelines due to the potential increased rate of adverse effects versus the marginal benefit derived. A trial of montelukast is required in this situation before other options, such as increasing corticosteroids, are considered.

A Add low-dose oral prednisolone

Given the side effect profile of long-term steroids, including osteoporosis, blood glucose derangement, fat redistribution and potential adrenal suppression, low-dose oral prednisolone is used only in refractory asthma. Typically, low-dose oral prednisolone is considered after trials of increasing inhaled steroid, adding montelukast and adding theophylline, if appropriate, have been exhausted first.

C Add theophylline

Theophylline is a nonselective phosphodiesterase inhibitor which results in bronchodilation. It has a narrow therapeutic window and can result in multiple adverse effects such as headaches, insomnia, nausea, reflux and seizures. It is therefore reserved for use if inhaled therapies and montelukast have failed to control the patient's symptoms.

E Add salmeterol

Adding salmeterol is only an option in the event that symptoms remain uncontrolled with the combination of montelukast and inhaled corticosteroids.

71365

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society. 2019. BTS/SIGN British guideline on the management of asthma.
brit-thoracic.org.uk/quality-improvement/guidelines/asthma/
(https://www.brit-thoracic.org.uk/quality-improvement/guidelines/asthma/)

NICE. 2017. Asthma: diagnosis, monitoring and chronic asthma management.
nice.org.uk/guidance/ng80
(https://www.nice.org.uk/guidance/ng80)

Asthma

Media

Expanded explanations

Flashcards

A 31-year-old asthmatic woman was seen in the Respiratory Outpatient Clinic. She said she had been generally well although she had not been sleeping well at night due to coughing. She had also noticed that when playing tennis she was getting wheezy and requiring her salbutamol inhaler more frequently – at least 4 times a week. She was taking budesonide 200 micrograms two puffs twice a day. She had no significant past medical history. She smoked 10 cigarettes/day and had done so for 2 years.

Auscultation of her chest revealed a mild expiratory wheeze. Examination was otherwise unremarkable.

How would you treat this patient?

Your answer was incorrect

A Increase Budesonide

B Add Montelukast

C Give a course of oral Prednisolone

D Add Aminophylline

E Start a regular Salbutamol inhaler

Explanation

B Add Montelukast

This woman is on step 2 of the asthma treatment ladder – she is on a low/moderate dose inhaled steroids (800 mcg/day). A trial of an alternative agent should be added prior to increasing inhaled steroids. The first choice of additional therapy would be a leukotriene receptor antagonist such as montelukast, for a trial, as per the NICE 2017 asthma guidelines.

C Give a course of oral Prednisolone

This is not an acute asthma exacerbation requiring oral steroid therapy, this lady has poor asthma control and needs her maintenance treatment increased.

A Increase Budesonide

She is already on a moderate dose of inhaled corticosteroid. The next most appropriate step is to add in a long acting bronchodilator. If this were to have no effect then the dose of inhaled corticosteroid should be increased.

D Add Aminophylline

Aminophylline is reserved for patients who have not responded to high dose inhaled corticosteroids in combination with a long acting inhaled beta 2 agonist.

E Start a regular Salbutamol inhaler

Salbutamol is a short acting beta 2 agonist and would need to be taken multiple times per day to relieve symptoms. It is more appropriate to give a long acting beta 2 agonist. Short acting beta 2 agonists like Salbutamol are reserved for acute relief of symptoms rather than prevention.

70123

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Asthma: diagnosis, monitoring and chronic asthma management
nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment

(https://www.nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment)

Asthma

Media

Expanded explanations

Flashcards

A 24-year-old asthmatic on PRN salbutamol inhaler and budesonide 200 µg inhaler twice per day is not well controlled and needs his treatment stepping up.

Which of the following would be recommended according to current (2017) NICE guidelines?

Your answer was incorrect

A Add oral cromoglicate

B Increase budesonide to 800 µg BD

C Add oral theophylline (modified release)

D Add montelukast

E Add tiotropium

Explanation

D Add montelukast

Montelukast is a leukotriene receptor antagonist. The 2017 NICE guidelines say:

"If asthma is uncontrolled in adults (aged 17 and over) on a low dose of ICS as maintenance therapy, offer a leukotriene receptor antagonist (LTRA) in addition to the ICS and review the response to treatment in 4–8 weeks."

C Add oral theophylline (modified release)

Theophylline would be added if asthma is not controlled after trying high-dose inhaled steroids.

A Add oral cromoglicate

Cromoglicate is only effective for asthma prophylaxis by inhalation 3–4 times a day and is generally only considered after patients have failed to gain control on multiple agents, including inhaled corticosteroids, long-acting beta-agonist and monteleukast.

B Increase budesonide to 800 µg BD

The standard dose of budesonide is 100–400 µg, the high dose is 0.8–2mg. Before using a high-dose steroid, a long-acting beta-agonist should be tried.

E Add tiotropium

Tiotropium is a long-acting anti-muscarinic licensed for maintenance treatment for COPD and for the treatment of asthma in patients not controlled on inhaled corticosteroids and long-acting beta-agonist.

2324

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Asthma: diagnosis, monitoring and chronic asthma management
nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment
(https://www.nice.org.uk/guidance/ng80/chapter/Recommendations#principles-of-pharmacological-treatment)

Asthma

Media

Expanded explanations

Flashcards

A 40-year-old farmer presents with a 1-month history of wheeze and shortness of breath associated with working on the farm and resolving about 6 h after stopping work.

On examination, his blood pressure (BP) is 136/82 mmHg, pulse is 82/min and regular and oxygen saturation is 94% on air. He has coarse inspiratory crackles and a dry cough. You suspect he is suffering from extrinsic allergic alveolitis.

Which feature is most likely to be seen on his chest X-ray?

Your answer was incorrect

A Bilateral hilar lymphadenopathy

B Lower zone nodular pattern of fibrosis

C Upper zone nodular pattern of fibrosis

D Pleural thickening

E Honeycomb pattern in the upper zones

Explanation

C Upper zone nodular pattern of fibrosis

Extrinsic allergic alveolitis usually has a nodular pattern bilaterally, with a predilection for the upper lobes .

B Lower zone nodular pattern of fibrosis

Extrinsic allergic alveolitis affects the upper lobes. Lower zone fibrosis usually occurs in those with asbestosis, rheumatoid arthritis or connective tissue diseases, or as a side effect of certain drugs (busulphan, bleomycin, nitrofurantoin, methotrexate, amiodarone).

A Bilateral hilar lymphadenopathy

This is rarely seen in extrinsic allergic alveolitis. It is more commonly associated with sarcoidosis, malignancy (lymphoma and carcinoma) and infection (tuberculosis, mycoplasma and silicosis).

D Pleural thickening

This is a non-specific feature that can occur with both benign (such as occupational – asbestos exposure) and malignant pleural disease (mesothelioma, pleural metastases) rather than extrinsic allergic alveolitis.

E Honeycomb pattern in the upper zones

This is usually a manifestation of end-stage interstitial fibrosis. This gentleman has a short presentation history and saturating 94% on air, making honeycomb pattern on radiology unlikely. Honeycombing occurs due to the airspaces consisting of irregular, thickened fibrotic walls and inflammatory cells.

32447

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Hypersensitivity Pneumonitis

Expanded
explanations

A 37-year-old Afro-Caribbean woman presents to the Emergency Department with a painful raised rash affecting both shins. She has been followed up on a number of occasions over the past few months by her General Practitioner because of complaints of shortness of breath and a dry cough. This has not responded to a salbutamol inhaler or two courses of amoxicillin. There is no past medical history of note.

On examination, her blood pressure is 145/89 mmHg; pulse is 75 bpm and regular. Her chest is clear on auscultation, and her abdomen is soft and non-tender. There is a rash on both shins consistent with erythema nodosum.

Investigations:

Investigation	Result	Normal value
Haemoglobin (Hb)	131 g/l	115–155 g/l
White cell count (WCC)	9.1×10^9/l	$4–11 \times 10^9$/l
Platelets (PLT)	178×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	138 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	103 μmol/l	50–120 μmol/l
Corrected calcium (Ca)	2.71 mmol/l	2.2–2.6 mmol/l
Alanine aminotransferase (ALT)	33 IU/l	5–30 IU/l
Alkaline phosphatase (ALP)	95 IU/l	30–130 IU/l
Bilirubin	11 μmol/l	2–17 μmol/l

Which of the following investigations should be performed next?

Your answer was incorrect

A Autoimmune profile

B Computed tomography (CT) thorax

C Chest X-ray

D Sputum for microscopy, culture and sensitivity

E Ultrasound scan (USS) abdomen

Explanation

C Chest X-ray

Given the patients ethnicity, symptoms of shortness of breath and a rash consistent with erythema nodosum. The next logical step would be a chest X-ray given the suspicions of a possible diagnosis of sarcoidosis. CT thorax and pulmonary function tests can be used to confirm the diagnosis. Controversy exists over the role of steroids in the short term; however, it has been widely seen that there is some symptom reduction and a chest X-ray therefore may be indicated.

D Sputum for microscopy, culture and sensitivity

The clinical presentation would be consistent with a patient suspected of sarcoidosis. Sputum for microscopy, culture and sensitivity would not be indicated.

A Autoimmune profile

Patients with erythema nodosum may require autoimmune profiling; however, the clinical scenario is more in keeping with a diagnosis of sarcoidosis. Therefore, an autoimmune profile is not clinically indicated.

B Computed tomography (CT) thorax

The patient’s clinical features would suggest a diagnosis of sarcoidosis. A chest X-ray would be able to confirm the presence of lymphadenopathy. CT thorax is used to confirm the diagnosis and identify areas of possible fibrosis; it is not the most useful next investigation.

E Ultrasound scan (USS) abdomen

Although the liver function tests are mildly elevated, it would not be appropriate next best investigation. The clinical scenario indicates a possible diagnosis of sarcoidosis and therefore a chest X-ray is the most logical next investigation.

39772

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 64-year-old man with a history of two previous transient ischaemic attacks (TIAs) is admitted to the Emergency Department by ambulance. He has a history of obesity, hypertension and diabetes and smokes 20 cigarettes per day. His only drug treatment is metformin for diabetes. His wife called the ambulance because he fell asleep while watching a drama and was snoring loudly and she was unable to wake him. Apparently, he has recently given up driving because he feels too tired to drive any distance at all. Examination reveals a blood pressure of 155/85 mmHg and a pulse of 80 bpm and regular. He has wheeze on auscultation of the chest, but no focal signs, and his saturations are 92% on air. His body mass index (BMI) is 35 kg/m^2. Neurological examination is normal.

Investigations:

Investigations	Results	Normal Values
Haemoglobin (Hb)	133 g/l	135–175 g/l
White cell count (WCC)	8.0 × 10^9/l	4–11 × 10^9/l
Platelets (PLT)	191 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	137 mmol/l	135–145 mmol/l
Potassium (K^+)	4.3 mmol/l	3.5–5.0 mmol/l
Creatinine	132 µmol/l	50–120 µmol/l
Glucose	11.2 mmol/l	3.5–5.5 mmol/l
HbA1c	66 mmol/mol (8.2%)	< 53 mmol/mol (< 7.0%)

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Epileptic seizure

B Hypoglycaemia

C Sleep apnoea

D Transient ischaemic attack

E Transient global amnesia

Explanation

C Sleep apnoea

The history of increasing tiredness, coupled with morbid obesity, is suggestive of sleep apnoea, with episodes of daytime sleepiness well recognised to occur in situations such as TV watching and while driving the car. A sleep study incorporating monitoring for apnoeic episodes is the obvious next step. If sleep apnoea is proven and is moderate to severe, then overnight continuous positive airways pressure (CPAP) is the initial intervention of choice. For mild sleep apnoea, a mandibular advancement device may be an appropriate alternative. Over the long term, weight reduction may lead to a regression of symptoms.

B Hypoglycaemia

The random glucose of 11.2 mmol/l and the suboptimal HbA1c control at 66 mmol/l (8.2%), coupled with use of metformin monotherapy only, rule out hypoglycaemia as a possibility.

A Epileptic seizure

The lack of motor features and other signs of epilepsy, including tongue biting and loss of continence, counts against epilepsy as the underlying diagnosis.

D Transient ischaemic attack

There are no focal signs, and the features of the episode here look like more severe sleepiness against a pattern of sleep apnoea symptoms.

E Transient global amnesia

A significant period of amnesia without other neurological features, followed by complete recovery, is most suggestive of transient global amnesia. In addition, the condition is not thought to be associated with vascular risk factors.

38299

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 58-year-old man was referred to the neurology clinic by his general practitioner with a 4-month history of progressive weakness. He described difficulty walking, with heavy, stiff legs. He had particular trouble in going upstairs and rising from a chair. He also occasionally had episodes of double vision. Apart from controlled hypertension, he had no significant past medical history. He smoked 20 cigarettes per day and drank 16 units of alcohol per week. On systems enquiry, he also admitted to weight loss of one stone, a persistent cough and a dry mouth. He reluctantly admitted that he had been having some sexual difficulties with his wife recently.

On examination, he looked reasonably well. There was no finger clubbing or lymphadenopathy. Cardiorespiratory examination revealed coarse crackles consistent with COPD, and abdominal examination was unremarkable. On neurological examination, his tone and sensation were normal, with no muscle wasting. He had predominantly proximal muscle weakness and his tendon reflexes were universally absent. Apart from mild bilateral ptosis, his cranial nerves were intact. Interestingly, his power and reflexes seemed to improve after a brief period of repetitive exercise of the affected muscle.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	140 g/l	135–175 g/l
White cell count (WCC)	5.0 × 10^9/l	4–11 × 10^9/l
Platelets	300 × 10^9/l	150–400 × 10^9/l
Urea	5.2 mmol/l	2.5–6.5 mmol/l
Mean corpuscular volume (MCV)	82 fl	76–98 fl
Sodium (Na^+)	130 mmol/l	135–145 mmol/l
Potassium (K^+)	3.6 mmol/l	3.5–5.0 mmol/l
Bilirubin	10 μmol/l	2–17 μmol/l
Alanine aminotransferase (ALT)	32 IU/l	5–30 IU/l
Creatinine	120 μmol/l	50–120 μmol/l
Alkaline phosphatase	80 IU/l	30–130 IU/l
γGT	42 IU/l	5–30 IU/l

Which of the following investigations would be the most useful in making a diagnosis?

Your answer was incorrect

A Tensilon (edrophonium) test

B Serum acetylcholine receptor antibodies

C Muscle biopsy

D Anti-calcium channel antibodies

E Serum creatine kinase

Explanation

D Anti-calcium channel antibodies

This man has Lambert–Eaton myasthenic syndrome (LEMS), which is associated with an underlying small cell carcinoma of the bronchus. This is a rare paraneoplastic phenomenon which can precede radiological evidence of the tumour by up to 5 years. LEMS is a presynaptic disorder of the neuromuscular junction which is caused by antibodies to the voltage-gated calcium channels of the nerve terminal, leading to failure of acetylcholine release. The clinical features are proximal muscle weakness and depressed tendon reflexes, which can return after a few minutes of muscular contraction. Autonomic symptoms are common. The diagnosis is made electrophysiologically, and auto-antibodies are found in 90% of cases. Treatment is with 3,4-diaminopyridine, prednisolone and occasionally plasma exchange.

C Muscle biopsy

A muscle biopsy would be useful in diagnosing an inflammatory or necrotising myopathy, but these diagnoses are less likely, given that testing improves with repetition in this patient, and one would expect power to decline with repeated testing in a myopathy.

A Tensilon (edrophonium) test

A tensilon test would be diagnostic for myasthenia gravis, an autoimmune muscle condition which is associated with fatigueability, ie power and function deteriorate with repeated testing.

B Serum acetylcholine receptor antibodies

ACh receptor antibodies would be part of the diagnostic testing for myasthenia gravis, an autoimmune muscle condition which is associated with fatigueability, ie power and function deteriorate with repeated testing.

E Serum creatine kinase

Raised creatine kinase is a non-specific marker of muscle damage. It can be elevated in, for example, inflammatory or necrotising myopathy, statin-induced myopathy, myocardial infarction, rhabdomyolysis or metabolic myopathy.

32404

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 71-year-old white man is admitted to your ward with a 3-day history of pleuritic right-sided chest pain, cough productive of green sputum, shortness of breath and a fever. He has smoked 20 cigarettes a day for 50 years and is usually fit and well. He has no penicillin allergy.

On examination, he is alert, has a fever of 38.1 °C and a respiratory rate of 40/min. He is tachycardic, pulse 120/min and his blood pressure is 86/61 mmHg. On chest auscultation, he has coarse inspiratory breath sounds and bronchial breathing in the right lower lobe.

Investigations:

Investigation	**Result**	**Normal values**
Sodium (Na^+)	131 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Urea	14.2 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	137 µmol/l	50 - 120 µmol/l
C-reactive protein (CRP)	314 µg/l	< 10 mg/l
White cell count (WCC)	16.8×10^9/l	$4.0 - 11.0 \times 10^9$/l
Haemoglobin (Hb)	113 g/l	135 - 175 g/l
Platelets (PLT)	534×10^9/l	$150 - 400 \times 10^9$/l

Chest X-ray shows shadowing in the right lower zone and blunting of the right heart border.

You want to start this patient on medication for his pneumonia.

Which of the following antibiotics/antibiotic combinations would be a reasonable choice for treatment?

Your answer was incorrect

A IV cefuroxime 1.5 g tds

B Oral co-amoxiclav 625 mg tds + oral clarithromycin 500 mg bd

C IV levofloxacin 500 mg od and oral rifampicin 600 mg od

D IV co-amoxiclav 1.2 g tds and PO clarithromycin 500mg BD

E IV ciprofloxacin 400 mg bd

Explanation

D IV co-amoxiclav 1.2 g tds and PO clarithromycin 500mg BD

This patient has severe community-acquired pneumonia (CAP) as defined by a CURB 65 score of 4. British Thoracic Society (BTS) guidelines have recommended that all patients with pneumonia are assessed using CURB 65. Where possible clarithromycin should be given orally due to the risk of phlebitis with IV route.

Patients get 1 point each for

- confusion
- urea >7 mmol/l
- respiratory rate >30/min
- blood pressure <90 mmHg systolic and 60 mmHg diastolic and
- patient older than 65.

Scores 3+ indicate high mortality.

Treatment for severe CAP is of course driven by local antibiotic protocols but an acceptable choice of antibiotics would be the combination of co-amoxiclav and clarithromycin delivered IV as a first-line option.

B Oral co-amoxiclav 625 mg tds + oral clarithromycin 500 mg bd

This treatment option could be considered for mild to moderate community-acquired pneumonia if amoxicillin had recently been used and shown to be ineffective.

A IV cefuroxime 1.5 g tds

IV cefuroxime can be used for severe sepsis with oral clarithromycin if the patient is penicillin allergic.

C IV levofloxacin 500 mg od and oral rifampicin 600 mg od

Rifampicin can be added if a staphylococcal cause of pneumonia is suspected.

A 2019 UK government drug safety update advises that fluoroquinolones should now be avoided in mild to moderate infections unless other commonly recommended antibiotics are not suitable. This is due to rare reports of long lasting or potentially irreversible side effects such as tendinitis and peripheral neuropathy. Fluoroquinolones should be used with special caution in patients older than 60 years or those with renal impairment; treatment should be discontinued promptly at fist signs of serious adverse reactions.

E IV ciprofloxacin 400 mg bd

Ciprofloxacin is effective against *Pseudomonas* but has little use in the empirical treatment of CAP.

33188

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 62-year-old man who has a history of COPD, for which he takes fluticasone and salmeterol (combined) and salbutamol, comes to the Emergency Department with sudden-onset shortness of breath and right-sided pleuritic chest pain. He takes a number of medications for hypertension and ischaemic heart disease, but has recently enrolled on a pulmonary rehabilitation programme and is walking up to 2 miles per day. On examination his BP is 155/90 mmHg with pulse 92/min and regular. He has diminished breath sounds over the right-hand side on auscultation. Oxygen saturation is reduced at 90%. CXR reveals a right-sided pneumothorax with a 1.5 cm rim of air.

Which of the following is the most appropriate intervention?

Your answer was incorrect

A Admit for high-flow oxygen

B Air aspiration

C Discharge and review in 24 h

D Discharge and review in 2 weeks

E Insert size 8–14 Fr chest drain

Explanation

E Insert size 8–14 Fr chest drain

Although the rim is <2 cm, in the presence of significant symptoms (i.e. shortness of breath) a chest drain is preferred to air aspiration and is therefore the correct course of action.

D Discharge and review in 2 weeks

This is the correct course of action after successful air aspiration of a primary pneumothorax.

A Admit for high-flow oxygen

This is the correct course of action after successful air aspiration for secondary pneumothorax, where patients are usually brought into hospital for a period of 24 h.

B Air aspiration

This is the recommended intervention for a secondary pneumothorax of size 1–2 cm that isn't associated with symptoms.

C Discharge and review in 24 h

Discharge and review in 2–4 weeks is recommended after successful air aspiration of a primary pneumothorax.

40118

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumothorax

Media Expanded explanations Flashcards

A 72-year-old gentleman is referred to the Respiratory Outpatients Clinic with a 2-month history of increasing shortness of breath and cough. Over the last few weeks he has coughed up a few streaks of blood. He has also noticed difficulty getting up from a chair. He has noticed that his clothes are looser without having tried to lose weight. He takes inhaled medication but cannot remember their names.

On examination he is clubbed. Auscultation of his chest reveals reduced expansion and breath sounds on the left. Heart sounds are normal. He has a 2 cm hepatomegaly on palpation of his abdomen. He has reduced power (4/5) on hip flexion bilaterally and generally reduced tendon reflexes. Plantars both show a flexor response; proprioception and sensation are normal.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	109 g/l	135 - 175 g/l
White cell count (WCC)	12.2 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	491 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	58 mm/h	1 - 20 mm/h
Sodium (Na^+)	126 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.6 mmol/l	3.5 - 5.0 mmol/l
Urea	5.9 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	125 μmol/l	50 - 120 μmol/l
Bilirubin	49 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	120 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	441 u/l	30 - 150 u/l
Albumin	32 g/l	35 - 55 g/l
Corrected calcium (Ca^{2+})	2.67 mmol/l	2.2 - 2.7 mmol/l
Phosphate (PO_4^{3-})	0.81 mmol/l	1.1 - 1.45 mmol/l

His chest X-ray is shown;



How would you treat this patient?

Your answer was incorrect

A	Radiotherapy
B	Lobectomy
C	Pneumonectomy
D	Chemotherapy
E	Best supportive care

Explanation

D Chemotherapy

The chest X-ray shows left lower lobe collapse. There is loss of volume of the left hemi-thorax and the classical 'sail sign' behind the heart.

This man has radiological and clinical features suggestive of a lung malignancy weight loss, haemoptysis, dyspnoea and hypercalcaemia.

He also has features of paraneoplastic syndromes, making small cell lung cancer the most likely diagnosis in this case:

- hyponatraemia secondary to the syndrome of inappropriate antidiuretic hormone secretion

- the proximal weakness and hyporeflexia which, along with autonomic features, represents Lambert-Eaton myasthenic syndrome.

Platinum-based combination chemotherapy represents the mainstay of treatment for small cell lung cancer (SCLC), as per current NICE guidelines, ± radiotherapy dependent on the stage of the disease at the time of diagnosis or response to chemotherapy. Those with limited-stage disease, or with extensive disease that shows a complete response to chemotherapy at distant sites, should be considered for adjunctive radiotherapy. Because SCLC has a predilection for early metastasis, surgery has a limited role in management in comparison to other lung cancer subtypes.

C Pneumonectomy

Surgery in the case of widely metastatic SCLC would not provide a curative option, and it would be inappropriate in this case; chemotherapy represents the treatment of choice.

A Radiotherapy

Radiotherapy is an inappropriate monotherapy in the treatment of SCLC - chemotherapy is the most appropriate choice. Some patients may benefit from chemo-radiotherapy depending on their disease stage and response to chemotherapy.

B Lobectomy

Surgery in the case of widely metastatic SCLC would not provide a curative option, and it would be inappropriate in this case; chemotherapy represents the treatment of choice.

E Best supportive care

SIGN guidelines state that ‘Patients have a high objective response rate to [systemic anti-cancer therapy], with useful symptomatic improvement’ and therefore it should be at least considered in all patients with SCLC. Age should not be a factor in itself, but consideration should be given to best supportive care in those patients with, for example, poor performance status (WHO PS 3-4) or other life-limiting disease.

70134

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Lung cancer: diagnosis and management
nice.org.uk/guidance/cg121/
(http://www.nice.org.uk/guidance/cg121/)

Lung Cancer

Media

Expanded explanations

Flashcards

A 23-year-old man presents to the Emergency Department following sudden-onset left-sided pleuritic chest pain. His past medical history includes recurrent shoulder dislocations. He is on no medications.

On examination, he is tall with a long arm span. He appears short of breath and clammy. Observations reveal a heart rate of 95 bpm, BP of 110/55 mmHg, respiratory rate of 23 breaths per minute and oxygen saturations of 96% on room air. There is evidence of reduced air entry on the left-hand side and a ‘click’ on auscultation. CXR reveals a 2.5-cm pneumothorax on the left.

What is the most appropriate management of this patient?

Your answer was incorrect

A 8– to 14–Fr chest drain

B Discharge with outpatient review in 2–4 weeks

C High-flow oxygen therapy

D Needle aspiration with 16– 18–G cannula

E Observation for 24 hours

Explanation

D Needle aspiration with 16– 18–G cannula

This patient is tall and thin, with a history of joint dislocation. The suspicion is that he may have Marfan syndrome, meaning he has presented with a primary pneumothorax. The pneumothorax is over 2 cm, warranting immediate intervention. This should initially be needle aspiration with a cannula to aspirate a maximum of 2.5 litres. If this fails, the insertion of a small-bore chest drain is recommended.

C High-flow oxygen therapy

High-flow oxygen therapy should be given due to this patient’s breathlessness. In addition, oxygen may promote the resolution of a pneumothorax. However, this patient requires active intervention with needle aspiration due to his breathlessness and the size of the pneumothorax.

A 8– to 14–Fr chest drain

If this patient shows evidence of haemodynamic instability, the first-line treatment would include chest drain insertion due to the risk of this being a tension pneumothorax. Additionally, if this patient does not show clinical improvement after needle aspiration, then a chest drain would be required.

B Discharge with outpatient review in 2–4 weeks

Guidelines from the British Thoracic Society for the management of pneumothorax indicate that patients with primary pneumothoraces of below 2 cm who are haemodynamically stable and not breathless can be discharged and reviewed in the outpatient setting in 2–4 weeks.

E Observation for 24 hours

This patient is clinically unwell as a result of his pneumothorax and therefore requires prompt management. Observation would not be appropriate.

72974

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society- Management of pneumothorax
bsuh.nhs.uk/library/wp-content/uploads/sites/8/2020/06/BTS-pneumothorax-guideline.pdf
(https://www.bsuh.nhs.uk/library/wp-content/uploads/sites/8/2020/06/BTS-pneumothorax-guideline.pdf)

Pneumothorax

Media

Expanded explanations

Flashcards

A 39-year-old Irish accountant was referred to the clinic with a 4-month history of progressive breathlessness. He noticed this mainly on climbing the stairs. Over the last few months, he had been using some eye drops from the local pharmacy for dry eyes. He was also complaining of fatigue, night sweats and polyuria. He was a non-smoker and drank 20 units of alcohol a week.

On examination, he had conjunctival injection, but examination of his eyes was otherwise unremarkable. He had no rash or joint swelling. On auscultation of his chest, he had fine inspiratory crackles. Heart sounds were normal. He had no organomegaly on palpation of the abdomen. Urinalysis was unremarkable.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	101 g/l	135 - 175 g/l
White cell count (WCC)	6.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	201 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	100 µmol/l	50 - 120 µmol/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	45 mm/h	1 - 20 mm/h
Urea	7.2 mmol/l	2.5 - 6.5 mmol/l
Bilirubin	23 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	51 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	191 u/l	30 - 150 u/l
Albumin	45 g/l	35 - 55 g/l
TB test	-	

Pulmonary function tests	**Results**
FEV_1	75% predicted
FVC	70% predicted
TLCO	80% predicted

His chest X-ray is shown below:



What is the most likely diagnosis?

Your answer was incorrect

A	Tuberculosis
B	Berylliosis
C	Small cell lung carcinoma
D	Sarcoidosis
E	Langerhans cell histiocytosis

Explanation

D Sarcoidosis

The most likely diagnosis is sarcoidosis. This is a relatively young patient with breathlessness, dry eyes, fatigue, night sweats and polyuria. He has a normocytic anaemia, a raised ESR and slightly deranged liver function tests.

Sarcoid is a multi-system granulomatous disorder of unknown aetiology, with a number of different presentations. It usually presents in those under 40, and is more common in women and persons of West Indian and African descent. Presentation may be with erythema nodosum, bilateral hilar lymphadenopathy on chest radiography with fever or arthralgia (Löfgren syndrome), or more insidiously with symptoms such as dyspnoea, dry cough, fever, malaise and weight loss. The main differential diagnoses are TB and lymphoma.

The diagnosis is suggested from a chest X-ray and high-resolution CT. High-resolution CT characteristically shows features such as hilar and mediastinal lymphadenopathy, peribronchovascular nodules, ground-glass shadowing, parenchymal bands, cysts and fibrosis.

Pulmonary involvement can be classified according to the radiographic stage of the disease as below:

Stage 0: Clear chest radiograph
Stage 1: Bihilar adenopathy
Stage 2: Bihilar adenopathy and interstitial infiltrates
Stage 3: Diffuse interstitial disease
Stage 4: Advanced fibrosis.

Transbronchial biopsy is the investigation with the highest yield, as positive results are seen in 90% of patients with pulmonary sarcoidosis. Endobronchial biopsies are also useful but less sensitive. Histology typically shows non-caseating granulomas composed of macrophages, lymphocytes and epithelioid histiocytes, which fuse to form multinucleate giant cells.

Serum ACE is not specific, as raised levels are also seen in pulmonary TB, asbestosis, silicosis and lymphoma. It is elevated in about 75% of patients with untreated sarcoid. The value of using serum ACE to monitor disease activity remains unclear.

24 h urinary calcium excretion is often raised.

The lung function tests depend on severity of the disease. The tuberculin test is negative in 80% of patients with sarcoidosis, but clinicians must be mindful that anergy can also be seen in HIV-positive patients or patients who have overwhelming TB.

Treatment is rarely required in stage 0 or 1 disease. Corticosteroids should be given to patients with stage 2 or higher disease with deterioration in lung function, or patients with ocular sarcoid, hypercalcaemia, severe or persistent erythema nodosum, or myocardial and neurological manifestations of sarcoidosis. Prednisolone dose is 0.5 mg/kg for the first four weeks of treatment and is then down-titrated to the lowest maintenance dose that controls the disease. Treatment withdrawal should be considered after 6–24 months. Steroid-sparing agents such as methotrexate or azathioprine may be used for relapsing disease.

Common clinical characteristics of sarcoidosis include:

- Skin:
  Erythema nodosum – caution, not always on the shins
  Lupus pernio – red crusty lesions, often around the nose
  Annular lesions
- Polyuria:
  Secondary to hypercalciuria/hypercalcaemia (can also cause renal calculi and nephrocalcinosis)
  Secondary to central diabetes insipidus
- Cardiac:
  Ventricular arrhythmias
  Conduction defects
  Cardiomyopathy
  Congestive cardiac failure
- Neurological:
  Involvement of CNS occurs in 2%
  Cranial diabetes insipidus
- Ocular:
  Anterior uveitis

  Conjunctivitis
  Retinal lesions
  Keratoconjunctivitis sicca and lacrimal gland enlargement
  Optic neuritis
- Metabolic:
  Hypercalcaemia and hypercalciuria
- Bone and joints:
  Arthralgia
  Bone cysts
- Others:
  Hepatosplenomegaly
  Löfgren syndrome (acute sarcodosis) – triad of bihilar lymphadenopathy, arthritis and erythema nodosum.

B Berylliosis

There is no history suggestive of beryllium exposure, and bilateral hilar lymphadenopathy is not a feature of berylliosis.

A Tuberculosis

We are not given a history of contact with TB, nor a history of fevers. Inflammatory markers are normal, and tuberculin test is negative (with no history to suggest immunosuppression).

C Small cell lung carcinoma

The patient is young and is a non-smoker, making SCLC a less likely diagnosis. The chest radiograph does not show any lung lesion suggestive of a lung malignancy; it shows bilateral lymphadenopathy and diffuse interstitial infiltrates.

E Langerhans cell histiocytosis

Langerhans cell histiocytosis is a disease more commonly seen in childhood (though is increasingly recognised in adults), and in those with lung involvement, the majority being smokers.

70083

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

A 70-year-old woman presents in acute respiratory distress. She gives a history of some chest pain earlier that morning, which was worse on coughing. She is an ex-smoker of 30 pack-years (20/day for 30 years).

On examination she looks unwell. Respiratory rate is 30/min, pulse 120/min, BP 80/50 mmHg and O_2 saturation is 80% on air. Heart sounds are normal and her chest is clear.

ABG on air:

pH	7.44
PO_2	7.0 kPa
PCO_2	3.25 kPa
Base excess	-2.6
Bicarbonate	19.5 mmol/l

You manage to get an urgent CT within the hour, shown below.



What is the most likely diagnosis?

Your answer was incorrect

A Congestive cardiac failure

B Myocardial infarction

C Pneumonia

D Pulmonary embolism

E Pneumothorax

Explanation

D Pulmonary embolism

This lady has had a massive pulmonary embolus, as shown on the CTPA. She presents with shortness of breath and tachypnoea, and her arterial blood gases show she has type I respiratory failure; note the low CO_2 – common in PE.

The CT shows large filling defects in the left and right main pulmonary arteries.

PE should be managed according to British Thoracic Society guidelines on the treatment of massive PE. The guidelines state that if a PE is sufficiently severe to cause circulatory collapse, thrombolysis should be given as early as possible. The current guidelines recommend alteplase as the thrombolytic agent of choice, as it can be given to the hypotensive patient, as in this example.

B Myocardial infarction

We are not given a history of cardiac chest pain, and the CTPA is clearly demonstrative of PE as the cause of her symptoms.

A Congestive cardiac failure

The patient’s symptoms are not in keeping with cardiac failure, and her clear chest on examination makes this a less likely option. The CTPA is also clearly demonstrative of PE as the cause of her symptoms.

C Pneumonia

The patient would be expected to demonstrate symptoms such as productive cough, fever, etc. if the diagnosis were pneumonia. Her clear chest on examination also makes this a less likely option. The CTPA is also clearly demonstrative of PE as the cause of her symptoms.

E Pneumothorax

Clear chest on examination, and a CTPA demonstrative of PE, make pneumothorax a less likely option. Pneumothorax would be demonstrated by CT if present.

70080

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society guidelines for the management of suspected acute pulmonary embolism
brit-thoracic.org.uk/index.asp
(http://www.brit-thoracic.org.uk/index.asp)

A 31-year-old, previously healthy man presents with a 6-month history of nasal congestion and sinusitis. Over a 6-month period he was treated with antibiotic therapy, but his symptoms persisted. His current complaints include frontal headache and a nose bleed.

On examination, he is pyrexial and dyspnoeic. He has a rash on his legs (image below) and a right foot drop.

Investigations:

Investigation	**Result**	**Normal Values**
Sodium (Na^+)	140 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.5 mmol/l	3.5 - 5.0 mmol/l
Urea	15 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	230 µmol/l	50 - 120 µmol/l
Haemoglobin (Hb)	108 g/l	135 - 175 g/l
White cell count (WCC)	12.0×10^9/l	$4.0 - 11.0 \times 10^9$/l
Erythrocyte sedimentation rate (ESR)	84 mm/h	1 - 20 mm/h
Urine dipstick	Blood ++ Protein ++	
Chest X-ray	Cavitating lesion in the right upper lobe	

Rash:



What is the most likely diagnosis?

Your answer was correct

A Eosinophilic Granulomatosis with Polyangiitis

B Granulomatosis with Polyangiitis (GPA)

C Polyarteritis nodosa

D Post-streptococcal glomerulonephritis

E Subacute bacterial endocarditis

Explanation

B Granulomatosis with Polyangiitis (GPA)

GPA is a necrotising vasculitis involving the respiratory tract and kidneys. It causes epistaxis, sinusitis, destruction of the nasal cartilage and glomerulonephritis and kidney disease. The serum contains antibodies reacting with the cytoplasm of neutrophils (cANCA).

GPA is an idiopathic, systemic inflammatory disease characterised by the presence of granulomas, necrosis and/or vasculitis. Although GPA typically affects the upper and lower airways and the kidneys, it may involve any organ system. GPA is classified, along with microscopic polyangiitis (MPA) and Churg–Strauss syndrome (CSS), as a primary systemic small-medium-sized vessel vasculitis associated with anti-neutrophil cytoplasmic antibodies (ANCAs). However, GPA can also affect medium and even large arteries, and may lack an association with ANCAs. It is a renal/pulmonary disease which often presents with haemoptysis, sinusitis, oral lesions and otitis.

The diagnosis of GPA is based on a combination of clinical, laboratory and, if necessary, pathological features. If a typical clinical picture is associated with a positive ANCA finding with specificity for proteinase 3 (PR3), the diagnosis of WG can be presumed. It remains incumbent for the clinician to be certain that GPA 'mimics', especially granulomatous infection, are not present. In the setting of only a moderate suspicion of GPA and a negative ANCA result, it would be judicious to pursue histological support for the diagnosis. The outcome of GPA has been dramatically altered by treatment with glucocorticoids (GCs) and cyclophosphamide (CP).

A Eosinophilic Granulomatosis with Polyangiitis

Eosinophilic Granulomatosis with Polyangiitis (eGPA), formally Churg–Strauss syndrome, typically presents with respiratory symptoms such as asthmatic wheeze. Investigations in eGPA will show microscopic evidence of red cells and casts in the urine, with protein on dipstick. In the full blood count there would be either prominent eosinophilia (>10% total white cell count or absolute count of $>1.5 \times 10^9$/l) or a predominance of eosinophils in the biopsy of affected tissues. This eosinophilia is highly steroid-responsive and resolves rapidly on treatment.

C Polyarteritis nodosa

Polyarteritis nodosa is a systemic vasculitis. There would be palpable purpura and livedo reticularis. Nerve involvement may include paraesthesia. There is no sinusitis. Investigations in polyarteritis nodosa may include a raised creatinine kinase due to muscle injury, raised ESR and ANCA negative. Hepatitis B would be positive in the minority. The diagnosis is made on angiography, which would show arterial aneurysms in the renal or mesenteric tree.

D Post-streptococcal glomerulonephritis

This generally presents with sore throat. The streptococcal titre would be positive. There would be no cavitating lung lesions on chest X-ray, as present in this case.

E Subacute bacterial endocarditis

You would expect mention of a murmur with endocarditis. Duke’s criteria are used for diagnosis of endocarditis.

70770

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Granulomatosis with Polyangiitis

Media

Expanded explanations

Flashcards

A 70-year-old lady is referred to the Medical Clinic by her general practitioner (GP) following shortness of breath. She has been seeing her GP over the last couple of months for increasing shortness of breath on exercise. Her GP had done some investigations but still could not explain her shortness of breath on exercise. She has had a dry cough which has been on and off for the last 4 months. On examination, her blood pressure is 135/80 mmHg, pulse 70 bpm and regular. Her jugular venous pulse is not raised and her heart sounds are normal. Her breathing is noisy but her chest is clinically clear on auscultation.

Which investigation is likely to be most helpful in determining where any obstruction lies?

Your answer was correct

A Chest X-ray

B Bronchoscopy

C Flow volume loop

D Polysomnography

E Arterial blood gases

Explanation

C Flow volume loop

This patient likely has a partial upper airway obstruction. The causes in this age group would include: a tumour, a multinodular goitre or an aneurysm. A flow loop volume would show characteristic patterns with intrathoracic/extrathoracic obstruction.

A Chest X-ray

This patient has signs suggestive of partial upper airway obstruction. A chest X-ray would not help determining where the upper airway obstruction lies. A significant multinodular goitre may, however, be an incidental finding on chest X-ray.

B Bronchoscopy

Though bronchoscopy is important in the diagnosis and management of central airway obstruction, its role in the diagnosis of upper airway obstruction is limited.

D Polysomnography

This test would not help determine the location of the upper airway obstruction. Polysomnography is used in the diagnosis of sleep disorders.

E Arterial blood gases

In the context of the history given, there is no indication for an arterial blood gas.

7506

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 42-year-old homeless man is admitted with confusion, fever and shortness of breath. He is well known to the Emergency Department and has presented on a number of occasions with alcohol intoxication. On examination he is pyrexial at 38.6°C. He has a cough productive of purulent and blood-stained sputum. Auscultation of the chest reveals findings consistent with right-sided consolidation.

His chest X-ray shown below.



Which of the following is the most likely diagnosis?

Your answer was correct

A *Chlamydia psittaci*

B *Haemophilus influenzae*

C *Klebsiella pneumoniae*

D *Staphylococcus aureus*

E *Streptococcus pneumoniae*

Explanation

C *Klebsiella pneumoniae*

This patient has pneumonia with evidence of lung abscess formation. Given his history of alcoholism and the fact there is evidence of cavitation in the right upper lobe on chest X-ray, *Klebsiella* is a likely cause. Third-generation cephalosporins or quinolones provide coverage for community-acquired infection but macrolides do not. Mortality is particularly high in patients with a history of alcoholism, potentially as a result of poor nutrition and defective immunity associated with chronic excess alcohol consumption.

Image source: http://www.svuhradiology.ie/case-study/cavitating-pneumonia/

A *Chlamydia psittaci*

C. psittaci is associated with atypical pneumonia as a result of exposure to infected birds. Symptoms may initially mimic typhoid fever with fever, joint pains, gastrointestinal (GI) disturbance, headache and lymphopaenia. A cough and evidence of respiratory tract infection develop later.

B *Haemophilus influenzae*

H. influenzae infection is associated primarily with exacerbations of chronic obstructive pulmonary disease (COPD) rather than the cavitating pneumonia seen here.

D *Staphylococcus aureus*

S. aureus is also associated with cavitation and abscess formation, but occurs more commonly after an episode of influenza.

E *Streptococcus pneumoniae*

S. pneumoniae is a common cause of community-acquired pneumonia, although it is not usually cavitating.

32429

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 54-year-old stonemason was referred to the clinic with a 3-year history of progressive breathlessness. He is an ex-smoker (40 pack-years) having stopped 14 years ago.

On examination at that time, he looked well, was not clubbed and there was no lymphadenopathy. His chest was hyper-expanded and he had basal inspiratory crackles.

His spirometry revealed:

FEV_1	38% predicted
FVC	70% predicted
KCO	55% predicted

His chest X-ray from the original referral is shown below:



Three years later, he presents to the Emergency Department with increasing shortness of breath, cough and wheeze. Over the last year, he has had recurrent chest infections and now has a cough productive of green sputum.

On examination, respiratory rate 30/min, temperature 36.2 ºC, blood pressure (BP) 160/100 mmHg, pulse 130/min, SpO_2 57% on air.

He is breathing through pursed lips. Auscultation of his chest reveals expiratory wheeze and crackles throughout.

Blood gases on 28% oxygen:

Investigation	Result	Normal Values
pH	7.22	7.35 - 7.45

PO_2	7.9 kPa	> 10.5 kPa
PCO_2	8.0 kPa	4.7 - 6.0 kPa
Bicarbonate	20 mmol/l	22 - 29 mmol/l
Base excess	-3	(-2 -> +2)

What do the blood gases show?

Your answer was incorrect

A Metabolic acidosis

B Respiratory acidosis

C Respiratory acidosis with metabolic compensation

D Mixed respiratory and metabolic acidosis

E Metabolic acidosis with respiratory compensation

Explanation

D Mixed respiratory and metabolic acidosis

This patient’s blood gases show a picture of mixed respiratory and metabolic acidosis. The chest X-ray shows predominantly upper lobe fibrosis consistent with chronic exposure to silica.

It is important when assessing blood gas results to do so in a systematic manner to make sure the correct diagnosis is reached.

Initially, looking at the pH in this case, it can be seen that there is an acidosis. pH normally ranges between 7.35 and 7.45, with values lower than 7.35 representing acidosis and values greater than 7.45 representing alkalosis.

When it has been established that an alkalosis or acidosis exists, look next at the CO_2 and bicarbonate to establish the cause, and the presence of compensation (if any).

pCO_2 is normally 4.5–6.0 kPa. As CO_2 is acidic, values greater >6.0 signify acidosis (respiratory acidosis) while values lower <4.5 signify alkalosis (respiratory alkalosis).

Bicarbonate normally ranges between 22 and 28 mEq/l. As bicarbonate is basic, values lower than 22 signify acidosis (metabolic acidosis) while values greater than 28 signify alkalosis (metabolic alkalosis).

In this case, it can be seen that CO_2 is elevated *and* bicarbonate is reduced, indicating a mixed respiratory and metabolic cause of this patient’s acidosis.

Compensation can be seen when there are abnormalities of either CO_2 or bicarbonate in the ‘opposite’ direction to the change in pH. For example, in the case of metabolic acidosis (with low pH and low bicarbonate) there may be respiratory compensation in which the patient hyperventilates to ‘blow off’ CO_2 and, as such, a reduced CO_2 is seen (which in isolation would normally signify a respiratory alkalosis).

B Respiratory acidosis

Because his CO_2 is raised, in addition to a low bicarbonate and base excess this option is incorrect.

A Metabolic acidosis

Because his CO_2 is raised, in addition to a low bicarbonate and base excess this option is incorrect.

C Respiratory acidosis with metabolic compensation

If he had metabolic compensation, his base excess and bicarbonate would be raised in response to the acidosis caused by the raised CO_2.

E Metabolic acidosis with respiratory compensation

In respiratory compensation, the carbon dioxide is reduced to compensate for acidosis caused by reduced bicarbonate.

70920

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 30-year-old man, who is fit and well, is admitted with sudden-onset shortness of breath associated with sharp chest pain made worse with coughing. The symptoms began suddenly while the patient was working on his computer.

He is not on any regular medications. His blood pressure (BP) is 145/70 mmHg, and his pulse is 70 bpm and regular. After clinical examination of the patient, a chest X-ray is requested, which is shown below:



Which of the following signs is most likely to have been elicited during the clinical examination?

Your answer was incorrect

A Deviation of the trachea

B Bilateral reduced expansion

C Saturation of 90% on air

D Dull percussion note

E Unilateral decreased breath sounds

Explanation

E Unilateral decreased breath sounds

This is another classical finding that occurs due to failure of the affected lung to expand appropriately.

Image source: https://commons.wikimedia.org/wiki/File:PNTX.png

D Dull percussion note

Localised hyperresonance is a common finding that occurs due the presence of air in conjunction with reduced amounts of tissue.

A Deviation of the trachea

A pneumothorax occurs due to the entry of air in the intrapleural space and the subsequent partial or total collapse of the lung. Pneumothoraces are often spontaneous and are more likely to occur in young, thin men. A secondary pneumothorax is more likely to occur in the presence of an underlying lung disease. Patients may be asymptomatic or they may present with sudden onset of shortness of breath and pleuritic chest pain. The classical signs of a pneumothorax on clinical examination of the chest include: reduced expansion on the affected side, hyperresonant percussion note and diminished breath sounds.

A deviated trachea (away from the affected side) is seen in cases of a tension pneumothorax and is usually accompanied by a combination of respiratory distress, tachycardia, tachypnoea and hypotension.

B Bilateral reduced expansion

Unilateral reduced expansion is a very common finding in a pneumothorax. Due to the entry of air, the associated lung collapses and cannot inflate, leading to reduced chest expansion on the affected side.

C Saturation of 90% on air

The patient described has unremarkable BP and heart rate measurements. Given that and the fact that spontaneous pneumothoraces are often small, normal saturation levels would be an expected finding.

71456

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumothorax

Media

Expanded explanations

Flashcards

A 90-year-old man is seen in the Respiratory Clinic with an 18-month history of progressive shortness of breath and dry cough. Before this, he was able to complete a full round of golf; he is now unable to walk further than 200 m. He has been a heavy smoker all his life. He has no known asbestos exposure. On examination, he is breathless and has a photosensitive rash on his face and neck, as well as clubbing. His heart rate is irregularly irregular. Auscultation of his chest reveals fine inspiratory crackles bibasally.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	121 g/l	135 - 175 g/l
White cell count (WCC)	9.2 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	6.1 × 10^9/l	1.5 - 7.0 × 10^9/l
Platelets (PLT)	203 × 10^9/l	150 - 400 × 10^9/l
Thyroid-stimulating hormone (TSH)	7.1 mu/l	0.4 - 5.0 mu/l

His CXR is shown below:



Spirometry:

Investigation	**Result**	**Predicted values**
Forced expiratory volume in one second (FEV_1)	1.5 L	2.0 - 2.4 L
Forced vital capacity (FVC)	2.0 L	3.2 - 3.8 L
Carbon monoxide transfer coefficient (KCO)	74% predicted	

What is the most likely cause of this patient’s breathlessness?

Your answer was correct

A Bronchial alveolar carcinoma

B Interstitial lung disease

C Chronic obstructive pulmonary disease

D Lymphangioleiomyomatosis

E Pulmonary oedema

Explanation

B Interstitial lung disease

This patient has interstitial lung disease. He has gradual-onset shortness of breath and dry cough and has clubbing and fine inspiratory crackles. Lung function tests show a mixed restrictive/obstructive picture and he has type I respiratory failure. The CXR shows small lung fields and reticular nodular shadowing.

The patient has a photosensitive rash on his face, atrial fibrillation and abnormal thyroid function, suggesting he is on amiodarone. Interstitial lung disease is an important side-effect of long-term amiodarone administration. Pulmonary fibrosis is a known side-effect of several other drugs, including beta-blockers, nitrofurantoin and methotrexate.
Diagnosis is made by a combination of clinical findings, lung function tests, CXR, bronchoalveolar lavage (BAL) and high-resolution CT.
Other side effects of amiodarone include:

- hyper- and hypothyroidism
- peripheral sensory neuropathy
- hepatitis
- photosensitivity
- ataxia
- corneal microdeposits (reversible)
- metallic taste
- arrhythmias (torsades de pointe).

Drugs causing interstitial lung disease include:

- antibiotics – nitrofurantoin
- cardiovascular – amiodarone, beta-blockers
- chemotherapeutic – bleomycin, busulfan, cyclophosphamide, carmustine, lomustine, methotrexate, melphalan, mitomycin C, nitrosoureas, sulfasalazine, 5-fluorouracil (5FU)
- drugs used to treat rheumatoid arthritis – sulfasalazine, penicillamine, cyclophosphamide, methotrexate, gold

- drugs used to treat migraine – ergotamine, ergots, methysergide

A Bronchial alveolar carcinoma

The long duration of symptoms, small lung fields and reticular nodular shadowing, with absence of a focal lesion on the CXR, coupled with the dry cough, are more consistent with interstitial lung disease than with bronchial alveolar carcinoma.

C Chronic obstructive pulmonary disease

While chronic obstructive pulmonary disease does cause progressive breathlessness, a productive cough, obstructive lung function and hyperinflated lung fields on this patient's CXR would be expected.

D Lymphangioleiomyomatosis

This condition usually affects women in their childbearing years and is associated with cystic destruction of the lungs and is not likely in this patient.

E Pulmonary oedema

This is a reasonable differential diagnosis based on the history. However, productive sputum may be expected and the CXR shows interstitial lung disease, not pulmonary oedema.

70116

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 72-year-old man is referred to the Respiratory Outpatients Clinic with a 2-month history of increasing shortness of breath and cough. Over the last few weeks, he has coughed up a few streaks of blood. He has also noticed difficulty getting up from a chair, and that his clothes have been looser recently despite his not trying to lose weight.

On examination, he is clubbed. Auscultation of his chest reveals reduced expansion and breath sounds on the left. Heart sounds are normal. He has a 2-cm hepatomegaly on palpation of his abdomen. He has reduced power (4/5) on hip flexion bilaterally and generally reduced tendon reflexes. Plantars both show a flexor response; proprioception and sensation are normal.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	109 g/l	135 - 175 g/l
White cell count (WCC)	12.2 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	491 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
Erythrocyte sedimentation rate (ESR)	58 mm/h	1 - 20 mm/h
Sodium (Na^+)	126 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.6 mmol/l	3.5 - 5.0 mmol/l
Urea	5.9 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	125 µmol/l	50 - 120 µmol/l
Bilirubin	49 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	120 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	441 u/l	30 - 150 u/l
Albumin	32 g/l	35 - 55 g/l
Corrected calcium (Ca^{2+})	2.74 mmol/l	2.2 - 2.7 mmol/l
Phosphate (PO_4^{3-})	0.81 mmol/l	1.1 - 1.45 mmol/l

His chest X-ray is shown below:



What does the chest X-ray show?

Your answer was correct

A	Left lower lobe consolidation
B	Left lower lobe collapse
C	Lingular consolidation
D	Left upper lobe collapse
E	Left apical pneumothorax

Explanation

B Left lower lobe collapse

The chest X-ray shows left lower lobe collapse. There is loss of volume of the left hemi-thorax, the classical ‘sail sign’ (double left heart border) behind the heart, and loss of the normal outline of the left hemi-diaphragm.

A Left lower lobe consolidation

The opacification seen in the left lower zone has a smooth border and is more in keeping with the ‘sail sign’ of left lower lobe collapse. Consolidation would have a more heterogeneous appearance and air bronchograms may be visible.

C Lingular consolidation

The left cardiac border can be visualised, making lingular consolidation unlikely. In lingular consolidation, loss of the left heart silhouette would be expected.

D Left upper lobe collapse

Left upper lobe collapse presents as veil-like shadowing over the remaining inflated lung.

E Left apical pneumothorax

Lung markings are visible throughout all lung fields and no lung edge is seen.

70131

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct: 82

Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Cancer

Media

Expanded explanations

Flashcards

A 69-year-old female was taken to the Emergency Department with a reduced GCS score; no history was available from relatives for the patient in A&E. The ambulance crew said that the call was put out by her husband as she had become increasingly short of breath. When they arrived she looked breathless and cyanosed. She denied any chest pain at that time.

On examination he had a GCS score of 10 (M5, V2, E3), respiratory rate 10/min, BP 130/70 mmHg, pulse 120/min, temperature 36.9°C, SaO_2 94% on 15 l oxygen. Auscultation of the chest revealed an expiratory wheeze throughout.

Heart sounds were normal and abdominal examination unremarkable. There was no obvious focal neurological abnormality and both plantars showed a flexor response.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	173 g/l	135 - 175 g/l
White cell count (WCC)	8.2 ×10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	400 ×10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
C-reactive protein (CRP)	21 mg/l	< 10 mg/l
Electrocardiogram (ECG)	Sinus tachycardia	

In her notes you find a flow-volume curve from 1 year ago shown below.



Her chest x-ray is shown below.



ABGs on high-flow oxygen on this admission:	
pH	7.29
PCO_2	10.91 kPa
PO_2	12.39 kPa
Bicarbonate	40.6 mmol/l
Base excess	10.4

What does the arterial blood gas analysis show?

Your answer was incorrect

A Mixed respiratory and metabolic acidosis

B Acute respiratory acidosis

C Metabolic acidosis

D Acute on chronic respiratory acidosis

E Metabolic alkalosis

Explanation

D Acute on chronic respiratory acidosis

This patient has an acute on chronic respiratory acidosis. The high bicarbonate and base excess show that there is an underlying chronic respiratory acidosis with renal compensation. The acidosis and high CO_2 indicate an acute respiratory acidosis. If this was acute without a chronic element then, for a PCO_2 of 10.91 one would expect the pH to be lower.

The diagnosis of COPD is suggested from the CXR, showing hyperexpansion and flattened diaphragms, and flow-volume curve, which shows the classic shape for COPD. In some patients with COPD, respiratory drive depends on their degree of hypoxia, rather than the usual dependence on hypercapnia. Although it is important to prevent life-threatening hypoxia, uncontrolled oxygen therapy should be used with caution. This patient has a suppressed respiratory drive and carbon dioxide narcosis secondary to high-flow oxygen. She was able to tell the ambulance crew that she had no chest pain at home; she was then given high-flow oxygen and her GCS dropped. Note on arrival at the Emergency Department that she scored 2/5 for verbal response and therefore would not be able to give any information. The immediate management is therefore to reduce the inspired oxygen concentration and repeat the arterial blood gases. If by doing this the patient becomes hypoxic or remains acidotic, they should then be referred for non-invasive ventilation.

Image source: https://radiopaedia.org/cases/emphysema-9?lang=gb

B Acute respiratory acidosis

In acute respiratory acidosis, one would expect the bicarbonate and base excess to be lower. The raised bicarbonate implies a chronicity to the respiratory acidosis with renal compensation.

A Mixed respiratory and metabolic acidosis

The ABG would be expected to show low bicarbonate and base excess if a metabolic component to the acidosis were present.

C Metabolic acidosis

The raised CO_2 implies a respiratory component to the acidosis. In a metabolic acidosis, the bicarbonate and base excess would be expected to be low.

E Metabolic alkalosis

The ABG demonstrates a low pH and raised CO_2, neither of which is in keeping with metabolic alkalosis. The bicarbonate and base excess are raised, likely representing renal compensation to a chronic respiratory acidosis.

70128

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 54-year-old Asian female, who had emigrated from Pakistan 9 months previously, presents to the Emergency Department with retrosternal discomfort at the site of her sternotomy wound. She had undergone aortic valve replacement two months previously and subsequently developed a keloid scar. A wound infection at the incision site is diagnosed and she is admitted by the surgical team for removal of the sternotomy sutures.

At 15 days post-procedure, she develops right pleuritic chest pain with a temperature of 38°C, which does not respond to amoxicillin. A CXR is performed which shows a raised right hemidiaphragm and a right hilar mass (also seen on the lateral film).

Two days after starting antibiotic therapy, the patient also develops dorsal back pain for which a CT scan of the thorax and a CT scan of her thoracic spine is performed. The result of the chest CT scan confirms the hilar lymphadenopathy as well as showing a mass within the right psoas muscle. The CT scan of her mid-thoracic spine shows the destruction of the body and the process of T8. She goes onto have a bone scan which identifies several 'hot spots', including over the thoracic spine.

What organism is most likely to be cultured from this abscess?

Your answer was correct

A *Escherichia coli*

B *Mycobacterium tuberculosis*

C *Streptococcus bovis*

D *Candida albicans*

E *Staphylococcus aureus*

Explanation

B *Mycobacterium tuberculosis*

Pott's disease is also known as tuberculous spondylitis and is one of the oldest diseases to afflict humankind. It is usually secondary to an extra-spinal source of infection, primarily from the lungs or the mediastinal glands through the blood stream. The classical symptoms of tuberculosis of the spine are pain, rigidity, deformity, cold abscess and paraplegia (following vertebral collapse). Although any vertebra may be involved, it is usually the lower thoracic and upper lumbar vertebrae that are affected. Patients also report constitutional symptoms, including fever and weight loss. This patient had tuberculosis of the lung which had spread to her spine. Although upper lobe involvement is most common, tuberculosis can be found in any part of the lung.

A *Escherichia coli*

This organism is associated with secondary psoas abscess in patients with inflammatory bowel disease or renal sepsis, but would not explain her respiratory or vertebral symptoms.

C *Streptococcus bovis*

This organism is associated with gastrointestinal malignancy, but would not explain her constellation of respiratory, vertebral and psoas symptoms.

D *Candida albicans*

Systemic candidial infection is most commonly seen in immunocompromised patients such as those with HIV, and is a less likely diagnosis than TB given the respiratory, vertebral and psoas features seen.

E *Staphylococcus aureus*

This is the most common cause of primary psoas abscess, and secondary psoas abscess where skeletal infection is the primary cause. In this case, this patient probably has pulmonary TB which has spread to her vertebrae and psoas.

6810

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219

Responses Total: 301

Responses - % Correct: 27%

Tuberculosis

Media

Expanded explanations

Flashcards

You are asked to review a 58-year-old care worker who has complained of diffuse, non-specific right-sided chest pain increasing over the past few months. She smokes 20 cigarettes per day and has no past medical history of note. Her husband works as a plumber and had worked as a boiler engineer for many years before that. A chest X-ray taken 12 months ago had shown pleural plaques, but was otherwise normal.

Chest x-ray taken following clinic:



Which of the following diagnoses fits best with the features of this case?

Your answer was incorrect

A Adenocarcinoma of the bronchus

B Small cell carcinoma of the bronchus

C Mesothelioma

D Benign pleural plaques

E Asbestosis

Explanation

C Mesothelioma

The chest x-ray shows opacification of the right lower zone with pleural plaques on the left side. The appearance of this change over the past year, coupled with the history of possible asbestos exposure in the family, is suggestive of mesothelioma. A spouse can contract the disease through secondary exposure such as handling the laundry of a spouse that has already been exposed. Patients may present with a history of diffuse, non-specific chest pain and profuse sweating. The diagnosis is a possibility in any patient with a suggestion of asbestos exposure and who presents with pleural thickening or pleural effusion. Diagnosis is with pleural fluid aspiration under ultrasound guidance and CT scanning. Repeated aspiration should be avoided if the diagnosis is suspected, due to the possible risk of tumour tracking. In addition, pleural biopsy is often helpful in making the diagnosis; this should be CT or ultrasound guided, if possible, rather than using a blind technique.

Image source: https://radiopaedia.org/cases/mesothelioma?lang=gb

A Adenocarcinoma of the bronchus

The presentation with initial pleural disease is more consistent with mesothelioma than a bronchial carcinoma.

B Small cell carcinoma of the bronchus

Small cell carcinoma primaries are located centrally in 90% or more of cases, and this doesn’t therefore fit with the clinical picture here.

D Benign pleural plaques

The rapid evolution of the CXR over the course of the past year to now show an area suggestive of an infiltrative lesion doesn’t fit with benign pleural plaque disease.

E Asbestosis

Asbestosis fits with a picture of pulmonary fibrosis, unlike that seen here.

9280

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Mesothelioma and Other Asbestos Related Conditions

Media

Expanded explanations

Flashcards

A 62-year-old man who works in a factory which sorts and recycles metals, comes to the clinic for review. He has progressive shortness of breath and a dry cough, which has worsened significantly over the last 6 months. He smokes 5 cigarettes per day. His GP has trialled a Salbutamol inhaler for him with no improvement. On examination his BP is 132/88 mmHg; pulse is 80/min and regular. He has inspiratory crackles on auscultation, worse in the mid- and lower zones.

Investigations;

Hb	131 g/l
WCC	9.4 x10^9/l
PLT	201 x10^9/l
Na^+	137 mmol/l
K^+	4.5 mmol/l
Creatinine	95 micromol/l
Glucose	5.4 mmol/l
FEV_1	65% of predicted
FVC	70% of predicted
CXR	Interstitial shadowing affecting both lung fields

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Berylliosis

B COPD

C Colophony lung

D Idiopathic pulmonary fibrosis

E Silicosis

Explanation

A Berylliosis

The answer is Berylliosis -

Berylliosis is a granulomatous lung disorder, which in chronic form leads to restrictive lung disease. One occupation known to be associated with beryllium exposure is metal recycling. The beryllium lymphocyte proliferation test (BeLPT) is the laboratory investigation of choice. High resolution CT reveals ground glass changes consistent with pulmonary fibrosis, and nodule formation. Lung biopsy reveals granuloma formation indistinguishable from sarcoidosis. Prevention is key, given the similarities of this disorder to sarcoidosis. Some response to corticosteroids is often seen.

C Colophony lung

Colophony lung is incorrect, it is associated with a primarily asthmatic type picture, and we are not given information that active soldering is taking place at the metal recycling plant.

B COPD

COPD is incorrect, it is associated with an obstructive, rather than the restrictive lung disease pattern seen here, and his tobacco exposure is relatively modest.

D Idiopathic pulmonary fibrosis

Idiopathic pulmonary fibrosis is incorrect, it is unlikely given the history of potential exposure to heavy metals.

E Silicosis

Silicosis is incorrect, it is associated with occupations where exposure to silica dust is likely, such as in patients who work as stone masons.

40404

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Berylliosis

Expanded explanations

A 64-year-old man was referred to the clinic with insomnia. He complained of falling asleep during the day and having an early morning headache. He had a past history of a cholecystectomy and gout. He took no other regular medication. Examination was unremarkable.

Investigation	Result	Normal Values
Weight	115 kg	
Height	175 cm	
Haemoglobin (Hb)	161 g/l	135 - 175 g/l
White cell count (WCC)	6.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	94 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	101 fl	80 - 100 fl
Thyroid-stimulating hormone (TSH)	1.2 mu/l	0.4 - 5.0 mu/l
Glucose	8.2 mmol/l	3.9 - 7.1 mmol/l

What would the optimal management of this patient be over the long term?

Your answer was correct

A Refer for non-invasive ventilation

B Diuretics

C Advise to lose weight

D Refer to ENT surgeons

E Doxapram

Explanation

C Advise to lose weight

This patient has obstructive sleep apnoea (OSA) – insomnia, daytime somnolence, morning headache and obesity. Other symptoms include poor concentration during the day and partners may describe snoring followed by apnoeic episodes.

Diagnosis is made using the Epworth score – which is a measure of daytime somnolence – and sleep studies, which would show apnoeic/hypopnoeic episodes associated with desaturations, increase in heart rate and arousal from sleep. Treatment will depend on severity of symptoms and desaturations. In most cases, weight loss is advisable.

A mandibular advancement device may control symptoms in mild to severe OSA syndrome, such as nocturnal non-invasive ventilation (NIV) using continuous positive airway pressure (CPAP). Early morning headaches can be caused by a transient hypercapnia. This normally corrects to normal when the patient is awake. If a patient has hypercapnia in waking hours (and not just on an early morning blood gas) then overlap syndrome with obesity hypoventilation syndrome should be considered, although severe OSA may cause persistent hypercapnia in some cases. Patients with persistent hypercapnia should be considered for bi-level positive airway pressure (BiPAP).

It is important to exclude underlying causes – this patient has a history of gout which, together with the raised MCV and low platelets, suggests alcohol consumption, which can precipitate the problem. Hypothyroidism, acromegaly and sedating drugs also need to be excluded. Retrognathia can cause OSA and large tonsils may obstruct the airway.

A Refer for non-invasive ventilation

Although NIV is optimal for relief of symptoms, over the long term, given significant risk of type 2 diabetes and diseases associated with insulin resistance, weight loss is preferred long term.

B Diuretics

Diuretics would be an appropriate treatment for pulmonary oedema.

D Refer to ENT surgeons

Surgery is used as a last resort as a treatment for OSA and is often unsuccessful.

E Doxapram

Doxopram is a respiratory stimulant with a vast array of side effects previously used to treat acute type 2 respiratory failure. It has been superceded by non-invasive ventilation now.

70140

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Obstructive Sleep Apnoea

Media

Expanded explanations

Flashcards

A 39-year-old woman from Zimbabwe presents to the Emergency Department with a 5-day history of increasing shortness of breath and dry cough. She has a 1-day history of sudden onset of sharp chest pain which is worse on coughing, movement and deep inspiration. She describes having had night sweats and a poor appetite for two months. Her weight has decreased by about 5 kg. She has a past medical history of genital herpes and depression. She is married and has two children who live in Zimbabwe. She moved to England 2 years ago and has not travelled since. She is a smoker of 20/day and drinks little alcohol.

On examination, she looked unwell. Observations: temperature 38.3 °C, blood pressure (BP) 108/72 mmHg, pulse 120/min, regular, respiratory rate 30/min. Auscultation of her chest revealed inspiratory crackles bi-basally.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	102 g/l	135 - 175 g/l
White cell count (WCC)	3.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	439 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	141 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	75 μmol/l	50 - 120 μmol/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
C-reactive protein (CRP)	73 mg/l	< 10 mg/l
Urea	5.0 mmol/l	2.5 - 6.5 mmol/l
Glucose	5.7 mmol/l	3.9 - 7/1 mmol/l

Blood gases on 80% oxygen:

pH	7.46
PCO_2	3.44 kPa
PO_2	7.77 kPa
Bicarbonate	17.6 mmol/l
Base excess	–3.5

Her chest X-ray is shown below:



What treatment should be started?

Your answer was incorrect

A Intravenous co-amoxiclav and clarithromycin

B Anticoagulation with warfarin

C Prednisolone

D Intravenous ganciclovir

E Intravenous high-dose co-trimoxazole ± steroids

Explanation

E Intravenous high-dose co-trimoxazole ± steroids

For the purpose of the exam, people who have lived abroad (especially Africa and South America), businessmen who work abroad, men who have sex with men (MSM) and intravenous drug users, are more likely to have HIV.

This lady has HIV – the history of living in Zimbabwe and the low white cell count are the clues. *Pneumocystis jirovecii* is the most common opportunistic infection to cause pneumonia in AIDS – especially when the CD4 count is <200/mm^3. It accounts for about 50% of cases of pneumonia in AIDS and 40% of all AIDS-defining illnesses.

Patients usually present with a fever, dry cough and breathlessness. They are usually hypoxic and desaturate on exercise.

In *P. jirovecii* pneumonia (PJP) the chest X-ray usually shows bilateral interstitial shadowing and cysts in the mid and lower zones. However, the CXR may be normal. Pneumothorax (because the cysts rupture) may be present in up to 10%.

Diagnosis is now usually made by PCR of sputum or BAL fluid. Other diagnostic methods include staining induced sputum or BAL with indirect immunofluorescence with monoclonal antibodies.

Treatment is with high-flow oxygen and high-dose co-trimoxazole. Prednisolone should be added in severe cases (PO_2 <9.5 kPa).

C Prednisolone

Prednisolone may be appropriate in combination with co-trimoxazole in significantly hypoxic patients with *Pneumocysti*s, but should not be given alone (without appropriate anti-*Pneumocystis* cover).

A Intravenous co-amoxiclav and clarithromycin

This is appropriate treatment for community-acquired pneumonia.

B Anticoagulation with warfarin

This is appropriate treatment for PE. The pulmonary infiltrates seen here count much more towards an inflammatory process as the underlying cause of her symptoms.

D Intravenous ganciclovir

This is an appropriate treatment for CMV pneumonitis. CMV pneumonitis has a similar presentation, although other features such as small pulmonary nodules and predominantly basal consolidation/infiltration may be seen.

70101

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 64-year-old man is admitted to the Emergency Department with a cough productive of green sputum and a severe deterioration in cough and shortness of breath related to underlying chronic obstructive pulmonary disease (COPD). He normally takes tiotropium and high-dose Seretide® and, at best, his peak expiratory flow rate (PEFR) is 260. Today he is only able to manage 140. He continues to smoke 15 cigarettes per day.

On examination, his blood pressure (BP) is 152/72 mmHg and pulse 85 bpm and regular, and his temperature is 37.9 °C. There is poor air entry and wheeze throughout both lung fields, and an area of crackles at the right base. Oxygen saturation on pulse oximetry is 91%. He is allergic to penicillin.

Investigations:

Investigation	**Result**	**Normal values**
Haemoglobin (Hb)	138 g/l	135 - 175 g/l
White cell count (WCC)	14.1×10^9/l	$4.0 - 11.0 \times 10^9$/l
Platelets (PLT)	200×10^9/l	$150 - 400 \times 10^9$/l
Sodium (Na^+)	137 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	115 µmol/l	50 - 120 µmol/l
C-reactive protein (CRP)	245 mg/l	< 10 mg/l

He is treated with nebulisers and corticosteroids.

His chest x-ray is shown below.



Which of the following is the most appropriate antibiotic therapy?

Your answer was incorrect

A	Piperacillin with tazobactam (Tazocin®)
B	Co-amoxiclav
C	Cephalexin
D	Clarithromycin
E	Ciprofloxacin

Explanation

D	Clarithromycin

The answer is Clarithromycin (Option D).

This man has evidence of a community-acquired pneumonia on top of underlying COPD. His chest x-ray shows evidence of right upper lobe consolidation. As he is penicillin-allergic, a macrolide, such as clarithromycin, is the next most logical step. Where clarithromycin is prescribed against a background of theophylline, this should be discontinued during the course of antibiotics. Another option would be doxycycline – not because of his underlying COPD, but rather due to a diagnosis of community-acquired pneumonia with a background of penicillin allergy.

Image source: https://commons.wikimedia.org/wiki/File:X-ray_of_COPD_exacerbation_-_anteroposterior_view.jpg

C Cephalexin

Cefalexin (Option C) is incorrect.

There is 20% cross-over between penicillin and cephalosporin allergy, and cephalosporins are not as effective as clarithromycin in this situation.

A Piperacillin with tazobactam (Tazocin®)

Piperacillin with tazobactam (Tazocin®) (Option A) is incorrect.

Tazocin® contains a penicillin and would therefore be inappropriate. It is also not indicated for community-acquired pneumonia, unless there is evidence of *Pseudomonas*.

B Co-amoxiclav

Co-amoxiclav (Option B) is incorrect.

Co-amoxiclav is not appropriate because of the history of penicillin allergy reported here.

E Ciprofloxacin

Ciprofloxacin (Option E) is incorrect.

Ciprofloxacin is a potential therapy for COPD exacerbation but is not recommended for the treatment of pneumonia. It is also now recommended that quinolone treatment is avoided wherever an effective alternative is available due to a small but significant risk of debilitating adverse effects.

36522

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 55-year-old asthmatic patient presented in December with increasing cough and wheeze. She described a cough productive of brown sputum. She had no other past medical history. She was taking a steroid inhaler regularly and increasing doses of beta-agonist with no alleviation of her symptoms. She worked in a bakery for many years. She is an ex-smoker and stopped smoking 20 years ago. She has a dog at home.

On examination, she had a temperature of 38 °C. She was breathless at rest. She had reduced expansion anteriorly on the left side, with corresponding reduced breath sounds. She had a mild expiratory wheeze.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	141 g/l	135 - 175 g/l
White cell count (WCC)	12.7×10^9/l	$4.0 - 11.0 \times 10^9$/l
Neutrophils	7.37×10^9/l	$1.5 - 7.0 \times 10^9$/l
Lymphocytes	2.55×10^9/l	$1.5 - 4.0 \times 10^9$/l
Monocytes	0.6×10^9/l	$0.0 - 0.8 \times 10^9$/l
Basophils	0.05×10^9/l	$< 0.1 \times 10^9$/l
Mean corpuscular volume (MCV)	87.3 fl	80 - 100 fl
Platelets (PLT)	314×10^9/l	$150 - 400 \times 10^9$/l
Erythrocyte sedimentation rate (ESR)	28 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	16 mg/l	< 10 mg/l
Sodium (Na^+)	138 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5 - 5.0 mmol/l
Urea	4.8 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	95 μmol/l	50 - 120 μmol/l
Bilirubin	12 μmol/l	1 - 22 μmol/l
Alanine aminotransferase (ALT)	49 u/l	7 - 55 u/l
Alkaline phosphatase (ALP)	61 u/l	30 - 150 u/l
Albumin	36 g/l	35 - 55 g/l

Chest X-ray on presentation is shown below.



What does the chest X-ray show?

Your answer was correct

A Left upper lobe consolidation

B Left upper lobe collapse

C Left lower lobe collapse

D Left lower lobe consolidation

E Lingular consolidation

Explanation

B Left upper lobe collapse

The chest X-ray shows left upper lobe collapse with a veil like opacity over the left hemithorax. This patient has allergic bronchopulmonary aspergillosis (ABPA).

ABPA is suspected in any patient with asthma who has an abnormal chest X-ray and high peripheral blood eosinophilia (in this case, by adding together all the differentials the eosinophil count is >2). The chest X-ray may show diffuse pulmonary infiltrates, and pulmonary, lobar or segmental collapse occur as transient features. The most common cause is sensitivity to *Aspergillus fumigatus* spores.

The diagnostic criteria include:

- Asthma (in most cases)
- Peripheral blood and sputum eosinophilia
- Abnormal chest X-ray (infiltrates, segmental or lobar collapse)
- Positive skin tests/RAST to an extract of *A. fumigatus*
- *A. fumigatus* IgG serum-precipitating antibodies
- Raised total IgE >1000 ng/ml
- Fungal hyphae of *A. fumigatus* on microscopy of sputum.

The lung and eosinophils:

- Loeffler syndrome (eosinophils about 10% of blood WCC), also known as acute eosinophilic pneumonia/simple eosinophilic pneumonia. Mild self-limiting illness with transient migratory pulmonary shadows. Associated with parasitic infections, drug allergies and exposure to inorganic chemicals
- Tropical pulmonary eosinophilia (eosinophils >20%) – in tropical countries usually due to migrating larvae of the filarial worms *Wucheria bancrofti* and *Brugia malayi*
- Chronic/prolonged pulmonary eosinophilia (eosinophils >20%) – eosinophilic pneumonia persisting for more than one month. Chronic debilitating illness characterised by malaise, weight loss, fever and dyspnoea
- Allergic bronchopulmonary aspergillosis (eosinophils 5–20%) as in this case
- Churg–Strauss syndrome (eosinophils 5–20%) associated with asthma
- Hyper-eosinophilic syndrome (eosinophils >20%: eosinophilic infiltrations of various organs – e.g. lungs, heart, bone marrow. Can be associated with an eosinophilic arteritis
- External agents – drugs, toxins, parasitic infection.

Image source: https://radiopaedia.org/cases/left-upper-lobe-collapse-5?lang=gb

A Left upper lobe consolidation

There is no left upper lobe consolidation in the chest radiograph. The left lung appears more opaque due to left upper lobe collapse.

C Left lower lobe collapse

Left lower lobe collapse presents as ‘sail sign’, a triangular opacification visualised behind the cardiac shadow.

D Left lower lobe consolidation

There is no evidence of left lower lobe consolidation and the left hemi-diaphragm is clearly visualised.

E Lingular consolidation

The left cardiac border is clear, making linguar lobe consolidation unlikely.

70093

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 73-year-old man is sent to the Emergency Department by ambulance after a General Practitioner's (GP) home visit. The GP found him to be extremely short of breath, with a cough productive of rust-coloured sputum. He smokes ten cigarettes per day. On examination, he has right-sided crackles and bronchial breathing consistent with pneumonia. On attendance at the hospital, his respiratory rate is 24 breaths per minute, and he requires 3 litres of oxygen via nasal cannulae to achieve oxygen saturations of 96%. Chest X-ray confirms right basal consolidation and he is started on antibiotics for community-acquired pneumonia (CAP). He makes a good recovery and after five days of treatment, he is off oxygen and ready for discharge.

Which of these features of CAP indicates the need for a repeat chest X-ray at six weeks?

Your answer was incorrect

A Age > 50 years

B CURB-65 (confusion, urea, respiratory rate, blood pressure and age > 65) score of 2 on admission

C Need for hospital admission

D Need for oxygen therapy during admission

E Previous history of pneumonia

Explanation

A Age > 50 years

The British Thoracic Society (BTS) guidelines for community-acquired pneumonia (CAP) do not recommend repeat X-ray before discharge or at six weeks routinely for all patients with pneumonia, as there is no evidence on which to base this recommendation. Instead, they recommend that a chest X-ray should be arranged after about six weeks for all those patients who have a persistence of symptoms or physical signs or who are at higher risk of underlying malignancy (especially smokers and those aged over 50 years), whether or not they have been admitted to hospital.

C Need for hospital admission

The BTS guidelines indicate that all patients treated for pneumonia who have persistent signs or symptoms and/or are at higher risk of underlying malignancy (especially smokers and those aged over 50 years) should receive imaging six weeks after treatment, whether or not they are admitted to hospital.

B CURB-65 (confusion, urea, respiratory rate, blood pressure and age > 65) score of 2 on admission

Using the CURB-65 score is a useful way of stratifying a patient's risk associated with CAP. It should not be used to guide the need for repeat imaging.

D Need for oxygen therapy during admission

The need for oxygen therapy during admission does not indicate a need for a follow-up chest X-ray at six weeks.

E Previous history of pneumonia

A previous history of pneumonia does not indicate the need for repeat imaging in itself, although persistent symptoms or signs following initial treatment should indicate repeat imaging.

72984

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 17-year-old student presents with a persistent cough productive of brown/green sputum. He occasionally produces a small amount of blood. His mother said he had always suffered from chest infections, requiring numerous courses of antibiotics. He also complained of offensive-smelling stools which were difficult to flush .

On examination, he looks small for his age. Auscultation of his chest revealed bi-basal coarse crackles.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	112 g/l	135–175 g/l
White cell count (WCC)	10.7 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	511 × 10^9/l	150–400 × 10^9/l
Mean corpuscular volume (MCV)	91 fl	80–100 fl
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	3.9 mmol/l	3.5–5.0 mmol/l
Urea	5.3 mmol/l	2.5–6.5 mmol/l
Creatinine (Cr)	99 μmol/l	50–120 μmol/l
Bilirubin	18 μmol/l	1–22 μmol/l
Aspartate aminotransferase (AST)	34 u/l	10–40 u/l
Alkaline phosphatase (ALP)	93 u/l	30–150 u/l
Albumin	39 g/l	35–55 g/dl
Total protein level	50 g/l	60–83 g/l

His chest X-ray is shown below.



How would you make a diagnosis?

Your answer was correct

A	Sweat test
B	Measure immunoglobulins
C	High-resolution CT scan
D	Electron microscopy of cilia
E	Full lung function tests

Explanation

B Measure immunoglobulins

This patient has X-linked hypogammaglobulinaemia. The main clue is the calculated globulin fraction, which is low, if you minus the albumin from the total protein.

Patients usually present in childhood with recurrent infections, which result in bronchiectasis if untreated, and malabsorption, in a similar way to patients with cystic fibrosis. All immunoglobulin classes and B cells and plasma cells are reduced.
The defect is in the differentiation of pre-B cells into B cells. T cells are normal. The gene defect is on the long arm of the X chromosome, making it an X-linked condition.

Treatment is with intravenous (iv) immunoglobulin therapy which, if started early, can prevent progression of disease. The chest X-ray shows bronchiectasis – parallel thickened bronchial walls with a 'tram track' appearance.

A Sweat test

This is a diagnostic test for cystic fibrosis. This can present in infancy with failure to thrive, typically due to pancreatic insufficiency. Cough, recurrent chest infections, malabsorption and recurrent sinusitis are typical features. However, the low globulins in this patient's case make X-linked hypogammaglobulinaemia a more likely diagnosis here.

C High-resolution CT scan

This would help diagnose bronchiectasis but would not reveal the underlying cause in this case.

D Electron microscopy of cilia

This is a diagnostic test for primary ciliary dyskinesia, associated with chronic/recurrent respiratory infections and sinusitis. Up to 50% of patients with PCD have situs inversus. Kartagener syndrome is a subset of this condition, associated with bronchiectasis, sinusitis and dextrocardia.

E Full lung function tests

Lung function tests would help assess the severity of lung function impairment but would not be diagnostic of hypogammaglbulinaemia.

70148

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 63-year-old man is admitted with shortness of breath and productive cough. He is pyrexial at 39 °C, has a blood pressure of 81/59 mmHg, and is tachycardic at 113 bpm. Concern is raised regarding sepsis, and he is treated with intravenous (IV) piperacillin/tazobactam.

Initially, he improves; however, you are asked to review him, as his oxygen saturations have dropped to 81% despite 15 litres of high-flow oxygen. On arterial blood gas, his $p_a(O_2)$ is 6.9 kPa. He has a background of diabetes and obstructive sleep apnoea, for which he has home continuous positive airway pressure (CPAP).

His chest X-ray is shown below.



What is the most appropriate next management step?

Your answer was incorrect

A Change antibiotics to IV co-amoxiclav and clarithromycin

B IV furosemide

C IV methylprednisolone

D Referral to Intensive Therapy Unit (ITU)

E Utilise the patient’s home CPAP

Explanation

D Referral to Intensive Therapy Unit (ITU)

This patient has **Acute respiratory distress syndrome** (ARDS), as indicated by diffuse bilateral coalescent opacities on the chest X-ray. Most ARDS patients require mechanical ventilation, and maintaining the cardiac output typically requires the use of a combination of inotrope, vasodilators and transfusion. ARDS is associated with a mortality as high as 75%, even with ITU involvement; therefore, early referral is essential.

C IV methylprednisolone

This is ARDS. In the past, this was occasionally managed with IV methylprednisolone; however, this was not proven to be associated with increased survival rates. These patients usually require mechanical ventilation and must be referred to ITU.

A Change antibiotics to IV co-amoxiclav and clarithromycin

This patient has ARDS as indicated by diffuse bilateral coalescent opacities on the chest X-ray. This is most commonly associated with sepsis, as in this scenario. ARDS may be managed by treating the underlying cause; therefore, a change in antibiotics may well prove beneficial. However, the main thrust of management is supportive and should be undertaken in ITU.

B IV furosemide

The X-ray demonstrates diffuse bilateral coalescent opacities, for which a differential diagnosis may be severe pulmonary oedema in the context of heart failure. However, given the history of sepsis, ARDS must be suspected and the patient should be managed in ITU.

E Utilise the patient’s home CPAP

The chest X-ray and history of sepsis are strongly suggestive of ARDS. While CPAP with oxygen may maintain oxygenation, recommended management is involvement of ITU as patients usually require mechanical ventilation.

71771

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Faculty of Intensive Care Medicine. 2018. Guidelines on the management of ARDS.
ficm.ac.uk/news-events-education/news/guidelines-management-ards
(https://www.ficm.ac.uk/news-events-education/news/guidelines-management-ards)

Acute Respiratory Distress Syndrome

Expanded
explanations

A 35-year-old patient presents with a productive cough and shortness of breath on exertion over the past month. He has no personal past medical history, but his father suffered hepatocellular carcinoma at a young age. He is a smoker of 10 pack-years. On examination he has a hyper-expanded chest and bi-basal wheeze on auscultation.

He has spirometry showing the following results:

Investigation	**Result**
FEV_1	54% predicted
FVC	82% predicted
Residual volume (RV)	115% predicted
T_{LCO}	68% predicted
K_{CO}	67% predicted

Which of the following diagnoses would best fit the above results?

Your answer was incorrect

A Asthma

B Emphysema

C Lymphangitis carcinomatosis

D Thoracic cage deformity

E Slowly expanding pneumothorax

Explanation

B Emphysema

The spirometry shows an obstructive picture with raised RV, suggesting air trapping. He also has reduced alveolar function, suggesting a lung parenchymal cause. It is unlikely to be asthma given the short history and poor spirometry; it is more likely to be a rapidly progressive disease such as emphysema in a smoker with α1-antitrypsin. All patients under the age of 35 years presenting with COPD should have their α1-antitrypsin tested. His symptoms do not suggest someone with bad asthma. There is also a risk of HCC in patients with cirrhosis.

C Lymphangitis carcinomatosis

This is lymphangitis as a result of a malignancy, most commonly adenocarcinoma such as breast, lung, stomach or prostate. The patient’s young age makes this a less likely diagnosis than emphysema secondary to α1-antitrypsin deficiency. Also, this would be expected to show a restrictive pattern of PFTs, rather than the obstructive picture seen here.

A Asthma

The short history and poor spirometry make a rapidly progressive disease such as emphysema in a smoker with α1-antitrypsin deficiency more likely.

D Thoracic cage deformity

This would be expected to show a restrictive pattern of PFTs, rather than the obstructive picture seen here. The short history also makes this a less likely diagnosis.

E Slowly expanding pneumothorax

This would be expected to show a restrictive pattern of PFTs, rather than the obstructive picture seen here.

7162

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 76-year-old man with a 10-year history of chronic obstructive pulmonary disease was admitted to hospital with a 2-day history of progressively worsening dyspnoea and a cough productive of mucoid sputum. His wife stated that he is on home oxygen therapy for up to 12 h per day and home nebulisers. He is breathless on minimal exertion and rarely leaves the house. He has several exacerbations per year requiring hospital admission and these have become more frequent over the last few years, although he has never required ventilation before. He stopped smoking 2 years ago.

On examination he was acutely dyspnoeic with a respiratory rate of 32/min and was using accessory muscles of respiration. He was apyrexial, pulse 120/min and regular, blood pressure 182/92 mmHg. He was able to speak in broken sentences. Auscultation of his chest revealed generalised poor air entry and widespread expiratory wheeze.

His chest radiograph is shown below.



Arterial blood gases on 24% oxygen via Venturi mask reveals the following:

Investigation	Result	Normal values
pH	7.32	7.35–7.45
$p_a(O_2)$	6.2 kPa	10.5–13.5 kPa
$p_a(CO_2)$	7.2 kPa	4.6–6.0 kPa

His inspired oxygen concentration was increased to 28% and he was treated with intravenous hydrocortisone, salbutamol and ipratropium nebulisers and an aminophylline infusion. After 1 h he had made little improvement and was tiring.

Repeat blood gases on 28% oxygen reveals the following:

Investigation	Result	Normal values
pH	7.26	7.35–7.45
$p_a(O_2)$	6.8 kPa	10.5–13.5 kPa
$p_a(CO_2)$	9.2 kPa	4.6–6.0 kPa

What is the next most appropriate step in management?

Your answer was incorrect

A Immediate intubation and ventilation

B Intravenous doxapram

C Bi-level positive airway pressure (BiPAP)

D Increase inspired oxygen concentration to 35%

E Continuous positive airways pressure (CPAP)

Explanation

C Bi-level positive airway pressure (BiPAP)

This man is in type 2 respiratory failure due to a severe exacerbation of chronic obstructive pulmonary disease (COPD), and is not improving despite maximal medical therapy. Non-invasive ventilation is the best option in this situation and may be life saving. BiPAP has been shown to reduce respiratory acidosis, increase clearance of secretions and re-expand collapsed lung segments. However, it is only suitable for patients who are conscious, cooperative and able to protect their airway.

B Intravenous doxapram

Doxapram is a respiratory stimulant and should only be used when BiPAP is unavailable or in the case of reduced respiratory drive due to sedatives or anaesthetic agents. Doxapram also has a number of contraindications, which include hypertension (relative contraindication), such that concomitant therapy and past medical history should be carefully reviewed prior to starting therapy.

A Immediate intubation and ventilation

His poor premorbid functional status and lack of an acute precipitating event make him a poor candidate for the Intensive Therapy Unit. If he had a better function baseline this could be the ideal management choice, especially as he is tiring.

D Increase inspired oxygen concentration to 35%

This is a patient with type 2 respiratory failure and acidosis. His main problem is hypercapnia, which is decompensated as indicated by the low pH. He will need support in ventilation rather than oxygenation, and increasing inspired oxygen is likely to increase hypercapnia. Ventilatory support can help both hypoxia and hypercapnia, and further oxygen can be given during ventilation.

E Continuous positive airways pressure (CPAP)

With respect to BiPAP vs CPAP, outcome evidence is much more widely available for BiPAP, which drives the correct answer here.

9143

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 30-year-old man from London, with no significant past medical history, is on holiday in Peru to climb the Inca Trail. After climbing above 3500 m, he has developed a headache, associated with nausea and vomiting. He also feels tired and complains of insomnia. There is no past medical history of note and he is a non-smoker.

On examination, he has a Glasgow Coma Scale score of 14/15. His respiratory rate is 22 breaths per minute and his pulse is 90 bpm. His chest is clear on auscultation.

What treatment should be initiated en route to the hospital?

Your answer was incorrect

A Acetazolamide

B Betahistine

C Cyclizine

D Dexamethasone

E Nifedipine

Explanation

D Dexamethasone

This patient has symptoms of acute mountain sickness (AMS) and high-altitude cerebral oedema (HACE) with fatigue, headache, nausea and vomiting during a hiking expedition. The exact cause is unknown, but it is thought to be secondary to cerebral oedema and raised intracranial pressure. It is more common in people who live at low altitudes.

Symptoms and signs of AMS are non-specific; however, diagnosis can be made using the criteria according to the Lake Louise Score, namely, the setting of the recent gain in altitude, the presence of headache and at least one of the following symptoms:

- gastrointestinal – anorexia, nausea and vomiting
- fatigue and weakness
- dizziness or light-headedness
- difficulty sleeping.

The patient does not improve and progresses to developing ataxia and drowsiness. This is the development of HACE, which can be considered as end-stage AMS. Treatment is immediate descent, oxygen and intravenous dexamethasone.

C Cyclizine

Cyclizine is an antiemetic used in the treatment of nausea and vomiting. Although it may be helpful in acute illness, it plays no role in the treatment of AMS or HACE.

A Acetazolamide

Acetazolamide may be used for the prevention and treatment of AMS. It is a carbonic anhydrase inhibitor that causes intracellular acidosis. It has a diuretic effect which can be useful in AMS/HACE and high-altitude pulmonary oedema (HAPE). It is also beneficial as it shifts the oxygen dissociation curve to the right, and therefore, at lower partial pressures, more oxygen is released to the tissues.

B Betahistine

Betahistine is used in the treatment of vestibular disorders but plays no role in the management of this patient.

E Nifedipine

Nifedipine is used in the management of HAPE. Given the absence of a respiratory history and clear lung fields, HACE is the more likely diagnosis.

72947

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 60-year-old man with known emphysema presented to the Emergency Department with sudden onset of severe right-sided pleuritic chest pain and shortness of breath. He is a smoker of 20/day. On examination he looks unwell. He is pale and sweaty. BP 140/85 mmHg, pulse 120/min, respiratory rate 30/min, SaO_2 88% on air. He has reduced expansion and reduced breath sounds on the right side. Heart sounds were normal and abdominal examination unremarkable.

ABGs on air:

pH	7.41
PO_2	7.5 kPa
PCO_2	6.5 kPa

His CXR is shown below;



He is given 28% oxygen and analgesia and his saturations improve.

What would your immediate management of this patient be?

Your answer was incorrect

A Chest aspiration

B Insertion of an intercostal chest drain

C Arrange an urgent CT scan of the chest

D Refer to the thoracic surgeons

E Insertion of an intercostal tube and apply suction

Explanation

B Insertion of an intercostal chest drain

This man has a secondary pneumothorax as he has underlying lung disease. The management is different for primary (no underlying pulmonary disease) and secondary (underlying pulmonary disease) pneumothoraces. Current guidelines[1] suggest that an intercostal drain should be inserted if the patient is breathless, over 50 years old with a significant smoking history and there is a rim of air of > 2 cm from the chest wall on a standard size chest X-ray. This patient fits all the above criteria.

Suction should not be applied immediately and it is recommended that it should not be routinely employed. It maybe considered after 48 hours whilst awaiting surgical review if the lung is not fully re-expanded or if there is a persistent air leak. Suction should be low-pressure (-10 to -20 cmH_2O), high-volume.

If this patient had a persistent air leak, despite suction, they should be referred to the thoracic surgeons. The exact timing of referral remains contentious but it is suggested that in cases of persistent air leak or failure of the lung to re-expand, an early (3-5 days) thoracic surgical opinion should be sought.
For secondary pneumothoraces, an early thoracic surgical referral is recommended.

Image source: https://commons.wikimedia.org/wiki/File:PNTX.png

C Arrange an urgent CT scan of the chest

A CT is not required here, the diagnosis of pneumothorax is clearly visible on CXR.

A Chest aspiration

Chest aspiration should only be considered for a secondary pneumothorax <2cm if the patient is clinically stable.

D Refer to the thoracic surgeons

It would be appropriate to try and resolve pneumothorax with an intercostal drain before assessing whether a surgical referral is required.

E Insertion of an intercostal tube and apply suction

Suction should be applied with caution due to risk of re-expansion pulmonary oedema. It should only be applied after consideration and specialist advice and monitoring.

70121

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 45-year-old woman is referred to the Respiratory Clinic for review. She has a history of SLE which is currently treated with hydroxychloroquine, but she continues to have multiple joint pains and a photosensitive rash. Over the past few weeks she has suffered from increasingly severe left-sided pleuritic chest pain and a dull chest ache. On examination her BP is 155/85 mmHg, with pulse 75/min and regular. There are decreased breath sounds at the left base on auscultation.

Investigations:

Investigation	**Result**	**Normal values**
Haemoglobin (Hb)	121 g/l	135 - 175 g/l
White cell count (WCC)	10.5×10^9/l	$4.0 - 11.0 \times 10^9$/l
Platelets (PLT)	155×10^9/l	$150 - 400 \times 10^9$/l
Sodium (Na^+)	137 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.4 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	133 µmol/l	50 - 120 µmol/l
Erythrocyte sedimentation rate (ESR)	72 mm/h	1- 20 mm/h

Her chest x-ray is shown below



Which of the following is the most likely underlying cause of her chest symptoms?

Your answer was incorrect

A Alveolar haemorrhage

B Bronchiolitis obliterans

C Chronic interstitial pneumonitis

D Multiple pulmonary emboli

E Pleuritis

Explanation

E Pleuritis

Pleuritis is the most common manifestation of SLE-related lung disease; up to 60% of patients with SLE may report pleuritic chest pain, with or without concomitant pleural effusion. Pleural biopsies are rarely performed, although when they are, usually to exclude alternative diagnoses, non-specific inflammation is seen.

Image source: https://radiopaedia.org/cases/pleural-effusion-small?lang=gb (https://radiopaedia.org/cases/pleural-effusion-small?lang=gb)

C Chronic interstitial pneumonitis

Chronic interstitial pneumonitis (CIP) is associated with more widespread inflammatory infiltrates and may be seen in up to 15% of individuals with SLE.

A Alveolar haemorrhage

Alveolar haemorrhage may be seen in SLE, although more rarely than pleuritis, and it is associated with interstitial shadowing rather than the pleural effusion seen here.

B Bronchiolitis obliterans

Bronchiolitis obliterans isn't classically associated with a pleural effusion – it is associated with hyperinflation, attenuation of vascular markings and reticulonodular shadowing in some cases.

D Multiple pulmonary emboli

Although pulmonary embolism is a possible differential, multiple pulmonary emboli would be unlikely to affect one segment of the lung only.

36027

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 41-year-old woman presents with a dry cough, fatigue and multiple joint pains.

Her calcium is elevated at 2.85 mmol/l. She has multiple skin lesions, one of which is shown on her shin and a biopsy is taken.

What is the most likely histology for this lesion on her shin



Your answer was correct

A	Granuloma with prominent epithelioid cells with sparse lymphocytic infiltrate at the margins.
B	Necrotising granuloma with few immune cells and evidence of haemorrhage
C	Necrotic granulomas with a cuff of lymphocytes
D	Necrobiotic granuloma (altered collagen surrounded by histiocytes and some multinucleated giant cells)
E	Suppurative granuloma with central neutrophil infiltration

Explanation

A Granuloma with prominent epithelioid cells with sparse lymphocytic infiltrate at the margins.

This is typical of sarcoid granuloma, the most likely diagnosis.

B Necrotising granuloma with few immune cells and evidence of haemorrhage

This is the picture you would expect to see in Granulomatosis with polyangiitis.

C Necrotic granulomas with a cuff of lymphocytes

This is typical of the picture seen in tuberculoid granulomas, seen in TB and leprosy.

D Necrobiotic granuloma (altered collagen surrounded by histiocytes and some multinucleated giant cells)

This is the picture seen in rheumatoid nodules and necrobiosis lipoidica.

E Suppurative granuloma with central neutrophil infiltration

Suppurative granulomas with central neutrophil inflammation are associated with chronic infection due to diseases such as blastomycosis and cat scratch disease.

39098

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Sarcoidosis

Expanded explanations

A 67-year-old man presents with increasing shortness of breath, cough and pleuritic chest pain. He has a past history of myocardial infarction and left ventricular failure and a 40 pack-year smoking history. Medication includes ramipril, furosemide and amlodipine. Last year he was prescribed amiodarone for paroxysmal ventricular tachycardia and he has taken the medication since then. On examination his blood pressure is 142/85 mmHg, pulse is 85/min and there are scattered crackles on auscultation with evidence of a left-sided pleural effusion. Chest X-ray (CXR) is shown below.



Which of the following diagnoses fits best with this clinical picture?

Your answer was incorrect

A Left ventricular failure

B Amiodarone toxicity

C Angiotensin-converting enzyme (ACE) inhibitor-induced lung fibrosis

D Idiopathic pulmonary fibrosis

E Chronic obstructive pulmonary disease (COPD)

Explanation

B Amiodarone toxicity

Amiodarone toxicity occurs in approximately 5% of patients. This can vary from acute respiratory distress to a picture of cough, pleuritic chest pain, pulmonary infiltrates and small pleural effusions, to interstitial pneumonitis and eventually progressive pulmonary fibrosis. The chest x-ray here shows several areas of reticular opacities, most noticeable in the right upper zone. Due to its long half-life, amiodarone toxicity takes some weeks to resolve; as such, corticosteroids are added when therapy is discontinued as this may lead to more rapid resolution of symptoms. Of course cardiology input is vital before amiodarone is discontinued – as in this case when paroxysmal VT is the indication for therapy, an implantable cardioverter defibrillator is the next logical step (and has been proven in trials to have a superior outcome).

Image source: https://radiopaedia.org/cases/amiodarone-lung-1 (https://radiopaedia.org/cases/amiodarone-lung-1)

C Angiotensin-converting enzyme (ACE) inhibitor-induced lung fibrosis

ACE inhibitors are associated with a dry cough and bronchospasm, but are not associated with pulmonary fibrosis.

A Left ventricular failure

Left ventricular failure is associated with infiltrates on the CXR consistent with fluid overload, but not with the focal opacities seen here. Upper lobe diversion is also expected on CXR examination where there is significant LVF.

D Idiopathic pulmonary fibrosis

Idiopathic pulmonary fibrosis is diagnosed once other causes of pulmonary fibrosis have been excluded. In this case, given the amiodarone use, this is a more likely cause of the lung changes.

E Chronic obstructive pulmonary disease (COPD)

COPD is not associated with the pulmonary fibrotic changes seen on CXR here, although smoking-related lung disease may be a contributor to his shortness of breath.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 64-year-old woman is referred with worsening COPD. She has a 10 pack-year smoking history. Her exercise tolerance has been steadily decreasing due to dyspnoea, and she is only able to walk a short distance on the flat now. She has previously responded to oral prednisolone with around a 25% improvement in her FEV1. FEV1 is now only 38% of predicted. She has tried PRN salbutamol but this has not helped.

Which of the following represents the most appropriate next step for this patient?

Your answer was correct

A Switch salbutamol to PRN ipratropium

B Start BD inhaled beclomethasone

C Start oral prednisolone 5 mg OD

D Start oral theophylline

E Start regular high-dose inhaled fluticasone and inhaled long-acting β-agonist

Explanation

E Start regular high-dose inhaled fluticasone and inhaled long-acting β-agonist

This patient has severe COPD. Recent studies have demonstrated that high-dose combination therapy with fluticasone and long-acting β-agonist may slow deterioration in FEV1. Low-dose fluticasone, beclomethasone, ipratropium and ventolin have no impact on the deterioration of lung function. The TORCH study (2007) showed a reduction in exacerbations with respect to dual LABA and inhaled steroid therapy. It did, however, show that the risk of pneumonia was increased in the treated group. Current NICE guidance states that for patients with FEV_1 <50%, LABA and LAMA as combination therapy is the intervention of choice, except as here when there is evidence of previous reversibility when patients have been treated with corticosteroids in the past.

A Switch salbutamol to PRN ipratropium

NICE guidelines state that a once-daily long-acting muscarinic antagonist should be offered in preference to four-times-daily short-acting muscarinic antagonists to people with stable COPD who remain breathless or have exacerbations despite using short-acting bronchodilators as required. Ipratropium is a short-acting muscarinic antagonist and is therefore not a suitable option here.

B Start BD inhaled beclomethasone

NICE guidelines state that if FEV_1 is <50% predicted then give either LABA with an inhaled corticosteroid (ICS) in a combination inhaler, or LAMA. Corticosteroid monotherapy for COPD is not recommended.

C Start oral prednisolone 5 mg OD

Prednisolone would be appropriate if the patient had presented acutely with an exacerbation of COPD (though a higher dose would be required), but NICE guidelines state that ‘Maintenance use of oral corticosteroid therapy in COPD is not normally recommended’.

D Start oral theophylline

NICE guidance states that ‘Theophylline should only be used after a trial of short-acting bronchodilators and long-acting bronchodilators’ and therefore does not represent the most appropriate next step for this patient.

6609

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

NICE CKS. COPD
cks.nice.org.uk/chronic-obstructive-pulmonary-disease
(https://cks.nice.org.uk/chronic-obstructive-pulmonary-disease)

TORCH Study. The New England Journal of Medicine
nejm.org/doi/full/10.1056/NEJMoa063070
(https://www.nejm.org/doi/full/10.1056/NEJMoa063070)

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 72-year-old man presents on the Emergency Medical Take with a worsening cough and bloodstained sputum. He has stable chronic obstructive pulmonary disease (COPD) treated with a high-dose fluticasone and salmeterol inhaler. On examination, his BP is 135/85 mmHg and his heart rate is 80 bpm and regular. There are scattered coarse crackles and wheeze on auscultation. His CXR shows a left hilar mass, suspicious of malignancy, with a small left pleural effusion.

Which of the following is the most appropriate next step?

Your answer was incorrect

A Blind pleural biopsy

B Bronchoscopy

C Contrast-enhanced CT

D Sputum cytology

E Thoracoscopy

Explanation

C Contrast-enhanced CT

In this situation, with a central lesion suspicious of bronchial carcinoma, thoracoabdominal CT, including the liver and adrenals, is the most appropriate next step. This is carried out before bronchoscopy because it can provide valuable information about the location of the lung malignancy and the nature of the pleural effusion.

D Sputum cytology

Sputum cytology is not of value in the majority of patients, as the sensitivity is low. It should only be considered when patients refuse or are unable to undergo bronchoscopy.

A Blind pleural biopsy

Blind pleural biopsy is not recommended because of the risk that an abnormal area of pleura may be missed and unintended trauma caused, compared to an ultrasound-guided attempt.

B Bronchoscopy

Bronchoscopy is recommended after CT has been conducted to define the anatomy of an underlying lesion to ensure any abnormal areas are not missed.

E Thoracoscopy

Thoracoscopy may be required for investigation and biopsy of pleural tissue. However, CT is more useful as an initial step to define the nature of the effusion and has the benefit of being non-invasive.

40252

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Lung cancer: diagnosis and management
nice.org.uk/guidance/cg121/
(http://www.nice.org.uk/guidance/cg121/)

Lung Cancer

Media

Expanded explanations

Flashcards

A 20-year-old man has recently started work in the bakery section of his local supermarket. He complains of increasing shortness of breath, cough and wheeze during the course of the working week. These symptoms resolve somewhat at weekends and improved markedly during a week's holiday in Spain. Examination in the clinic on a Monday morning reveals no evidence of wheeze; peak flow is 520 (540 predicted).

Which of the following initial actions is most appropriate in this case?

Your answer was incorrect

A Trial of salbutamol

B 2-hourly peak flow recordings during the week, comparing work to rest periods

C Chest X-ray

D Bronchial provocation testing

E Patch testing

Explanation

B 2-hourly peak flow recordings during the week, comparing work to rest periods

It is important to establish the relationship of deterioration in peak flow to work periods as there is a suspicion of occupational asthma here. Occupational asthma in bakers may occur due to flour dust or to other allergens used in making breads and cakes. Review of the worker's work pattern, allowing for removal of exposure to the potential allergen, is optimal therapy, of course with the cooperation of the employer.

D Bronchial provocation testing

Bronchial provocation testing should only be undertaken in specialised centres, with appropriate anaesthetist support, and only when there is doubt as to the underlying trigger.

A Trial of salbutamol

A trial of salbutamol is inappropriate given the likely history of occupational triggers for this patient's asthma. Removal of exposure to any potential allergen is much more important than starting asthma therapy.

C Chest X-ray

A chest x-ray is not useful when evaluating potential bronchoconstriction related to asthma.

E Patch testing

Patch testing is rarely useful in evaluating occupational asthma.

32435

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Occupational Asthma

Expanded explanations

A 65-year-old patient presents with increased shortness of breath (SOB) to the Respiratory Outpatient Clinic. This has been progressively worsening over the past 6 months. He is a farmer, a smoker and has two daughters with asthma. There was obvious wheeze and coarse end-inspiratory crackles on examination of the chest. A chest X-ray shows diffuse non-specific changes consistent with lung disease.

His chest x-ray is shown below.



Which would be the next most appropriate investigation?

Your answer was incorrect

A Peak expiratory flow rate (PEFR)

B Spirometry and reversibility

C Serum precipitins

D Bronchoalveolar lavage

E CT thorax

Explanation

B Spirometry and reversibility

This man either has asthma, chronic obstructive pulmonary disease or farmer’s lung. Spirometry and reversibility would be the investigation of choice. A restrictive defect would support a diagnosis of farmer’s lung; an obstructive defect with reversibility would support a diagnosis of asthma; respiratory obstruction without reversibility would support a diagnosis of COPD. Management of farmer’s lung involves avoidance of exposure to hay moulds. High-dose steroid therapy may be useful in initial resolution of symptoms.

The chest x-ray here shows mildly expanded lung fields with slightly flattened diaphragms favouring obstructive pathology.

Image source: https://commons.wikimedia.org/wiki/File:Chest_x-ray_-_posteroanterior_view.jpg

C Serum precipitins

Precipitating antibodies will show evidence of exposure to actinomycetes but may not necessarily indicate disease.

A Peak expiratory flow rate (PEFR)

PEFR would only provide hints as to a diagnosis of obstructive lung disease and wouldn’t differentiate between asthma and COPD.

D Bronchoalveolar lavage

This may indicate lymphocytosis in extrinsic allergic alveolitis, but is invasive and wouldn’t therefore be an appropriate next test.

E CT thorax

CT thorax is likely to show structural changes, but wouldn’t indicate the severity of obstructive or restrictive lung disease.

6604

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 56-year-old man presented to his General Practitioner (GP) with a 4-week history of feeling generally unwell. He described nasal congestion, with pain under his left eye, flu-like symptoms and lethargy. His GP made a diagnosis of sinusitis and gave him a course of antibiotics. He did not improve and developed breathlessness, cough and chest pain. He had previously been fit and his only past medical history was of an inguinal hernia repair 10 years previously. He worked in a shop. He smoked 10 cigarettes a day and drank about 10 units of alcohol a week. He was taking no regular medication.

On examination he looked unwell and pale. Observations: respiratory rate 34/min, BP 140/85 mmHg, pulse 120 bpm, temperature 36.6 °C. He had crusting of his nasal septum. His JVP was not elevated, and heart sounds were normal. Auscultation of his chest revealed fine crackles bilaterally. His abdominal and neurological examinations were unremarkable.

Urinalysis:

Protein	++
Blood	++
Nitrites	-
Leucocytes	-
Bilirubin	-

Haematology:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	81 g/l	135 - 175 g/l
White cell count (WCC)	6.3 × 10^9/l	4.0 - 11.0 × 10^9/l
Mean corpuscular volume (MCV)	81 fl	80 - 100 fl
Platelets (PLT)	510 × 10^9/l	150 - 400 × 10^9/l
Erythrocyte sedimentation rate (ESR)	114 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	45 mg/l	< 10 mg/l
Sodium (Na^+)	142 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.9 mmol/l	3.5 - 5.0 mmol/l
Urea	21 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	341 µmol/l	50 - 120 µmol/l
Bilirubin	18 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	31 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	191 u/l	30 - 150 u/l

Albumin	34 g/l	35 - 55 g/l

Spirometry:

FEV1	72% predicted
FVC	77% predicted
KCO	122% predicted

His chest X-ray is shown below:



How would you treat this patient?

Your answer was correct

A Intravenous antibiotics

B Intravenous diuretics

C Cyclophosphamide and methylprednisolone

D Methylprednisolone and Methotrexate

E Plasma exchange

Explanation

C Cyclophosphamide and methylprednisolone

This is a question about pulmonary–renal syndromes (for a list see below).

The raised KCO and the CXR should make one think of pulmonary haemorrhage. This should narrow the diagnosis down to a vasculitis, Goodpasture syndrome and SLE.

This patient has granulomatosis with polyangiitis. The clinical diagnosis is made by the triad of upper respiratory tract (nasal polyps and sinusitis), lower respiratory tract and renal involvement. It is a small-vessel vasculitis associated with granulomas.

First-line immunosuppressive therapy is with cyclophosphamide and methylprednisolone.

Causes of pulmonary–renal syndromes:

- Systemic diseases:
  Granulomatosis with polyangiitis
  Microscopic polyangiitis
  Goodpasture syndrome
  Systemic lupus erythematosus
  Polyarteritis nodosa
  Henoch–Schönlein purpura
  Churg–Strauss syndrome (renal involvement less common)
  Polyarteritis nodosa (Rare)
- Primary pulmonary disease:
  Legionella pneumonia and interstitial nephritis
  Bacterial pneumonia with renal compromise secondary to sepsis
- Others:
  Pulmonary oedema with acute kidney disease
  Uraemic pneumonitis
  Right-sided bacterial endocarditis – may cause pulmonary embolic lesions and glomerulonephritis
  Iatrogenic glomerulonephritis with ciprofloxacin, e.g. in patients given ciprofloxacin for cystic fibrosis.

Image source: https://upload.wikimedia.org/wikipedia/commons/7/77/Wegeners_Granulomatosis_CXR.jpg (https://upload.wikimedia.org/wikipedia/commons/7/77/Wegeners_Granulomatosis_CXR.jpg)

A Intravenous antibiotics

This would be an appropriate treatment for severe pneumonia or sepsis.

B Intravenous diuretics

This would be appropriate for acute pulmonary oedema.

D Methylprednisolone and Methotrexate

This combination may be considered as initial treatment for non-organ or life-threatening granulomatosis with polyangiitis. This gentleman is clearly very unwell and has renal involvement.

E Plasma exchange

Most patients with granulomatosis with polyangiitis are treated initially with immunosuppression rather than plasma exchange. Patients with rapidly worsening renal function (> 500 μmol/l), despite immunosuppression, or marked pulmonary haemorrhage are usually referred earlier for plasma exchange.

70090

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Granulomatosis with Polyangiitis

Media

Expanded explanations

Flashcards

A 30-year-old patient with grade 3 Medical Research Council (MRC) dyspnoea had a transthoracic echocardiogram showing severe pulmonary hypertension, with normal left ventricle size and function and normal function of all his heart valves. His chest X-ray ordered by the GP showed clear lung fields and his lung function tests were within normal limits. On examination, his BP is 145/79 mmHg, his pulse is 80/min and regular, and his BMI is 32. His chest is clear on auscultation. There is bilateral pitting oedema of both legs and marked varicosities on the left leg. He tells you the legs have been swollen and painful intermittently over the past year.

Which of the following tests is the most appropriate first line investigation?

Your answer was correct

A High-resolution CT chest

B Ultrasound abdomen and venous ultrasound lower limbs

C Left heart catheterisation

D Right heart catheterisation

E Sleep study

Explanation

B Ultrasound abdomen and venous ultrasound lower limbs

It is important to determine whether pulmonary hypertension is primary or secondary, because any underlying cause of secondary pulmonary hypertension may be treatable. Primary pulmonary hypertension is associated with connective tissue disease, vasculitis, HIV infection and use of appetite suppressants (e.g. fenfluramine), but there are no clues in this question pointing towards any of these associated features.

Given that the question alerts us that the patient has bilateral leg oedema as well as left leg varicosities, this gives us a clue that this is likely secondary pulmonary hypertension, possibly due to chronic venous thromboembolism. Therefore, ultrasound scanning of the lower limb to confirm left leg deep vein thrombosis, and of the abdomen to look for any potential sources of inferior vena cava obstruction (*bilateral* leg oedema), would be prudent, as these are widely available, non-invasive investigations.

A High-resolution CT chest

The commonest cause of secondary pulmonary hypertension is COPD, but this patient has normal pulmonary function, so this is less likely. This response is a reasonable suggestion, but it is not necessarily the first-line investigation to be carried out, given that the patient has normal pulmonary function tests, making a structural abnormality of the lung parenchyma less likely.

C Left heart catheterisation

Left heart catheterisation is unlikely to give us any further diagnostic evidence. Causes of secondary pulmonary hypertension include structural problems with the heart, such as mitral valve disease and left-to-right shunt, but again, this is less likely as the transthoracic echocardiogram does not demonstrate any of this. We already know from the patient's transthoracic echocardiogram that his left heart has normal size and function, so there is nothing more to be gained from this invasive investigation.

D Right heart catheterisation

The invasive investigation of right heart catheterisation is going to add little more diagnostic utility. We already know that the patient has pulmonary hypertension from his transthoracic echocardiogram, and a right heart catherisation is only going to give more detail regarding pressures. It will not give us more information regarding the aetiology of the patient's pulmonary hypertension, which at this stage, we want to ascertain so that any underlying cause can be treated.

E Sleep study

This is a reasonable option, because we know that the patient is obese (BMI 32) and obstructive sleep apnoea is a common cause of secondary pulmonary hypertension. However, given his history of leg oedema and left leg varicosity and the relative availability of lower limb ultrasound scanning, it would be more prudent to rule out chronic venous thromboembolism first before moving on to sleep studies.

32380

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pulmonary Hypertension

Media

Expanded explanations

Flashcards

A 70-year-old woman presents in acute respiratory distress. She gives a history of some chest pain earlier that morning, which was worse on coughing. She is an ex-smoker of 30 pack-years (20/day for 30 years).

On examination, she looks unwell. Respiratory rate is 30/min, pulse 120/min, blood pressure (BP) 80/50 mmHg and O_2 saturation is 80% on air. Heart sounds are normal and her chest is clear.

ABG on air:

pH	7.44
PO_2	7.0 kPa
PCO_2	3.25 kPa
Base excess	-2.6
Bicarbonate	19.5 mmol/l

You manage to get an urgent CT within the hour, shown below.



The patient deteriorates.

What is the immediate management?

Your answer was correct

A IV diuretic

B IV antibiotics

C Streptokinase

D Alteplase

E Low-molecular weight heparin

Explanation

D Alteplase

This woman has had a massive pulmonary embolus. She presented with shortness of breath and tachypnoea. Her arterial blood gases show she has type I respiratory failure; note the low CO_2 – common in PE.

The CT shows large filling defects in the left and right main pulmonary arteries.

British Thoracic Society guidelines on the treatment of massive PE state that if a PE is sufficiently severe to cause circulatory collapse, thrombolysis should be given as early as possible. The current guidelines recommend alteplase as the thrombolytic agent of choice as it can be given to the hypotensive patient, as in this example.

A IV diuretic

Diuretics would be an appropriate treatment for acute pulmonary oedema.

B IV antibiotics

Intravenous antibiotics would be appropriate treatment if clinically she had severe pneumonia or sepsis.

C Streptokinase

The thrombolytic therapy of choice for PE with haemodynamic instability is alteplase, as streptokinase can worsen hypotension.

E Low-molecular weight heparin

Low-molecular weight heparin is used to treat patients with PE and no haemodynamic compromise while they are initiated on long-term anticoagulation, such as warfarin, to therapeutic levels. Unfractionated heparin can be used alongside thrombolysis in patients with PE and haemodynamic instability.

70081

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 69-year-old woman who is known to have severe COPD managed with high-dose fluticasone and salmeterol (combined) and tiotropium comes to the Emergency Department with an infectious exacerbation. She has had a cough productive of thick, yellow sputum for the past few days and now has a severe cough and shortness of breath. She is unable to mobilise more than a few metres from her chair at home.

On examination, she is pyrexial (38.2 °C), her pulse is 90/min and there is atrial fibrillation. Her blood pressure is 123/82 mmHg. There is bilateral wheeze and poor air entry on auscultation of her chest.

Investigations and arterial blood gas analysis (after antibiotics, one hour of back-to-back nebulisers and hydrocortisone IV), on 28% O_2:

Investigation	Result	Normal values
Haemoglobin (Hb)	134 g/l	135 - 175 g/l
White cell count (WCC)	8.9 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	204 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	138 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.8 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	139 μmol/l	50 - 120 μmol/l
Bicarbonate (HCO_{3-})	28 mmol/l	22 - 29 mmol/l
pH	7.29	7.35 - 7.45
pO_2	8.8 kPa	> 10.5 kPa
pCO_2	8.0 kPa	4.7 - 6.0 kPa

Which of the following is the most appropriate next step?

Your answer was correct

A Further nebulisers

B Intubation and ventilation

C Non-invasive positive pressure ventilation (NIPPV)

D IV doxopram

E IV Aminophylline

Explanation

C Non-invasive positive pressure ventilation (NIPPV)

Guidelines suggest that NIPPV should be considered in all patients with COPD who have respiratory acidosis despite maximal medical therapy. NIPPV should be instigated in a timely fashion, before there is further deterioration.

A Further nebulisers

Further nebulisers are unlikely to impact on air entry or CO_2 retention, only extending the time before NIPPV is instigated.

B Intubation and ventilation

In those with a pH of <7.26, there is evidence that intervention should be stepped up further, with ITU review to consider intubation and ventilation the next step.

D IV doxopram

IV doxopram is a respiratory stimulant which was the only realistic alternative option to intubation and ventilation before widespread use of NIV. It is inferior with respect to outcomes.

E IV Aminophylline

Aminophylline is no longer routinely used because of the risk of inducing atrial arrhythmias, with little added beta agonist impact.

36477

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

The Use of Non - Invasive Ventilation in the management of patients with chronic obstructive pulmonary disease
brit-thoracic.org.uk/document-library/clinical-information/niv/niv-guidelines/the-use-of-non-invasive-ventilation-in-the-management-of-pati...
(https://www.brit-thoracic.org.uk/document-library/clinical-information/niv/niv-guidelines/the-use-of-non-invasive-ventilation-in-the-management-of-patients-with-copd-admitted-to-hospital-with-acute-type-ii-respiratory-failure/)

Ventilatory Support

Expanded explanations

An 82-year-old man is referred to the Respiratory Clinic with a chronic cough. He has a 30-pack-year history of smoking.

His General Practitioner had referred him after an abnormal CXR, shown below:



Exposure to which industrial substance is most likely to give these findings?

Your answer was incorrect

A Asbestos

B Coal dust

C Silica

D Beryllium

E Iron dust

Explanation

B Coal dust

Coal dust exposure causes the occupational disease coal worker's pneumoconiosis (CWP). This can be classified into two subsets of disease. Simple CWP generally leads to multiple small, round opacities, typically 1–10 mm in diameter, mostly in the upper lobes. Complicated CWP shows much larger opacities in the upper lobes, with associated fibrosis, particularly when there is coexisting smoking.

The X-ray in this patient shows complicated CWP. Cases of CWP can be mistaken for malignancy, and further investigation following an initial X-ray is always required.

C Silica

Silica exposure is associated with diffuse small nodules scattered throughout the lung fields and with eggshell calcification of hilar lymph nodes. The patient presented with none of these making it unlikely here.

A Asbestos

Asbestos exposure is associated with pleural plaques and fibrosis, more marked in the lower zones rather than what this patient has.

D Beryllium

Beryllium exposure presents with acute pneumonitis that may continue to a granulomatous reaction (compared to sarcoidosis) with lymphadenopathy.

E Iron dust

Iron dust exposure presents with small nodules which appear denser than those associated with coal dust exposure, although there is less associated fibrosis.

7623

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 49-year-old woman presents to the Emergency Department complaining of haemoptysis. She has attended the GP on three previous occasions over the past 6 months complaining of recurrent nose bleeds and sinusitis.

Examination of the chest reveals scattered crackles and signs of consolidation, particularly on the left-hand side. On further questioning, she admits to smoking 30 cigarettes per day.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	110 g/l	135 - 175 g/l
White cell count (WCC)	8.1 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	430 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	140 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.7 mmol/l	3.5 - 5.0 mmol/l
Creatinine (Cr)	190 µmol/l	50 - 120 µmol/l
c-ANCA	Positive	
Chest X-ray	Consolidation with hilar lymphadenopathy and two well defined rounded opacities	

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Anti-GBM disease

B Eosinophilic granulomatosis with polyangiitis

C Sarcoidosis

D Tuberculosis

E Granulomatosis with polyangiitis (GPA)

Explanation

E Granulomatosis with polyangiitis (GPA)

The picture seen here fits well with a diagnosis of GPA, with positive c-ANCA, anaemia with leukocytosis, thrombocytosis, granulomas on CXR and evidence of renal involvement. Diagnosis is confirmed by taking a histology sample from affected tissue. Easiest to obtain is usually a biopsy of nasal/ upper respiratory tract mucosa, but if this is impossible, lung or renal biopsy is the alternative. Most physicians begin treatment with a course of 3 days' high-dose IV methylprednisolone and cyclophosphamide, with the aim of inducing remission from granulomatosis with polyangiitis. This is followed by conversion to lower doses of orals. IV immunoglobulin may be used in patients who are severely ill. More recently, IV rituximab (an anti-CD20 antibody) has shown promising results in inducing a clinical response in patients resistant to medical treatment.

C Sarcoidosis

Although sarcoid is a granulomatous disease, it is not associated with a positive ANCA.

A Anti-GBM disease

The are no granulomas on chest X-ray in anti-GBM disease. This most typically presents with pulmonary haemorrhage in the context of rapidly deteriorating renal function.

B Eosinophilic granulomatosis with polyangiitis

Eosinophilic granulomatosis with polyangiitis typically presents with asthma and peripheral blood eosinophilia. There is eosinophilic vasculitis. However, unlike this scenario, there is no granuloma formation in the lungs and it is associated more with p-ANCA positivity.

D Tuberculosis

Pulmonary tuberculosis may present with haemoptysis. However, C-ANCA would not be positive.

71101

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 73-year-old man with severe COPD managed with high-dose combination steroid and long-acting beta-agonist (LABA) inhaler is admitted to the Emergency Room. He has suffered increasing shortness of breath and a worsening cough over the past few days and is now completely unable to get his breath.

On examination he is pyrexial (38.2 °C) and there are chest signs consistent with both right-sided consolidation and pulmonary oedema. He looks tired and his respiratory rate has fallen to 13/min.

Investigations reveal the following:

Investigation	**Result**	**Normal value**
Haemoglobin (Hb)	131 g/l	135–175 g/l
White cell count (WCC)	10.1 × 10^9/l	4.0–11.0 x 10^9/l
Platelets (PLT)	295 × 10^9/l	150–400 x 10^9/l
Sodium (Na^+)	134 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	128 μmol/l	50–120 μmol/l
C-reactive protein (CRP)	52 mg/l	< 10 mg/l
Chest X-ray (CXR)	Right-sided consolidation, with evidence of pulmonary oedema	

Blood gases on controlled O_2 via a 40% venturi reveal the following:

Investigation	**Result**	**Normal values**
pH	7.28	7.35–7.45
pO_2	7.8 kPa	10.5–13.5 kPa
pCO_2	7.1 kPa	4.6–6.0 kPa

Which of the following is the most appropriate way to manage his O_2 replacement?

Your answer was incorrect

A Continue 40% O_2 by mask

B Decrease inspired pO_2 to 28% by mask

C Continuous positive airway pressure (CPAP)

D BiPAP

E Increase inspired O_2 to 60% by mask

Explanation

D BiPAP

This patient has evidence of both LVF and right-sided pneumonia, a long history of COPD and is beginning to tire. The acidosis seen here can drive worsening respiratory muscle weakness, and risks increasing CO_2 retention further. BiPAP can reverse acidosis and significantly reduces progression to intubation and ventilation.

C Continuous positive airway pressure (CPAP)

BiPAP reduces the work of breathing during inspiration more than CPAP, and hence confers a potential advantage in COPD versus CPAP alone.

A Continue 40% O_2 by mask

Acidosis and CO_2 retention is potentially progressive, and continuing the status quo here with respect to O_2 replacement is considered inadequate.

B Decrease inspired pO_2 to 28% by mask

This will drive hypoxia in addition to CO_2 retention, putting the patient at risk of cardiac rhythm disturbance and worsening pulmonary oedema.

E Increase inspired O_2 to 60% by mask

Once acidosis is established, respiratory muscle weakness is progressive; increasing inspired O_2 won't reverse this and will at best lead to a minor, short-term incremental improvement in oxygenation.

18669

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 70-year-old man with type 2 diabetes was admitted from a care home with a cough, fever and chest pain. He has been managed with IV co-amoxiclav for four days but is not improving. Other medication of note includes ramipril, atorvastatin, aspirin and mixed insulin BD.

On examination he is pyrexial at 38.5 °C, with BP 100/60 mmHg and pulse 95/min. Auscultation of the chest reveals coarse crackles at the right base consistent with continuing pneumonia.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	121 g/l	135–175 g/l
White cell count (WCC)	13.4 × 10^9/l	4–11 × 10^9/l
Platelets	231 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	4.9 mmol/l	3.5–5.0 mmol/l
Creatinine	145 μmol/l	50–120 μmol/l
Glucose	12.1 mmol/l	3.5–5.5 mmol/l
Sputum culture	MRSA sensitive to fusidic acid, gentamycin, vancomycin and rifampicin, resistant to co-amoxiclav	

Which of the following is the most appropriate treatment?

Your answer was incorrect

A Continue co-amoxiclav

B Add oral fusidic acid

C Switch to oral rifampicin

D Switch to IV vancomycin

E Add IV gentamicin

Explanation

D Switch to IV vancomycin

This man has nosocomial pneumonia, most likely acquired from his care home. The usual dose for vancomycin is 1 g BD 12-hourly, or 15 mg/kg. Vancomycin is effective as a single agent. It is only effective IV for this indication, due to poor absorption from the GI tract. It is implicated in nephrotoxicity and this patient already has a degree of renal impairment. Along with commencing vancomycin and careful monitoring during its use, other changes should be made to his medications and measures to prevent further decline in his renal function should be taken.

C Switch to oral rifampicin

Oral rifampicin is an option for MRSA, but it is associated with rapid development of resistance when used as a single agent.

A Continue co-amoxiclav

The patient has failed to improve on this agent and the sputum culture has proven resistance to it. Persisting with this would be detrimental to the patient and increase the risk of antibiotic resistance.

B Add oral fusidic acid

Widespread use of topical fusidic acid has encouraged the development of *S. aureus* resistance and therefore the use of fusidic acid alone as an anti-staphylococcal agent is not recommended.

E Add IV gentamicin

Although the sputum culture has shown that the MRSA in this case is sensitive to gentamicin, this patient already has a degree of kidney disease and therefore gentamicin is contraindicated. There would also be no sense in continuing the co-amoxiclav.

23313

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 43-year-old solicitor with breast carcinoma presents to the A&E department complaining of a cough and shortness of breath, associated with fever and sharp chest pain. She has only recently finished her last chemotherapy treatment course. She is currently on no treatment and has no allergies. She has no other significant past medical history.

On examination, she is dyspnoeic and tachypnoeic at rest and has a fever of 39°C. Pulse oximetry shows her oxygen saturations to be 90%. Examination of her chest shows a dull percussion note in the right upper region of the lung associated with increased vocal resonance and, on auscultation, bronchial breathing can be heard in the same area associated with coarse crackles. In order to confirm the suspected diagnosis, the patient undergoes a transbronchial biopsy. The specimen stains with Gomori methenamine silver and haematoxylin & eosin stains. Galactomannan testing is positive.

What is the most likely diagnosis?

Your answer was correct

A Secondary lung tumour

B Tuberculosis

C Invasive aspergillosis

D *Pneumocystis jirovecii* pneumonia

E Sarcoidosis

Explanation

C Invasive aspergillosis

Aspergillus fumigatus is a fungus transmitted to humans via inhalation. Hence, it primarily affects the lungs, causing five main syndromes: asthma, allergic bronchopulmonary aspergillosis (ABPA), aspergilloma, invasive aspergillosis and extrinsic allergic alveolitis (EAA). Asthma is a type-I hypersensitivity reaction to fungal spores whereas ABPA is a type-III hypersensitivity reaction to *A. fumigatus*. An aspergilloma is a fungal ball within a pre-existing lung cavity that may have resulted from tuberculosis, sarcoidosis or another necrotising pulmonary process. It is usually asymptomatic but may cause cough, torrential haemoptysis and fever. Invasive aspergillosis typically manifests with fever, cough, dyspnoea and pleuritic chest pain in patients with prolonged neutropenia or immunosuppression, such as that which occurs after chemotherapy. The CXR in invasive aspergillosis shows consolidation, and the patient has rapidly progressing and worsening hypoxaemia. This CXR appearance progresses despite the use of broad-spectrum antibiotics. The definitive diagnosis of invasive aspergillosis depends upon the demonstration of the organism in tissue. Haematoxylin and eosin (H&E) stain does not stain most fungi, except *Aspergillus* species and the zygomycetes. In *Pneumocystis jirovecii* pneumonia the typical chest X-ray appearance shows a diffuse central alveolar or interstitial infiltrate, although it may appear normal in early disease.

A Secondary lung tumour

GMS is used to stain fungi and *Pneumocystis jirovecii*, and would not be expected to stain a tumour.

B Tuberculosis

We are not given any history of contact with TB, or a history of symptoms such as weight loss or night sweats. GMS is used to stain fungi and *Pneumocystis jirovecii*.

D *Pneumocystis jirovecii* pneumonia

While this does stain with GMS, PJ typically occurs as opportunistic infection in patients with HIV. The positive galactomannan testing also directs us towards aspergillus as the diagnosis.

E Sarcoidosis

There is no mention of symptoms such as erythema nodosum or bilateral hilar lymphadenopathy, which is commonly seen on CXR, nor of non-caseating granulomas on biopsy, which are the hallmark of the disease.

6803

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Aspergillosis

Expanded explanations

A 28-year-old man from Botswana presents suffering from increased shortness of breath for the last three months. He has recently returned to the United Kingdom after a visit to his family. He has a dry, non-productive cough and admits to night sweats over the last two weeks. He is a non-smoker.

On examination, he is thin and not cyanosed at rest. You note multiple enlarged lymph nodes in the axillae and both groins. There is no finger clubbing. There are no signs on auscultation of the chest, and the chest X-ray is shown below.



Oxygen saturations reveal the following:

Investigation	Result
$S_p(O_2)$ at rest	98%

$S_p(O_2)$ on exertion	82%

What investigation is most likely to provide the diagnosis?

Your answer was incorrect

A Pulmonary function tests

B Ventilation/perfusion (V/Q) scan

C Bronchoalveolar lavage and transbronchial biopsy

D Computed tomography (CT) thorax

E Exercise electrocardiogram (ECG)

Explanation

C Bronchoalveolar lavage and transbronchial biopsy

The presence of exercise-induced oxygen desaturation in a young person should always raise the possibility of *Pneumocystis jiroveci* pneumonia (PCP), particularly in an individual who has lived in an area where undiagnosed human immunodeficiency virus (HIV) is common. *Pneumocystis jiroveci* is traditionally considered a protozoan, although recent data suggest that it may be taxonomically closer to a fungus. It is a ubiquitous pathogen, and disease most likely occurs as a result of latent infection acquired via the respiratory route early in life. Physical examination may reveal fever and tachypnoea, but examination of the chest is frequently normal. Radiographic features vary considerably, with diffuse, bilateral interstitial or alveolar infiltrates characteristic, but with unilateral or focal infiltrates and lobar or segmental consolidation also occurring. In this case, the X-ray shows diffuse bilateral interstitial changes.

Laboratory findings in PCP are generally unhelpful. Diagnosis can be made by staining spontaneous or induced sputum samples. In patients with HIV, bronchoalveolar lavage diagnostic yield approaches 95%, and this can be raised to nearer 100% with the use of transbronchial biopsy.

By -- Samir, 14 January 2007 (UTC) (http://en.wikipedia.org/wiki/Image:PCP_CAP_CXR.JPG) [GFDL (http://www.gnu.org/copyleft/fdl.html) or CC-BY-SA-3.0 (http://creativecommons.org/licenses/by-sa/3.0/)], via Wikimedia Commons

D Computed tomography (CT) thorax

CT thorax will confirm areas of infiltration consistent with PCP, although it will not deliver a microbiological diagnosis.

A Pulmonary function tests

Pulmonary function tests are likely to reveal decreased transfer factor, consistent with pulmonary infiltration, and potentially a restrictive defect, although they will not inform as to the underlying diagnosis.

B Ventilation/perfusion (V/Q) scan

V/Q scanning may show multiple ventilation scan defects due to pulmonary infiltration, and there is also likely to be evidence of pulmonary vessel shunting.

E Exercise electrocardiogram (ECG)

The hypoxia seen here is not due to cardiac decompensation, and the patient is highly unlikely to be able to complete the test because of the underlying *Pneumocystis* infection.

9836

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 54-year-old patient presents to the General Medical Clinic. He has a 6-month history of slowly worsening breathlessness and is finding it more difficult to carry out daily activities. He denies any chest pain but has an occasional cough, during which he brings up some sputum.

He is an ex-smoker with a 15 pack-year history. He has a minimal alcohol history. He travelled to the USAs last year. He has not been in contact with anyone with known tuberculosis. He has a long history of working as an electrician since he was 18 years old, but has lately struggled with his work due to breathlessness.

On examination, he has reduced air entry on the right lower lobe. There is also a slight general wheeze.

His hands are pictured below.



A chest X-ray demonstrates a small-to-moderate right-sided pleural effusion and evidence of multiple pleural plaques.

Which investigation is most likely to differentiate between benign and malignant disease?

Your answer was incorrect

A	Pleural biopsy
B	Pleural aspiration for cytology
C	Computed tomography (CT) chest with contrast
D	Liver function tests

E Blood film

Explanation

A Pleural biopsy

This patient has a long history of a profession associated with asbestos exposure and has developed breathlessness. In addition, there are pleural plaques which are highly suggestive of asbestos-related disease. The presence of a pleural effusion and clubbing raises the possibility of malignant mesothelioma.

C Computed tomography (CT) chest with contrast

A CT will be helpful to delineate any masses, but it will still struggle to differentiate between a malignant and a benign lesion unless there is clear evidence of metastatic or invasive disease. This is unusual with mesotheliomas, and therefore a biopsy would be more appropriate. In addition, the patient is relatively young and should be able to tolerate a biopsy.

B Pleural aspiration for cytology

Pleural aspiration can be done for this patient as he is symptomatic with a pleural effusion. However, the cytology from aspiration has a high rate of false-negative results. Therefore, it is important to send a large volume to be centrifuged to allow for a better yield. Because of this, the most definitive diagnosis is a biopsy which has a much lower rate of false-negative results.

D Liver function tests

Liver function tests can be useful and if severely deranged can indicate liver metastases; however, this is unusual for mesotheliomas and highly non-specific.

E Blood film

A blood film would be unlikely to give any abnormal results. It would be far more useful if acute haematology malignancy was suspected, such as in the presence of lymph nodes or night sweats.

71597

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Mesothelioma and Other Asbestos Related Conditions

Media

Expanded explanations

Flashcards

A 39-year-old lady from Zimbabwe presents to the Emergency Department with a 5-day history of increasing shortness of breath and dry cough. She has a 1-day history of sudden onset of sharp chest pain which is worse on coughing, movement and deep inspiration. She describes having had night sweats and a poor appetite for 2 months. Her weight has decreased by about 5 kg. She has a past medical history of genital herpes and depression. She is married and has two children who live in Zimbabwe. She moved to England 2 years ago and has not travelled since. She is a smoker of 20/day and drinks little alcohol.

On examination she looked unwell. Observations: temperature 38.3°C, BP 108/72 mmHg, pulse 120/min, regular, respiratory rate 30/min. Auscultation of her chest revealed inspiratory crackles bibasally.

Investigations:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	102 g/l	135 - 175 g/l
White cell count (WCC)	9.4 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	8.6 × 10^9/l	1.0 - 7.0 × 10^9/l
Lymphocytes	0.6 × 10^9/l	1.5 - 4.5 × 10^9/l
Platelets (PLT)	439 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	79 fl	80 - 100 fl
C-reactive protein (CRP)	73 mg/l	< 10 mg/l
Sodium (Na^+)	141 mmol/l	135 - 145 mmol/l
Potassium (K^+)	4.2 mmol/l	3.5 - 5.0 mmol/l
Urea	5.0 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	75 µmol/l	50 - 120 µmol/l
Glucose	5.7 mmol/l	3.9 - 7.1 mmol/l

Blood gases on 80% oxygen	**Result**
pH	7.46
PCO_2	3.44 kPa
PO_2	10.77 kPa
Bicarbonate	17.6 mmol/l
Base excess	-3.5

Her chest X-ray is shown:



What is the most likely diagnosis?

Your answer was incorrect

A *Legionella* pneumonia

B *Pneumocystis jirovecii* pneumonia

C Pulmonary embolus

D Nocardiosis

E CMV pneumonitis

Explanation

B *Pneumocystis jirovecii* pneumonia

For the purpose of the exam, people who have lived abroad (especially Africa and South America), businessmen who work abroad, men who have sex with men (MSM) and intravenous drug users are more likely to have HIV.

This lady has HIV – the history of living in Zimbabwe and the low lymphocyte count are the clues. *Pneumocystis jirovecii* is the most common opportunistic infection to cause pneumonia in AIDS – especially when the CD4 count is <200/mm^3. It accounts for about 50% of cases of pneumonia in AIDS and 40% of all AIDS-defining illnesses.

Patients usually present with a fever, dry cough and breathlessness. They are usually hypoxic and desaturate on exercise.

In PJP the chest X-ray usually shows bilateral interstitial shadowing and cysts in the mid and lower zones. However, the CXR may be normal. Pneumothorax (because the cysts rupture) may be present in up to 10%.

Diagnosis is now usually made by PCR of sputum or BAL fluid. Other diagnostic methods include staining induced sputum or BAL with indirect immunofluorescence with monoclonal antibodies.

Treatment is with high-flow oxygen and high-dose co-trimoxazole. Prednisolone should be added in severe cases ($PO_2 < 9.5$ kPa).

Other respiratory diseases associated with immunosuppression, in particular with HIV and AIDS, include:

- Tuberculosis
- Other bacterial pneumonias, including *Mycobacterium avium-intracellulare*
- *Cryptococcus* (other fungi include – *Candida*, *Histoplasma*, *Nocardia* and *Aspergillus fumigatus*
- CMV pneumonitis (other viruses include herpes simplex, Epstein–Barr virus)
- Toxoplasmosis
- Kaposi's sarcoma
- Lymphoma

C Pulmonary embolus

The pleuritic chest pain in this case is most likely due to pleurisy, although clinicians must be aware that there is a risk of pneumothorax in *Pneumocystis* pneumonia.

A *Legionella* pneumonia

The history is clearly suggestive of underlying HIV (patient from Zimbabwe, 2-month history of being unwell and night sweats, weight loss). The chest radiograph shows widespread changes in keeping with *Pneumocystis* pneumonia. There is very little in the history or investigation results to suggest *Legionella* pneumonia.

D Nocardiosis

HIV-positive individuals are at risk of nocardiosis, but you would expect a productive cough and history of abscess formation (e.g. skin abscesses, neurological symptoms to suggest brain abscess or lung abscess and cavitation). Diffuse pulmonary shadowing can occur, but the overall history is more in keeping with *Pneumocystis*.

E CMV pneumonitis

HIV-positive individuals are at risk of CMV pneumonitis, and this is the main differential here. One might expect GI or visual symptoms suggestive of systemic CMV disease. The lack of these symptoms with the history of pleuritic chest pain makes *Pneumocystis* pneumonia more likely.

70098

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 55-year-old man presents to the Emergency Department with a dry cough and shortness of breath, which he has had for the past four days. He also complains of some abdominal discomfort, nausea, vomiting, diarrhoea and a headache. He was previously well before he developed these symptoms and is not on any medication, having recently returned from a convention in Spain.

On examination, he is unwell with a blood pressure of 123/67 mmHg, a pulse of 92/min and a respiratory rate of 22/min. On auscultation of his chest, he has bi-basal crackles. He is pyrexial, 38.9 °C. His cardiovascular examination is unremarkable.

Investigations reveal the following:

Investigation	Result	Normal Value
Haemoglobin (Hb)	143 g/l	135–175 g/l
White Cell Count (WCC)	16.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	278 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	129 mmol/l	135–145 mmol/l
Potassium (K^+)	3.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	112 µmol/l	50–120 µmol/l
Mean corpuscular volume (MCV)	89 fl	76–98 fl
Urea	10.0 mmol/l	2.5–6.5 mmol/l
C-reactive protein (CRP)	169 mg/l	< 0 mg/l
Urinalysis	protein ++, blood ++, glucose –	

What is the likely diagnosis?

Your answer was incorrect

A Streptococcal pneumoniae

B Mycoplasma pneumoniae

C Chlamydia pneumoniae

D Legionella pneumophila

E Viral pneumonia

Explanation

D Legionella pneumophila

Legionella pneumophilia usually infects middle-aged or older men, is more severe in smokers and has an incubation period of 2–10 days. Male to female ratio is 2:1. It causes moderate leucocytosis (neutrophilia, lymphopenia), hyponatraemia, deranged LFTs, proteinuria, haematuria and myoglobinuria. Infection may be acquired from infected water-based air-conditioning systems, hence the link to a convention in Spain. Both fluoroquinolones and clarithromycin have activity against the bacterium. Rifampicin may be used as combination therapy in severe disease.

B Mycoplasma pneumoniae

Mycoplasma pneumonia tends to follow a more subacute course, with pleuritic chest pain and a dry cough both prominent features. Often appearance on chest X-ray is of a more severe pneumonia than first appears to be the case clinically.

A Streptococcal pneumoniae

Streptococcal pneumonia is less likely to be associated with hyponatraemia or GI upset. The attendance at a convention is also a stronger pointer towards Legionnaire's.

C Chlamydia pneumoniae

Chlamydial pneumonia, like mycoplasma, follows a more subacute course, and may be seen in patients who keep or are in close proximity to birds.

E Viral pneumonia

Viral pneumonitis is more likely to be associated with generalised crackles on auscultation of the chest due to more widespread infiltration, and a lesser rise in CRP would be expected.

7508

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 69-year-old man was taken to the Emergency Department with a reduced GCS score; no history was available from Accident & Emergency for the patient. The ambulance crew said that the call was put out by his wife as he had become increasingly short of breath. When they arrived he looked breathless and cyanosed. He denied any chest pain at that time.

On examination, he had a GCS score of 13, respiratory rate 10/min, BP 130/70 mmHg, pulse 120/min, temperature 36.9 °C, SaO_2 94% on 15 l oxygen. Auscultation of the chest revealed an expiratory wheeze throughout.

Heart sounds were normal and abdominal examination unremarkable. There was no obvious focal neurological abnormality and both plantars showed a flexor response.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	173 g/l	135 - 175 g/l
White cell count (WCC)	8.2 × 10^9/l	4.0 - 11.0 × 10^9/l
Platelets (PLT)	400 × 10^9/l	150 - 400 × 10^9/l
Mean corpuscular volume (MCV)	84 fl	80 - 100 fl
C-reactive protein (CRP)	21 mg/l	< 10 mg/l
ECG	Sinus tachycardia	
Chest X-ray	Hyperinflated lung fields with low, flattened diaphragm; no evidence of a pneumothorax	
Renal function & liver tests	Unremarkable	



ABGs on high-flow oxygen on this admission:	
pH	7.29
PCO_2	10.91 kPa
PO_2	12.39 kPa
Bicarbonate	40.6 mmol/l
Base excess	10.4

What is the most appropriate next step in the management of this patient?

Your answer was incorrect

A Intubation and invasive ventilation

B Start non-invasive ventilation

C Hydrocortisone 100 mg IV

D Salbutamol 2.5 mg via nebuliser

E Reduce inspired oxygen concentration to 28% and repeat ABG

Explanation

E Reduce inspired oxygen concentration to 28% and repeat ABG

This patient has an acute on chronic respiratory acidosis. The high bicarbonate and base excess shows that there is an underlying chronic respiratory acidosis with renal compensation. The acidosis and high CO_2 indicate an acute respiratory acidosis. If this was acute without a chronic element then, for a PCO_2 of 10.91 one would expect the pH to be lower.

This question is about oxygen therapy in COPD. The diagnosis of COPD is suggested from the CXR and flow-volume curve, which shows the classic shape for COPD. In some patients with COPD, CO2 retention is more prevalent. Two reasons are now thought to underlie this, 1) there is greater v/q mismatch in COPD, and secondly, the Haldane effect, a shift in the CO2 dissociation curve occurs.

The immediate management on arrival in the Emergency Department is to reduce the inspired oxygen concentration and repeat the arterial blood gases. If by doing this the patient becomes hypoxic or remains acidotic, they should then be referred for non-invasive ventilation.

National Collaborating Centre for Chronic Conditions. 2004. Chronic obstructive pulmonary disease. National clinical guideline on management of chronic obstructive pulmonary disease in adults in primary and secondary care.Thorax, 59 (suppl 1), 1-232

C Hydrocortisone 100 mg IV

While NICE recommend corticosteroids (prednisolone orally for a maximum of 7-14 days) for patients without contraindications who present to hospital with an exacerbation of COPD, in this case the patient has become obtunded due to excessive oxygen therapy. Corticosteroids may play a role in his medical management, but the optimal next step is to provide controlled oxygen therapy.

A Intubation and invasive ventilation

An anaesthetic opinion would be warranted for consideration of intubation if he did not respond to controlled oxygen ± a trial of NIV, but it does not represent the optimal next step for this patient.

B Start non-invasive ventilation

Non-invasive ventilation (NIV) would be appropriate if controlled oxygen therapy and medical therapies did not result in clinical improvement including resolution of acidosis, but does not represent the optimal next step for this patient.

D Salbutamol 2.5 mg via nebuliser

Salbutamol (and/or ipratropium) via nebuliser is recommended therapy in the treatment of COPD exacerbation, but the most pressing concern in this patient's case is his reduced GCS secondary to excessive oxygen therapy. The most appropriate next step is to provide controlled oxygen therapy and repeat an ABG.

70129

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

Chronic obstructive pulmonary disease
ncbi.nlm.nih.gov/pubmed/15041752
(http://www.ncbi.nlm.nih.gov/pubmed/15041752)

Ventilatory Support

Expanded explanations

A 70-year-old female smoker presents with a 3-day history of loss of sensation of both her lower legs. She also reports a 1-stone (6.53 kg) weight loss and unresolving cough over the last month. Her chest X-ray shows a large hilar mass, not seen on a chest X-ray from 2 months ago. She has no other past medical history and is not on any medication.

On examination, she has loss of light touch and pain sensation in a stocking distribution, reduced air entry in her right chest, palpable supraclavicular lymph node and a large craggy liver edge.

Investigations reveal the following:

Investigation	Result	Normal Value
White cell count (WCC)	6.2 × 10^9/l	4.0–11.0 × 10^9/l
Haemoglobin (Hb)	116 g/l	115–155 g/l
Platelets (PLT)	347 × 10^9/l	150–400 × 10^9/l

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Small-cell carcinoma

B Non-Hodgkin's lymphoma

C Adenocarcinoma

D Sarcoid

E Squamous cell carcinoma

Explanation

A Small-cell carcinoma

Given the rapid progression and spread of her pathology and the probable paraneoplastic peripheral neuropathy, this is the most likely diagnosis. Small cell carcinoma accounts for 20% of all lung cancers (National Cancer Intelligence Network, 2012), and it is the morphological subtype of lung cancer most closely associated with smoking. Small cell carcinoma often presents with unilateral hilar enlargement, a perihilar or a mediastinal mass. Less commonly, it may present with a central lesion, often cavitating, and given its rapid growth, with satellite lesions elsewhere in the thorax, characteristically with extensive, bulky mediastinal lymphadenopathy.

C Adenocarcinoma

Though the most common type of non-small-cell lung cancer, it is usually slow growing. It typically presents as a peripheral mass lesion, or less commonly as an area of segemental ground-glass attenuation.

B Non-Hodgkin’s lymphoma

The rapid spread of disease in this patient would be associated with an abnormal full blood count in non-Hodgkin’s lymphoma.

D Sarcoid

The history of smoking and rapid progression of pathology are suggestive of a malignant process. Sarcoid could cause the neurology seen, but would cause a smooth, enlarged granulomatous liver and widespread generalised lymphadenopathy rather than the craggy liver edge and isolated palpable supraclavicular lymph node noted.

E Squamous cell carcinoma

Like adenocarcinoma, squamous cell carcinoma is much slower growing. It typically presents as a central lesion on chest X-ray, with or without lung collapse due to intraluminal obstruction.

7175

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 56-year-old man is referred to the outpatients department with a 2-year history of increasing breathlessness and deteriorating exercise tolerance. He has a non-productive cough and always feels tired, so much so that he has needed to ask his wife to help him look after his pigeons, which he has been racing for many years. Over the last year he has lost about 1 stone in weight. He has a past medical history of asthma as a child. In his social history, he has had no history of recent foreign travel, except to France. He has always worked in a bank and is an ex-smoker who drinks minimal amounts of alcohol. On examination he is not clubbed. On auscultation of his chest, he has fine inspiratory crackles.

ABGs on air reveal:

pH	7.36
PO_2	7.4 kPa
PCO_2	4.7 kPa

Lung function tests:

FEV_1	70% predicted
FVC	64% predicted
KCO	53% predicted
TLCO	55% predicted

His CXR is shown below;



Which of the following would help make a diagnosis?

Your answer was incorrect

A *Aspergillus fumigatus* precipitins

B Avian serum precipitins

C Blood cultures

D Bronchoscopy

E Serum ACE

Explanation

B Avian serum precipitins

The patient has interstitial lung disease, demonstrated by the slowly progressive shortness of breath, type I hypoxia and restrictive lung function tests with reduced TLCO and KCO. The chest radiograph shows reticular nodular shadowing. The diagnosis is extrinsic allergic alveolitis secondary to pigeons – bird fancier's lung. This is caused by inhaled avian serum proteins (usually IgA) present in the pigeon's feathers and excreta. It is a hypersensitivity pneumonitis caused by a specific immunological response to inhaled organic dusts.

EAA may present either as an acute or a chronic disease. Acute disease develops several hours after exposure to high concentrations of the allergen. Breathlessness and flu-like symptoms occur sometimes with headaches, fever and myalgia. Symptoms usually resolve within 48 h. The patient has chronic disease as it has progressed to interstitial lung disease with fibrosis.

The diagnosis of extrinsic allergic alveolitis is made by:

- Identifying a potential source of antigen
- Characteristic clinical features: this patient has chronic EAA and presents with similar features to pulmonary fibrosis – clubbing tends to occur later (idiopathic pulmonary fibrosis would typically be associated with clubbing)
- Characteristic radiology (usually upper lobe fibrosis)
- Lung function tests – showing a restrictive defect with reduced carbon monoxide gas transfer
- Demonstrating precipitating antibodies (precipitins) in the patient's serum
- Resolution or improvement of symptoms following avoidance of exposure to the cause
- Histology of lung tissue biopsy showing a mononuclear cell infiltrate with the formation of non-caseating granulomas
- Fluid obtained from bronchoalveloar lavage has a high lymphocyte count (typically >40%).

Treatment is avoidance of the antigen. Corticosteroids may improve recovery in acute attacks but do not seem to provide long-term benefit. They are not usually helpful once fibrosis has occurred.

Other causes of EAA include:

- Farmer's lung, due to *Micropolyspora faeni* and *Thermactinomyces* in mouldy hay, straw and grain
- Bagassosis due to *Thermactinomyces sacchari* in sugar cane processing
- Malt worker's lung due to *Aspergillus clavatus*
- Mushroom worker's lung due to thermophilic actinomycetes
- Ventilation and water-related contamination, e.g. due to thermophilic actinomycetes contaminating air-conditioning systems
- Veterinary workers and animal handlers
- Workers in milling and construction, e.g. wood dust pneumonitis due to *Alternaria* spp.

Causes of upper-zone fibrosis:

- Progressive massive fibrosis/coal worker's pneumoconiosis
- Ankylosing spondylitis
- Sarcoid/silicosis
- Tuberculosis
- Extrinsic allergic alveolitis
- Radiation (depending on site).

C Blood cultures

There is nothing to suggest sepsis here.

A *Aspergillus fumigatus* precipitins

This is a test for *Aspergillus* lung disease.

D Bronchoscopy

Bronchoscopy may be helpful as part of the work-up here, but is not the test most likely to give the correct diagnosis.

E Serum ACE

This is not a specific test for any condition, but may be raised in sarcoidosis and TB.

70115

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 52-year-old man who is a known case of decompensated chronic liver disease presents to the Respiratory Clinic with gradual worsening of exertional dyspnoea for the past four weeks. He also reports easy fatiguability and worsening abdominal distension over the past few weeks despite being compliant with medications.

On examination, his blood pressure is 100/60 mmHg and his heart rate is 74 bpm. Pulse oximetry reveals an oxygen saturation (SpO_2) of 91% in the recumbent position and 84% in the upright position. The respiratory examination was normal and the abdominal examination was significant for the presence of ascites.

Which of the following is the most definitive management for this patient?

Your answer was incorrect

A Liver transplantation

B Long-term oxygen therapy

C Lung transplantation

D Spironolactone and furosemide

E Transjugular intrahepatic portosystemic shunt (TIPS)

Explanation

A Liver transplantation

Patients with hepatopulmonary syndrome (HPS) and a $p_a(O_2)$ of < 8 kPa should be evaluated for liver transplantation. A number of studies indicate complete or near-complete resolution of HPS, with improved oxygenation in the majority of patients within the initial 6–12 months.

C Lung transplantation

Lung transplantation is not a recommended treatment for patients with HPS.

B Long-term oxygen therapy

This patient, who is a known case of decompensated chronic liver disease now presenting with orthodeoxia and worsening exertional dyspnoea, can be diagnosed to have HPS. Although long-term oxygen therapy is the most frequently recommended treatment for HPS, it is not curative. It only improves the features related to intrapulmonary vascular shunts, eg dyspnoea, fatigue, desaturation.

D Spironolactone and furosemide

Although spironolactone and furosemide are the most frequently used diuretics for the management of decompensated chronic liver disease with ascites, they have not been proven to be beneficial in the management of HPS.

E Transjugular intrahepatic portosystemic shunt (TIPS)

TIPS has been found to be beneficial in reducing portal pressure. However, it is not recommended as a treatment strategy in patients with HPS. In addition, TIPS may worsen HPS by increasing the hyperkinetic state, leading to more pulmonary vasodilatation, thereby increasing the intrapulmonary shunting and worsening hypoxaemia.

72782

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 57-year-old man with a 40 pack per year smoking history presents to his General Practitioner with fatigue and lack of concentration, making it difficult for him to even attend work. His wife tells you that he is frequently sleeping in front of the television.

On examination, his heart rate is 88 bpm and his blood pressure is 136/90 mmHg. He has a normal cardiovascular examination and auscultation of the lung fields demonstrates mild expiratory wheeze, while oxygen saturation is 91%. His height measures 180 cm, weight 120 kg and neck circumference 44 cm.

Which of the following is the most appropriate treatment option?

Your answer was incorrect

A Intra-oral device to prevent snoring

B Long-term oxygen therapy

C Overnight continuous positive airway pressure (CPAP)

D Modafinil

E Nocturnal lorazepam

Explanation

C Overnight continuous positive airway pressure (CPAP)

Overnight continuous positive airway pressure (CPAP) is the treatment of choice as it has been shown in studies to be effective in reducing obstructive sleep apnoea (OSA) symptoms and is the first line in moderate to severe disease requiring intervention. However, it does not prevent cardiovascular mortality associated with OSA. Hypnotics worsen OSA, as do alcohol, smoking and obesity, and so patients should alter their lifestyle appropriately, with weight loss and alcohol and smoking cessation as first-line treatment. Long-term oxygen therapy has not been shown to have any long-term benefit in OSA patients.

D Modafinil

Modafinil is a daytime stimulant used in the treatment of narcolepsy. It is not used in the management of OSA. Modafinil has been shown in some small studies to have beneficial effects in use with continuous positive airway pressure (CPAP), but not as an alternative. Intra-oral devices may be useful for mild OSA, but not in daytime sleepiness.

A Intra-oral device to prevent snoring

Long-term oxygen therapy has not been shown to have any long-term benefit in OSA patients. Intra-oral devices may be useful for mild OSA, but not in daytime sleepiness.

B Long-term oxygen therapy

This can be used as an adjunct to CPAP therapy or in cases where patients are unable to adhere to CPAP. However, it should not be offered as first-line in preference to CPAP. Given this patient’s smoking history, it is likely that he is beginning to develop underlying chronic obstructive pulmonary disease. Although further investigation is warranted, including arterial blood gas analysis, oxygen saturations of 91% on air are not an indication for long-term oxygen therapy.

E Nocturnal lorazepam

This is an inappropriate treatment option for this patient as benzodiazepines are contraindicated in the context of OSA and their use can lead to respiratory failure.

72941

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 36-year-old woman who has visited her GP four times in the past year for sinusitis, and has been feeling that she has begun to develop a nose deformity, visits her GP complaining of shortness of breath. Additionally, she has suffered one episode of haemoptysis. On examination her facial appearance suggests a saddle-nose deformity.

Investigations reveal the following:

Investigation	**Result**	**Normal value**
Haemoglobin (Hb)	109 g/l	115–155 g/l
White cell count (WCC)	6.9×10^9/l	$4.0–11.0 \times 10^9$/l
Platelets (PLT)	195×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	139 mmol/l	135–145 mmol/l
Potassium (K^+)	5.6 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	178 μmol/l	50–120 μmol/l
Urine	Blood+, protein+	
cANCA	+	

A chest X-ray (CXR) is shown below:



Given the likely diagnosis, which of the following is the most appropriate treatment for her?

Your answer was correct

A	Prednisolone
B	Cyclosporine
C	Pulsed methylprednisolone and cyclophosphamide
D	Azathioprine

E Methotrexate

Explanation

C Pulsed methylprednisolone and cyclophosphamide

This woman has a classical picture of Granulomatosis with polyangiitis (GPA) with sinusitis initially, now progressing to pulmonary infiltration with haemoptysis and renal involvement as evidenced by urinary blood and protein signifying glomerulonephritis. The anaemia seen here and positive c-ANCA serve to further support the diagnosis. The chest x-ray shows bilateral upper zone nodules. Initial preferred therapy for GPA is with pulsed IV methylprednisolone and cyclophosphamide, although azathioprine or methotrexate may both be used for maintenance therapy or when there is not significant pulmonary vasculitis or renal impairment. Prognosis generally depends on early diagnosis and intervention, so recognising the features seen here is crucial.

Image source: https://www.flickr.com/photos/30950973@N03/5958079204 (https://www.flickr.com/photos/30950973@N03/5958079204)

A Prednisolone

Prednisolone alone is inadequate with respect to controlling Granulomatosis with polyangiitis (GPA), where use of a steroid-sparing agent allows the dose of prednisolone to be reduced and improves outcomes.

B Cyclosporine

Cyclosporine is an alternative to cyclophosphamide where it is not tolerated or is unsuitable. It is, however, used in combination with steroid therapy.

D Azathioprine

Azathioprine is an option as oral maintenance therapy for Granulomatosis with polyangiitis, or when there is mild disease.

E Methotrexate

Like azathioprine, methotrexate is an oral option for maintenance therapy in stable Granulomatosis with polyangiitis.

20543

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Granulomatosis with Polyangiitis

Media

Expanded explanations

Flashcards

A 49-year-old woman experiences long-standing, worsening shortness of breath. She smokes 15 cigarettes per day and has been provided with a salmeterol inhaler from her general practitioner (GP), to minimal benefit. She describes 3 kg of unintentional weight loss over the past six months but denies night sweats. She suffers from a chronic cough, which is non-productive.

Examination reveals crackles on auscultation and cervical lymphadenopathy.

Investigations:

C-reactive protein (CRP)	161 mg/l	(0–10 mg/l)
White cell count (WCC)	12.4 × 10^9/l	(4–11 × 10^9/l)

Her chest X-ray is shown below.



What is the most appropriate investigation to undertake to establish a diagnosis?

Your answer was incorrect

A Bronchoscopy and lavage

B Computed tomography (CT) thorax, abdomen and pelvis

C High-resolution CT

D Spirometry

E Tuberculin skin test

Explanation

A Bronchoscopy and lavage

In this case, the history is relatively non-specific. However, the chest X-ray reveals likely miliary tuberculosis (TB). This would benefit from staining with Ziehl–Neelsen stain and rapid direct microscopy for acid-fast bacilli. The patient has a non-productive cough; therefore, a sputum sample is not straightforward. Bronchoscopy and lavage would provide the diagnosis here.

Image original source: https://westjem.com/case-report/miliary-tuberculosis-coinfection-with-human-immunodeficiency-virus.html

C High-resolution CT

High-resolution CT would be of benefit to patients with lung fibrosis. However, this patient's chest X-ray reveals likely miliary TB, and therefore, bronchoscopy and lavage is the most appropriate diagnostic test.

B Computed tomography (CT) thorax, abdomen and pelvis

Any patient with significant weight loss should have malignancy considered in the differential diagnosis, which would benefit from CT. However, this chest X-ray reveals miliary TB. Therefore, bronchoscopy and lavage is the more appropriate answer.

D Spirometry

The history in this case is suggestive of chronic obstructive pulmonary disease (COPD). However, the chest X-ray indicated miliary TB. Therefore, spirometry would not be diagnostic for this patient.

E Tuberculin skin test

The diagnosis in this case is miliary TB, as suggested by the chest X-ray. However, the tuberculin skin test often leads to negative results in patients with miliary TB. Negative results would not exclude TB, and therefore, this is not a useful diagnostic test.

71774

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

National Institute for Health and Care Excellence. 2016. Tuberculosis. NICE guideline [NG33].
nice.org.uk/guidance/ng33/chapter/recommendations#managing-active-tb-in-all-age-groups
(https://www.nice.org.uk/guidance/ng33/chapter/recommendations#managing-active-tb-in-all-age-groups)

Tuberculosis

Media

Expanded explanations

Flashcards

A 54-year-old farmer’s wife with a history of rheumatoid arthritis presents to the clinic with increasing shortness of breath, cough and intermittent pleuritic chest pain. Previous agents have included corticosteroids and hydroxychloroquine, and she has commenced methotrexate, with appropriate folic acid replacement 2 years ago. On examination her BP is 135/85, with pulse 73 and regular. She has scattered crackles throughout both lung fields on auscultation, and there is no pitting oedema of the ankles. Lung function tests reveal a restrictive lung defect previous lung function tests were normal some 4 years ago.

Chest x-ray is shown below.



What diagnosis fits best with this clinical picture?

Your answer was incorrect

A	Extrinsic allergic alveolitis
B	Methotrexate-related lung toxicity
C	Hydroxychloroquine-related lung toxicity

D Obliterative bronchiolitis

E Idiopathic pulmonary fibrosis

Explanation

B Methotrexate-related lung toxicity

In this case the time course of lung disease (commencement of methotrexate 2 years ago, against a background of normal lung function 4 years ago) helps to point to methotrexate-related lung toxicity as the most likely cause of her symptoms. The chest x-ray shows diffuse interstitial markings. A spectrum of disease may occur with methotrexate toxicity, from interstitial pneumonitis to cough, pleuritic chest pain and shortness of breath related to subacute lung infiltration, associated with bilateral pleural effusions. Histology may demonstrate alveolitis, pneumonitis, epithelial cell hyperplasia, eosinophil infiltration and granuloma formation, and is more useful than when investigating other drug toxicities. Treatment is methotrexate withdrawal - in this case substitution with an anti-TNF or anti-CD20 antibody would be the next most logical step.

Image source: https://radiopaedia.org/cases/pulmonary-fibrosis-early?lang=gb (https://radiopaedia.org/cases/pulmonary-fibrosis-early?lang=gb)

C Hydroxychloroquine-related lung toxicity

Hydroxychloroquine toxicity primarily involves the eye, leading to retinal damage which may be permanent, and to deposition of the drug within the cornea.

A Extrinsic allergic alveolitis

Extrinsic allergic alveolitis typically presents with fever, fatigue, shortness of breath and cough. The symptoms are usually seen in the context of a trigger and resolve after removal of the trigger before reoccurring on re-exposure.

D Obliterative bronchiolitis

Obliterative bronchiolitis is characterised by fixed airways obstruction, rather than the restrictive lung defect seen here.

E Idiopathic pulmonary fibrosis

The fact that lung function was normal 4 years ago anchors the defect seen here to the initiation of methotrexate, making this far more likely to be the cause of the restrictive lung defect seen.

32440

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 55-year-old woman presents to the Emergency Department with a dry cough and shortness of breath, which she has had for the past four days. She has recently returned from a winter holiday to an all-inclusive hotel in Spain. She also complains of some abdominal discomfort, nausea, vomiting, diarrhoea and a headache. She was previously well before she developed these symptoms and is not on any medication.

On examination, she is unwell with a BP of 123/67 mmHg, a heart rate of 92 bpm and a respiratory rate of 22/min. On auscultation of her chest, he has bi-basal crackles. Her cardiovascular examination is unremarkable.

Investigations reveal the following:

Investigation	**Result**	**Normal Value**
Haemoglobin (Hb)	143 g/l	115–155 g/l
White cell count (WCC)	16.1 × 10^9/l	4.0–11.0 × 10^9/l
Platelets (PLT)	278 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	129 mmol/l	135–145 mmol/l
Potassium (K^+)	3.5 mmol/l	3.5–5.0 mmol/l
Creatinine (Cr)	112 µmol/l	50–120 µmol/l
Urea	8.1 mmol/l	2.5–6.5 mmol/l
C-reactive protein (CRP)	69 mg/l	< 10 mg/l

Her CXR is shown below:



What initial investigation is most likely to be helpful in elucidating the diagnosis?

Your answer was incorrect

A Urinary immunoglobulin titres

B Sputum microscopy and culture

C Stool microscopy and culture

D Bronchoalveolar lavage

E Urinary antigen testing

Explanation

E Urinary antigen testing

The most likely diagnosis is *Legionella pneumophila*. This usually infects middle-aged patients, more severely in smokers, is twice as common in men and has an incubation period of 2–10 days. It causes moderate leucocytosis (neutrophilia, lymphopenia), hyponatraemia, deranged LFTs, proteinuria, haematuria and myoglobinuria. The CXR here shows predominantly mid to lower zone opacity, worse on the left than the right. Diagnosis: urinary antigen testing, rise in specific IgM and IgG titres in the urine, blood and sputum or by direct immunofluorescent staining of the organism via bronchial washings. With a CURB-65 score of 1, indicating mild disease, treatment would be dependent on local guidelines but may consist of a macrolide, such as azithromycin or clarithromycin.
Image source: https://radiopaedia.org/cases/legionella-pneumonia

B Sputum microscopy and culture

Patients with pneumonia should have a sputum MCS sample sent. *Legionella*, as this patient has, can be diagnosed from growth by a sputum sample, but that takes time and is therefore not as helpful as urinary antigen testing, which is much faster.

A Urinary immunoglobulin titres

This is not a commonly used investigation in infection.

C Stool microscopy and culture

Despite the patient having diarrhoea, abdominal pain and a travel history, there are also respiratory signs and symptoms. This makes a respiratory pathogen far more likely than a gastrointestinal one, and so a stool sample is unlikely to provide any useful information.

D Bronchoalveolar lavage

A bronchoalveolar lavage can be a useful way to obtain sputum in a patient who is unable to expectorate a sample. It can be particularly useful in ITU patients, to diagnose TB or to diagnose PCP. It is generally not necessary when the patient is able to cough up a sputum sample.

7509

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 21-year-old man with a history of cystic fibrosis comes to the Respiratory Clinic for review. He had been admitted with a significant exacerbation some 2 months earlier and has been gradually losing weight since then. On examination, his blood pressure is 132/82 mmHg and his heart rate is 75/min and regular. You can hear coarse crackles and wheeze throughout both lung fields on auscultation. On examining his blood results, you note that he has a fasting plasma glucose of 8.5 mmol/l and his HbA1c is 53 mmol/mol.

Which of the following is the most appropriate way to manage his blood sugar?

Your answer was correct

A Empagliflozin

B Gliclazide

C High-carbohydrate diet and basal-bolus insulin

D Low-carbohydrate diet

E Metformin

Explanation

C High-carbohydrate diet and basal-bolus insulin

In this situation, where patients with cystic fibrosis and gradual weight loss because of exocrine pancreatic insufficiency and frequent infections, a diagnosis of diabetes only compounds their difficulties. Treatment involves maintaining carbohydrate intake and using enough insulin to maintain glucose control.

A Empagliflozin

SGLT-2 inhibitors are not an appropriate option for control of blood glucose in this situation; they in essence remove carbohydrate by promoting urinary excretion of glucose, which leads to a calorie deficit, exactly what patients with cystic fibrosis don’t need.

B Gliclazide

Although gliclazide may stimulate residual insulin production to attain some improvement in glucose control, this is likely to be short term and inadequate compared to a basal-bolus insulin regime.

D Low-carbohydrate diet

A low-carbohydrate diet here is not appropriate because it will potentially drive further weight loss without achieving adequate glucose control.

E Metformin

Insulin resistance is not the primary problem in diabetes associated with cystic fibrosis; as such, metformin is ineffective in reducing blood sugar here.

7624

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 56-year-old woman presents to the Emergency Department with palpitations and shortness of breath. She is known to have mild chronic obstructive pulmonary disease (COPD) and is on the following inhalers: salbutamol, ipratropium bromide and beclomethasone; according to her notes, she does have a mild degree of reversibility. On examination, the patient is dyspnoeic with BP 154/88 mmHg and a heart rate of >100/min (AF). She underwent echocardiography six months ago, which showed normal ventricular function and no structural abnormality.

Her admission arterial blood gases (ABG) are:

Investigation	Result	Normal Value
pH	7.35	7.35–7.45
$p_a(O_2)$	8.1 kPa	10.5–13.5 kPa
$p_a(CO_2)$	6.2 kPa	4.6–6.0 kPa
Bicarbonate (HCO_3^-)	28 mmol/l	24–30 mmol/l

You start the patient on 28% oxygen, nebulisers (salbutamol 5 mg and ipratropium 0.5 mg) and repeat the blood gases after 30 min:

pH	7.36	7.35–7.45
$p_a(O_2)$	9.6 kPa	10.5–13.5 kPa
$p_a(CO_2)$	5.8 kPa	4.6–6.0 kPa
Bicarbonate (HCO_3^-)	29 mmol/l	24–30 mmol/l

You reassure her that she is improving on the current medication, but she tells you that she is still having palpitations. You do a 12-lead electrocardiogram (ECG), which reveals multi-focal atrial tachycardia (MAT) with a rate of 118/min. Her BP has fallen to 110/70 mmHg.

How would you control this arrhythmia?

Your answer was incorrect

A DC – cardioversion

B Start metoprolol

C Start verapamil

D Increase inspired oxygen to 40%

E Observation only

Explanation

C Start verapamil

MAT commonly occurs in critically ill patients with obstructive airways disease who may be hypoxic and hypercapnic. It is characterised by at least three different P-wave morphologies with varying PP and PR intervals. Aim to improve both $p_a(O_2)$ and $p_a(CO_2)$.

Whilst AF control may not be required as her pulmonary status improves, it is likely that in the acute phase, especially with repeated salbutamol nebulisers, that she will suffer from worsening symptoms of tachycardia. As such, verapamil would be an appropriate option. It can be given in small boluses of 5 mg intravenously and then as an oral dose of 40–120 mg TDS. It is a moot point whether she will require long-term AF control. If following the acute episode she remains in sinus rhythm, a 72 h ECG monitor may be appropriate prior to committing her to new medication.

B Start metoprolol

Metoprolol could make her COPD worse when used acutely as she is noted to have a degree of reversibility. This is not the same as chronic use of beta blockers in COPD, where evidence suggests they don't impact significantly on lung function.

A DC – cardioversion

DC cardioversion would be an inappropriate option in this patient, particularly given it may lead to more serious rhythm disturbance in multi-focal atrial tachycardia. It should be done in an emergency situation if the patient becomes haemodynamically unstable or develops myocardial infarction as a result of the arrhythmia. If this happens and the patient is conscious, then sedation can be given. It is important that the shock delivered is synchronised to avoid shocking on a T wave, which could lead to asystole.

D Increase inspired oxygen to 40%

This would be inappropriate as it would not deal immediately with the underlying problem of the arrhythmia, which is important as the patient's blood pressure has become lower. Also, the hypoxia is already improving, and increasing oxygen may worsen the patient's hypercapnia.

E Observation only

This arrhythmia has become significant in that it has caused a drop in blood pressure in a very unstable patient. Therefore it would be inappropriate to simply observe.

7504

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

You are asked to comment on some arterial blood gas results. You understand the sample was taken from a 64-year-old patient attending one of the outpatient clinics. The clinical information says 'female smoker with shortness of breath, wheeze on auscultation of the chest'.

Investigations reveal the following:

Investigation	Result	Normal value
pH	7.38	7.35–7.45
pCO_2	6.5 kPa	4.6–6.0 kPa
pO_2	8.0 kPa	10.5–13.5 kPa
Bicarbonate (HCO_3^-)	31 mmol/l	24–30 mmol/l

Which of the following is the most likely underlying diagnosis?

Your answer was correct

A Diuretic therapy

B Bartter syndrome

C Chronic COPD

D Acute exacerbation of COPD

E Conn syndrome

Explanation

C Chronic COPD

This picture is consistent with chronic COPD, as exhibited by CO_2 retention in the presence of a pH within the normal range. What has occurred over time is renal compensation with conservation of bicarbonate, leading to the elevated HCO_3^- level of 31 mmol/l which is seen here.

A Diuretic therapy

Although diuretic therapy does result in a metabolic alkalosis, the raised bicarbonate seen here is as a result of compensation for chronic respiratory acidosis.

B Bartter syndrome

Bartter syndrome is associated with metabolic alkalosis and normal blood pressure. It often presents in infancy as a result of salt wasting. The elevated bicarbonate is inconsistent with Bartter as the underlying diagnosis.

D Acute exacerbation of COPD

An acute exacerbation is likely to be associated with respiratory acidosis due to further CO_2 retention, and it's unlikely the patient would be attending the outpatient's clinic.

E Conn syndrome

Conn syndrome is associated with metabolic alkalosis and hypertension, but not with the CO_2 retention seen here.

18627

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Chronic Obstructive Pulmonary Disease

Media

Expanded explanations

Flashcards

A 47-year-old hotel cleaner presents with a fever of 39°C, malaise and myalgia followed by the development of a dry and non-productive cough. Other non-specific symptoms include headache, anorexia, nausea and non-specific abdominal pain. Positive findings on general examination include tachycardia, tachypnoea and bibasal crackles on auscultation of the chest. The CXR shows early bibasal consolidation. Blood investigations show hyponatraemia and deranged LFTs, and urine testing shows the presence of protein and blood.

Which of the following is the most appropriate initial antibiotic regimen?

Your answer was correct

A Benzylpenicillin and gentamicin

B Amoxicillin

C Azithromycin

D Oxytetracycline

E Rifampicin

Explanation

C Azithromycin

The picture here, with marked fever, hyponatraemia, GI disturbance and blood and proteinuria, fits well with a diagnosis of Legionnaires' disease. His CURB-65 score classifies the pneumonia as mild. A reasonable option therefore are macrolides such as azithromycin or clarithromycin.

A Benzylpenicillin and gentamicin

This was the standard empirical intervention for bacterial endocarditis, although gentamicin now appears not to impact significantly on prognosis versus benzylpenicillin alone.

B Amoxicillin

Whilst amoxicillin is an appropriate first line option in community acquired pneumonia the constellation of symptoms here hints toward an underlying diagnosis of Legionnaires' disease. As such, amoxicillin is ineffective.

D Oxytetracycline

Oxytetracycline is an alternative, rather than a first- or second-line option, for the treatment of Legionnaires' disease.

E Rifampicin

Rifampicin is used in combination with a quinolone or a macrolide for treatment of severe *Legionella* infection.

71082

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media Expanded explanations Flashcards

A 60-year-old man presents to the Respiratory Clinic for follow-up. He was diagnosed with chronic obstructive pulmonary disease ten years ago and is currently on long-term oxygen therapy. He reports slight worsening of exertional dyspnoea. He has had no exacerbations and has not required hospitalisation in the past year. He has not smoked cigarettes for the past two years.

On examination, his blood pressure is 124/78 mmHg and his heart rate is 80 bpm. The respiratory physician wants to consider listing the patient for lung transplantation.

Which of the following findings would prompt the respiratory physician to list the patient for lung transplantation?

Your answer was incorrect

A Body mass index 20 kg/m^2

B Forced expiratory volume in one second (FEV1) 14% of predicted (490 ml)

C No acute exacerbations in the past year

D $p_a(O_2)$ 8.4 kPa

E $p_a(CO_2)$ 6.9 kPa

Explanation

B Forced expiratory volume in one second (FEV1) 14% of predicted (490 ml)

To be eligible for a lung transplant, patients must be on maximal medical therapy and be predicted to have a survival of at least 5 years, at an acceptable quality of life for the patient, post-transplant. In addition, patients with chronic obstructive pulmonary disease (COPD) must meet one of the following criteria:

- FEV1 < 20% of predicted and a diffusion capacity of the lungs for carbon monoxide of less than 20% predicted, or uniform distribution of emphysema throughout the lungs.
- Increasing frequency of hospitalisation with hypercapnia and hypoxia.
- Pulmonary hypertension or cor pulmonale.
- BODE score ≥ 7

A BODE classification (B – Body mass index; O – airflow Obstruction; D – Dyspnoea; E – Exercise capacity) score is an indicator functional capacity. The score ranges from 0-10, a higher score indicates a poorer prognosis.

C No acute exacerbations in the past year

No acute exacerbation in the past year is not an indication for lung transplantation. Increasing frequency of hospitalisations is an indication to list the patient for lung transplantation.

A Body mass index 20 kg/m^2

A body mass index of < 21 kg/m^2 indicates a more severe disease, according to the BODE classification. A BODE score of ≥ 7 is a criterion for transplantation listing. However, the body mass index is not a criterion to list for transplantation independently.

D $p_a(O_2)$ 8.4 kPa

A $p_a(O_2)$ of 8.4 kPa is indicative of hypoxaemia and it is a criterion for referring the patient for further evaluation regarding the need for transplantation, rather than an indication for transplantation by itself.

E $p_a(CO_2)$ 6.9 kPa

A $p_a(CO_2)$ of 6.9 kPa is indicative of hypercapnia and it is a criterion for referring the patient for further evaluation regarding the need for transplantation, rather than an indication for transplantation by itself. 72781

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Transplantation

Expanded explanations

A 56-year-old man is referred to the outpatients department with a 2-year history of increasing breathlessness and deteriorating exercise tolerance. He has a non-productive cough and always feels tired, so much so that he has needed to ask his wife to help him look after his pigeons, which he has been racing for many years. Over the last year he has lost about 1 stone in weight. He has a past medical history of asthma as a child. In his social history, he has had no history of recent foreign travel, except to France. He has always worked in a bank and is an ex-smoker who drinks minimal amounts of alcohol.

On examination, he is not clubbed. On auscultation of his chest, he has fine inspiratory crackles.

ABGs on air reveal:

pH	7.36
PO_2	7.4 kPa
PCO_2	4.7 kPa

Lung function tests:

FEV_1	70% predicted
FVC	64% predicted
KCO	53% predicted
TLCO	55% predicted

His chest X-ray is shown below:



What is the most likely diagnosis?

Your answer was correct

A	Bronchopulmonary aspergillosis
B	Extrinsic allergic alveolitis
C	Idiopathic pulmonary fibrosis
D	Psittacosis
E	Sarcoidosis

Explanation

B Extrinsic allergic alveolitis

The patient has interstitial lung disease, demonstrated by the slowly progressive shortness of breath, type I hypoxia and restrictive lung function tests with reduced TLCO and KCO. In the chest radiograph there is reticular nodular shadowing. The diagnosis is extrinsic allergic alveolitis secondary to pigeons – bird fancier's lung. This is caused by inhaled avian serum proteins (usually IgA) present in the pigeon's feathers and excreta. It is a hypersensitivity pneumonitis caused by a specific immunological response to inhaled organic dusts. EAA may present either as an acute or a chronic disease. Acute disease develops several hours after exposure to high concentrations of the allergen. Breathlessness and flu-like symptoms occur sometimes with headaches, fever and myalgia. Symptoms usually resolve within 48 h. The patient has chronic disease, as it has progressed to interstitial lung disease with fibrosis.

The diagnosis of extrinsic allergic alveolitis is made by:

- Identifying a potential source of antigen
- Characteristic clinical features: this patient has chronic EAA and presents with similar features to pulmonary fibrosis – clubbing tends to occur later (idiopathic pulmonary fibrosis would typically be associated with clubbing)
- Characteristic radiology (usually upper lobe fibrosis)
- Lung function tests – showing a restrictive defect with reduced carbon monoxide gas transfer
- Demonstrating precipitating antibodies (precipitins) in the patient's serum
- Resolution or improvement of symptoms following avoidance of exposure to the cause
- Histology of lung tissue biopsy showing a mononuclear cell infiltrate with the formation of non-caseating granulomas
- Fluid obtained from bronchoalveloar lavage has a high lymphocyte count (typically >40%).

Treatment is avoidance of the antigen. Corticosteroids may improve recovery in acute attacks but do not seem to provide long-term benefit. They are not usually helpful once fibrosis has occurred.

Other causes of EAA include:

- Farmer's lung, due to *Micropolyspora faeni* and *Thermactinomyces* in mouldy hay, straw and grain
- Bagassosis due to *Thermactinomyces sacchari* in sugar cane processing
- Malt worker's lung due to *Aspergillus clavatus*
- Mushroom worker's lung due to thermophilic actinomycetes
- Ventilation and water-related contamination, e.g. due to thermophilic actinomycetes contaminating air-conditioning systems
- Veterinary workers and animal handlers
- Workers in milling and construction, e.g. wood dust pneumonitis due to *Alternaria* spp.

Causes of upper-zone fibrosis:

- Progressive massive fibrosis/coal worker's pneumoconiosis
- Ankylosing spondylitis
- Sarcoid/silicosis
- Tuberculosis
- Extrinsic allergic alveolitis
- Radiation (depending on site).

A Bronchopulmonary aspergillosis

The chest radiograph shows diffuse infiltrates. CXR findings in ABPA are more likely to include proximal bronchiectasis, lobar collapse or discrete pulmonary infiltates.

C Idiopathic pulmonary fibrosis

There is a clear precipitant to interstitial lung disease here.

D Psittacosis

Psittacosis is an infection caused by *Chlamydia psittaci*. It usually presents with a sudden onset of fever, dry cough and headache following a recent exposure to birds. Psittacosis is typically linked to parrots. The duration of symptoms is too long for psittacosis, and the chest X-ray changes are too widespread.

E Sarcoidosis

Exposure to pigeons and weight loss make EAA more likely than sarcoidosis.

70114

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Average

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 38-year-old woman underwent a heart and lung transplantation for pulmonary hypertension. Whilst she was initially well post surgery she presents to the clinic some 15 months later with progressive shortness of breath. On examination her BP is 125/82 mmHg and there is wheeze on auscultation of her chest.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin	121 g/l	115–155 g/l
White cell count (WCC)	4.9 × 10^9/l	4–11 × 10^9/l
Platelets	184 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	140 mmol/l	135–145 mmol/l
Potassium (K^+)	4.0 mmol/l	3.5–5.0 mmol/l
Urea	6.9 mmol/l	2.5–6.5 mmol/l
Creatinine	120 µmol/l	50–120 µmol/l
Pulmonary function tests	Obstructive picture, irreversible with salbutamol, worsening over the past 3 months	

Which of the following is the most likely diagnosis?

Your answer was incorrect

A Post-transplant lymphoproliferative disorder

B Acute rejection

C Bronchiolitis obliterans syndrome

D New-onset asthma

E Recurrence of pulmonary hypertension

Explanation

C Bronchiolitis obliterans syndrome

Bronchiolitis obliterans syndrome occurs due to progressive obliterative bronchiolitis affecting terminal bronchioles. The condition is manifest in increasing shortness of breath and a progressively worsening fixed airways obstruction picture on pulmonary function testing. Risk factors for the development of the condition include early acute rejection and CMV infection. Bronchiolitis obliterans has been seen as early as 3 months post-transplant, and is responsible for over 50% of deaths after the first year.

Pulmonary hypertension (PAH) is defined by a mean pulmonary artery pressure greater than 25 mmHg at rest and a normal pulmonary capillary wedge pressure of 15 mmHg or less with a pulmonary vascular resistance greater than 3 Wood units. Prior to lung transplantation being available, median survival for patients with pulmonary hypertension was 2.8 years.

B Acute rejection

Acute rejection may occur any time from the first week after the transplant to 3 months afterwards. All recipients have some amount of acute rejection.

A Post-transplant lymphoproliferative disorder

In this condition there is controlled B-cell proliferation due to the need for immunosuppressive therapy after transplantation. The clinical presentation can vary from simple lymphoid hyperplasia to aggressive disease that closely resembles non-Hodgkin's lymphoma. Post-transplant lymphoproliferative disorder is still a rare disease, despite being more commonly observed after lung transplantation than with other organs.

D New-onset asthma

In asthma, you would see obstructive spirometry, but you would expect reversibility with salbutamol.

E Recurrence of pulmonary hypertension

Recurrence of the disease is rare after transplantation.

21436

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Lung Transplantation

Expanded
explanations

A 41-year-old man is referred to the hospital with a 1-week history of fever, malaise, increasing confusion, cough and breathlessness. The cough was initially dry but has become productive of yellow sputum in the last few days. He saw the GP 4 days ago, who prescribed amoxicillin with no benefit. His wife reports that he has also been suffering from abdominal pain and vomiting. He is a smoker of 20/day and works in a large law firm.

On examination he looks unwell and dehydrated. Temperature 39°C, BP 100/50 mmHg, pulse 120/min, respiratory rate 30/min, SpO_2 89% on air and Glasgow Coma Scale score 12. Auscultation of his chest reveals crackles in the left lower zone. His abdomen is generally tender on palpation. Examination is otherwise unremarkable. Urinalysis shows blood +++.

Investigations:

Investigation	Result	Normal Values
Haemoglobin (Hb)	111 g/l	135 - 135 g/l
White cell count (WCC)	10.6 × 10^9/l	4.0 - 11.0 × 10^9/l
Neutrophils	9.2 × 10^9/l	1.5 - 7.0 × 10^9/l
Platelets (PLT)	151 × 10^9/l	150 - 400 × 10^9/l
Sodium (Na^+)	127 mmol/l	135 - 145 mmol/l
Potassium (K^+)	3.4 mmol/l	3.5 - 5.0 mmol/l
Urea	9.0 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	120 μmol/l	50 - 120 μmol/l
Bilirubin	28 μmol/l	1 - 22 μmol/l
Aspartate aminotransferase (AST)	220 u/l	10 - 40 u/l
Alanine aminotransferase (ALT)	190 u/l	7 - 55 u/l
Alkaline phosphatase (ALP)	140 u/l	30 - 150 u/l
Albumin	28 g/l	35 - 55 g/l
C-reactive protein (CRP)	110 mg/l	< 10 mg/l

Arterial blood gases (ABG) on air	Result
pH	7.39
PCO_2	4.66 kPa
PO_2	7.3 kPa
Bicarbonate	21.4 mmol/l

Which organism is most likely to be responsible?

Your answer was incorrect

A *Streptococcus pneumoniae*

B *Legionella pneumophila*

C *Mycoplasma pneumoniae*

D *Coxiella burnetti*

E *Chlamydia psittaci*

Explanation

B *Legionella pneumophila*

Legionella pneumophila is an aerobic, waterborne, Gram-negative, non-motile bacillus and has three epidemiological patterns:

- Outbreaks due to contaminated water-cooling systems, air-conditioning or showers. May be an occupational history or recent hotel holiday. Usually occurs in previously fit individuals. This man worked in a law firm and it is presumed this contains an air-conditioning system.
- Sporadic cases. Rare at extremes of age. Usually age 40–70. More common in men than in women. At-risk groups: smokers, chronic illness – especially chest disease, diabetes mellitus – and alcoholics.
- Outbreaks occurring in immunosuppressed individuals, e.g. those on corticosteroids.

pneumophila can cause two distinct disease entities:

- Legionnaires' disease, which is characterised by pneumonia.
- Pontiac fever, which is a milder, self-limiting illness consisting of fever and myalgia but no pneumonia.

2–15% of community-acquired pneumonias requiring hospitalisation globally are secondary to *L. pneumophila*. Mortality is 10% in treated patients who are not immunocompromised, rising to nearly 50% in nosocomial infections.

The incubation period of *L. pneumophila* is 2–10 days and it typically presents with flu-like symptoms – fevers, malaise, myalgia and headaches; the patient later develops a cough (often non-productive), breathlessness and confusion; 50% develop gastrointestinal symptoms (usually diarrhoea and vomiting). Patients may also experience pleuritic chest pain and episodes of haemoptysis.

Investigations may show: hyponatraemia (due to SIADH), lymphopenia, haematuria, proteinuria and hypoxia. Abnormal liver biochemistry (especially aminotransferases) and renal function occur in 50% of patients.

Diagnosis: the quickest way is to test for *Legionella* antigen in urine, and direct immunofluorescent staining of the organism in sputum, pleural fluid or bronchial washings. *L. pneumophila* is a fastidious organism. Gram stain does not usually reveal an organism and culture on standard media is often negative.

Chest X-ray usually shows lobar and then multilobar shadowing. Small pleural effusions may occur.

Treatment is with a macrolide. The other clue in this question is the lack of response to beta-lactam antibiotics.

A *Streptococcus pneumoniae*

Whilst *Streptococcus pneumoniae* is the commonest cause of community-acquired pneumonia (accounting for 50–70% of cases), this gentleman has features of an atypical pneumonia: initial dry cough, lack of response to beta-lactam antibiotics and gastrointestinal disturbance, in addition to hyponatraemia and deranged liver function. Herpetic cold sores are associated with pneumococcal pneumonia and may be specifically mentioned in an exam question relating to pneumococcal pneumonia.

C *Mycoplasma pneumoniae*

Pneumonia caused by *Mycoplasma pneumoniae* typically affects young adults but usually has a longer prodrome of 2 weeks or more. White cell count may be normal. Cold agglutinins may be present in 50% of patients. Extra-pulmonary manifestations include pericarditis, myocarditis, rashes such as erythema multiforme and erythema nodosum, neurological pathology such as cranial or peripheral nerve palsies, and meningoencephalitis. Thrombocytopenia and haemolytic anaemia may occur, which are not suggested by the blood results given for this question.

D *Coxiella burnetti*

Coxiella burnetii infection is associated with Q fever. In acute Q fever, 60% of patients are asymptomatic. The incubation period varies from 2 to 6 weeks, and there are three main clinical presentations:

- A self-limiting, flu-like illness with abrupt onset of fever, headache, myalgia and arthralgia. This presentation does not fit with the case described.
- Pneumonia (predominant in North America), usually of mild severity and often an incidental finding on CXR. If patients have symptoms, typically there will be a dry cough, breathlessness and pleuritic chest pain. Again this presentation does not fit with the case described.
- Hepatitis (predominant in Europe), usually with mild elevation of LFTs. Nausea and vomiting can occur but diarrhoea is rare. Whilst some elements of the case described fit with this, there are many other symptoms and results in the case which do not, making *L. pneumophila* more likely.

E *Chlamydia psittaci*

Pneumonia due to *Chlamydia psittaci* is termed psittacosis. It is an infection acquired from avian reservoirs, and is classically associated with parrots. Incubation period is 5–14 days. Common symptoms include fever, chills and malaise, cough (usually dry), sore throat, mild pharyngitis, epistaxis and severe headache. Whilst some of these symptoms are described in this case, the lack of mention of avian contact, along with several features more in keeping with *L. pneumophila*, make this option unlikely.

70073

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 31-year-old woman presented to the Respiratory Outpatients Department with a 3-year history of increasing shortness of breath and cough. Her symptoms are constant and not episodic. She had required numerous courses of antibiotics in the last few years. She smoked five packs a day for about five years. Respiratory examination revealed a hyperinflated chest and a mild wheeze throughout.

Lung function tests:

Investigation	**Result**
FEV_1	50% predicted
FVC	80% predicted
TLCO	68% predicted
KCO	71% predicted
RV:TLC No bronchodilator reversibility	Increased

Which investigation would be most useful in making a definitive diagnosis?

Your answer was incorrect

A Histamine challenge test

B Sweat test

C Alpha-1 anti trypsin deficiency genotyping

D Reversibility testing

E High-resolution CT

Explanation

C Alpha-1 anti trypsin deficiency genotyping

Alpha-1-antitrypsin deficiency is a rare autosomal recessive disease associated with the development of premature pan-lobular emphysema, most severe in the basal areas of the lungs. Emphysema is thought to result from an imbalance in the lung between neutrophil elastase (which destroys elastin) and the elastase inhibitor α1-antitrypsin (which protects against proteolytic degradation by elastin). The decline in lung function is accelerated in smokers.

Patients usually present with progressive increasing shortness of breath and weight loss. Cor pulmonale and polycythaemia occur later in the course of the disease. A chest X-ray usually shows bilateral bi-basal emphysema. Lung function tests show an obstructive picture and the RV:TLC ratio is raised, indicating gas trapping. Transfer factor will be reduced. There are a number of different phenotypes depending on the serum level of α1-antitrypsin:
Pi – protease inhibitor
M – normal allele
S, X and null are disease alleles

Phenotype	α1-Antitrypsin level	Disease
PiMM	Normal	
PiMS	80%	
PiMZ	60%	
PiSS	60%	
PiSZ	40%	Liver cirrhosis/premature emphysema
Pi Null Null	Nil	Premature emphysema

Treatment is similar to that of asthma. Alpha1-antitrypsin replacement can be given weekly or monthly and is recommended for those with very low serum levels and abnormal lung function. Lung transplantation should be considered in advanced disease. Patients should be encouraged to stop smoking.

A Histamine challenge test

This would be used to confirm a diagnosis of asthma. Histamine challenge is rarely performed because of technical difficulties in performing it, including the risk of provoking severe bronchoconstriction.

B Sweat test

This would be used to confirm a diagnosis of cystic fibrosis, where levels of sodium and chloride are measured in sweat, production of which is stimulated by pads soaked in pilocarpine.

D Reversibility testing

This would be used to confirm a diagnosis of asthma, rather than the picture consistent with alpha-1-antitrypsin deficiency seen here.

E High-resolution CT

This would be helpful in confirming basal predominant pan-lobular emphysema, making the diagnosis likely, but α1-antitrypsin levels would be more helpful in making definitive diagnosis.

70193

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Alpha-1 Antitrypsin Deficiency

Expanded explanations

A 68-year-old man who has a history of asthma is admitted to hospital with severe community-acquired pneumonia. You understand that he has a part-time job in a pet shop. He is severely hypoxic on admission and is transferred immediately to the Intensive Therapy Unit (ITU). He is treated with meropenem but, after 5 days, is still pyrexial and requiring significant inotropic and ventilatory support.

Investigations:

Investigation	**Result**	**Normal value**
Haemoglobin (Hb)	119 g/l	135–175 g/l
White cell count (WCC)	13.2 × 10^9/l	4–11 × 10^9/l
Platelets (PLT)	221 × 10^9/l	150–400 × 10^9/l
Sodium (Na^+)	135 mmol/l	135–145 mmol/l
Potassium (K^+)	4.5 mmol/l	3.5–5.0 mmol/l
Creatinine	145 µmol/l	50–120 µmol/l
C-reactive protein (CRP)	82 mg/l (< 10)	0–10 mg/l
Chest X-ray (CXR)	Shown below	



By James Heilman, MD [CC BY-SA 3.0 (https://creativecommons.org/licenses/by-sa/3.0) or GFDL (http://www.gnu.org/copyleft/fdl.html)], from Wikimedia Commons

Which of the following is the most likely diagnosis?

Your answer was incorrect

A *Aspergillus*

B *Staphylococcus aureus*

C *Streptococcus pneumoniae*

D *Klebsiella pneumoniae*

E *Chlamydia psittaci*

Explanation

E *Chlamydia psittaci*

The suspicion is that the poor response to broad-spectrum antibiotics is indicative of either an infection with an atypical pneumonia, such as *Chlamydia*, or the possibility of a fungal infection. The chest X-ray appearance of a left upper lobe consolidation is more typical of *Chlamydia* than *Aspergillus*. Tetracyclines are the therapy of choice for *Chlamydia* pneumonia, with macrolides as the second option, although there is a significant percentage of treatment failures on erythromycin. The other clue in the history to this man's underlying diagnosis is the fact that he works in a pet shop and may have had exposure to bird faeces.

C *Streptococcus pneumoniae*

Streptococcal pneumonia is the most common type of pneumonia but has a higher incidence in the elderly, post-splenectomy, and in those with pre-existing heart or lung disease. Patients usually present with fever and pleurisy and can develop sepsis. Chest X-ray demonstrates consolidation of the left upper lobe, and diagnosis can be confirmed with urinary antigen.

A *Aspergillus*

Aspergillus can cause various infections of the lungs. In asthmatics, there can be a type I hypersensitivity reaction or allergic bronchopulmonary aspergillosis. The latter causes bronchoconstriction with persistent inflammation, leading to bronchiectasis. Aspergilloma can occur within a pre-existing cavity, such as from previous tuberculosis or sarcoidosis, and invasive aspergillosis can occur in immunocompromised patients. This patient does not have risk factors for an acute *Aspergillus* reaction.

B *Staphylococcus aureus*

Staphylococcus pneumonia can occur as a complication of influenza or in intravenous drug users, as well as at extremes of age. It causes bilateral cavitating bronchopneumonia. The absence of risk factors makes this an unlikely cause.

D *Klebsiella pneumoniae*

Klebsiella can cause cavitating pneumonia, particularly in alcoholics and the elderly. It often affects the upper lobes. It commonly displays antibiotic resistance. The location of the pneumonia makes this less likely.

23355

Rate this question:

Next Question

Previous Question Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Pneumonia

Media

Expanded explanations

Flashcards

A 64-year-old man presents to the Emergency Department with a 1-day history of increasing shortness of breath. He also complains of sharp chest pain over the left side of his chest. His past medical history includes chronic obstructive pulmonary disease (COPD), for which he takes high-dose fluticasone and salmeterol (combined).

On examination, he has a respiratory rate of 25 breaths per minute, with oxygen saturations of 80% on room air. There is reduced air entry on the left side of his chest and there is hyperresonance to percussion.

A chest X-ray demonstrates a 1.6-cm left-sided pneumothorax. Needle aspiration is completed and the pneumothorax has reduced to 1.1 cm.

Which of the following is the most appropriate next step?

Your answer was incorrect

A Admit for 24 hours supplemental oxygen

B Chest drain

C Discharge with routine respiratory follow-up

D Chemical pleurodesis

E Repeat needle aspiration

Explanation

B Chest drain

Inserting a chest drain would be most appropriate for this patient. This patient has a secondary pneumothorax, given the past medical history of chronic obstructive pulmonary disease (COPD). According to British Thoracic Society (BTS) guidelines, in patients with a secondary pneumothorax following needle aspiration, if the pneumothorax has not reduced to < 1 cm in size, a chest drain requires insertion.

C Discharge with routine respiratory follow-up

Discharge of patients with pneumothorax applies if a primary pneumothorax has occurred and has been successfully aspirated. These patients are then typically followed up in 2–4 weeks. However, this is not true for patients with a secondary pneumothorax.

A Admit for 24 hours supplemental oxygen

Admission for supplemental oxygen would not be appropriate here. If the pneumothorax has reduced to < 1 cm in size, then supplemental oxygen would be necessary. However, a pneumothorax of > 1 cm in a patient with a secondary pneumothorax requires chest drain insertion.

D Chemical pleurodesis

Pleurodesis is usually reserved for recurrent pneumothoraces. It is too premature to refer this patient for chemical pleurodesis without having been treated with a chest drain initially.

E Repeat needle aspiration

Repeat aspiration is not required. In the event that air aspiration is unsuccessful, insertion of an 8–14 Fr chest drain is the next step.

72927

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Difficult

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

External Links

British Thoracic Society (BTS): BTS Pleural Disease Guideline 2010 Management of spontaneous pneumothorax
thorax.bmj.com/content/65/Suppl_2/ii18
(https://thorax.bmj.com/content/65/Suppl_2/ii18)

Pneumothorax

Media

Expanded explanations

Flashcards

An 80-year-old gentleman presents to the Emergency Department with gradually worsening shortness of breath of 4 days' duration. He has a fever and cough productive of greenish sputum. He is a known patient with chronic obstructive pulmonary disease (COPD) and is on home nebulisers (ipratropium bromide). He can walk only 50 metres on the flat without shortness of breath and cannot climb the stairs. His past medical history includes: a myocardial infarction 7 years ago, intensive care admission for his COPD 2 years ago and type 2 diabetes mellitus (diet-controlled). On examination, he is alert but febrile, with a blood pressure of 159/92 mmHg and a pulse of 92/min and regular. He has basal crackles more marked on the right, and scattered rhonchi in both lung fields. His abdominal examination is unremarkable. He has received optimum initial treatment in ED and been placed on controlled oxygen at FiO2 40% and SpO2 90% for the last 30 minutes.

Investigations reveal the following:

Investigation	**Result**	**Normal value**
Haemoglobin	113 g/l	135–175 g/l
White cell count (WCC)	15.2×10^9/l	$4–11 \times 10^9$/l
Platelets	233×10^9/l	$150–400 \times 10^9$/l
Sodium (Na^+)	136 mmol/l	135–145 mmol/l
Potassium (K^+)	3.7 mmol/l	3.5–5.0 mmol/l
Creatinine	111 μmol/l	50–120 μmol/l
pH	7.26	7.35–7.45
paO_2	7.9 kPa	10.5–13.5 kPa
$paCO_2$	7.4 kPa	4.6–6.0 kPa
Bicarbonate	30 mmol/l	24–30 mmol/l
Urea	7.0 mmol/l	2.5–6.5 mmol/l
Mean corpuscular volume (MCV)	83 fl	76–98 fl

His chest x-ray is shown below.



Which one of the following would be the optimal way to manage his ventilatory support?

Your answer was incorrect

A High flow oxygen (15 l)

B Bi-level positive airway pressure (BiPAP)

C 24% oxygen

D Continuous positive airway pressure (CPAP)

E Intubate and ventilate the patient

Explanation

B Bi-level positive airway pressure (BiPAP)

We can see from the patient's arterial blood gases that he is in type 2 respiratory failure with CO_2 retention, probably secondary to a right sided lobar pneumonia. BIPAP will help with gas transfer, reducing CO_2 levels. BIPAP is highly effective in this situation and can have a significant impact on mortality and morbidity.

Image source: https://commons.wikimedia.org/wiki/File:X-ray_of_COPD_exacerbation_-_anteroposterior_view.jpg (https://commons.wikimedia.org/wiki/File:X-ray_of_COPD_exacerbation_-_anteroposterior_view.jpg)

C 24% oxygen

We can see from the patient's arterial blood gases that he is hypoxic on 40% O_2. Whilst the rationale for reducing his oxygen to try to reduce his CO_2 levels might be reasonable with a lesser degree of hypoxia, simply reducing his oxygen in this situation is unlikely to improve his clinical situation.

A High flow oxygen (15 l)

We can see from the patient's arterial blood gases that he is in a respiratory acidosis (with CO_2 retention), so increasing his F_iO_2 further is going to exacerbate his CO_2 retention and worsen his clinical situation.

D Continuous positive airway pressure (CPAP)

We can see from the patient's arterial blood gases that he is in type 2 respiratory failure with CO_2 retention. CPAP provides a pneumatic splint to the airway and is effective in improving oxygenation in patients requiring high concentration of oxygen by the recruitment of collapsed airways. In this patient's case, this is not such a good option; it would be more prudent to use BIPAP in order to maximise gas exchange and help him to 'blow off' his increased levels of CO_2.

E Intubate and ventilate the patient

Intubation and ventilation would not be the first line of treatment in this patient. Following a trial of BIPAP, if the patient failed to respond and following a careful assessment of his pre-morbid state and co-morbidities, a decision to intubate could then be considered.

32400

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A high-resolution thoracic CT of a patient complaining of shortness of breath shows patchy ground-glass shadowing. However, there is much movement artefact.

What would be the best test to confirm restrictive lung disease due to a parenchymal disorder?

Your answer was incorrect

A CT in expiration phase

B FEV_1

C Arterial blood gas

D Chest X-ray

E KCO

Explanation

E KCO

Restrictive lung diseases are characterised by reduced lung volume, either because of an alteration in lung parenchyma or because of a disease of the pleura, chest wall or neuromuscular apparatus. The transfer coefficient will be reduced in patients with interstitial lung disease. With the presence of an abnormal CT, despite the fact that it is of poor quality, this would be enough evidence to suggest the diagnosis. Without the presence of a CT lesion, a normal KCO would merely rule out significant restrictive lung disease as a cause. With the abnormal CT seen here, abnormal KCO is useful supporting evidence for an underlying parenchymal disorder.

D Chest X-ray

Parenchymal lung disease is often of normal appearance in the early stages on a chest X-ray.

A CT in expiration phase

CT in expiration phase is useful for showing air-trapping in COPD.

B FEV_1

FEV$_1$1 is the forced expiratory volume in one second. It can be measured by a peak flow meter or spirometry. From that, airway calibre can be estimated. Based on FEV$_1$, obstructive airway disease can be diagnosed. It is limited in that it is effort dependent.

C Arterial blood gas

In an ABG, arterial blood is taken in a heparinised syringe to establish pH, PaO$_2$ and PaCO$_2$, amongst other measurements. It is very useful in diagnosing hypoxia or hypercapnia as well as analysing pH, and can therefore differentiate between types of respiratory failure. However, in this patient it may be of limited value as the patient may not be sufficiently unwell to have an abnormal ABG.

2322

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

A 56-year-old man presented to his GP with a 4-week history of feeling generally unwell. He described nasal congestion, with pain under his left eye, flu-like symptoms and lethargy. His GP made a diagnosis of sinusitis and gave him a course of antibiotics. He did not improve and developed breathlessness, cough and chest pain. He had previously been fit and his only past medical history was of an inguinal hernia repair 10 years previously. He worked in a shop. He smoked 10 cigarettes a day and drank about 10 units of alcohol a week. He was taking no regular medication.

On examination he looked unwell and pale. Observations: respiratory rate 34/min, BP 140/85 mmHg, pulse 120 bpm, temperature 36.6 °C. He had crusting of his nasal septum. His JVP was not elevated, and heart sounds were normal. Auscultation of his chest revealed fine crackles bilaterally. His abdominal and neurological examinations were unremarkable.

Urinalysis:

Protein	++
Blood	++
Nitrites	-
Leucocytes	-
Bilirubin	-

Haematology:

Investigation	**Result**	**Normal Values**
Haemoglobin (Hb)	81 g/l	135 - 175 g/l
White cell count (WCC)	6.3×10^9/l	$4.0 - 11.0 \times 10^9$/l
Mean corpuscular volume (MCV)	81 fl	80 - 100 fl
Platelets (PLT)	510×10^9/l	$150 - 400 \times 10^9$/l
Erythrocyte sedimentation rate (ESR)	114 mm/h	1 - 20 mm/h
C-reactive protein (CRP)	45 mg/l	< 10 mg/l
Sodium (Na^+)	142 mmol/l	135 - 145 mmol/l
Potassium (K^+)	5.9 mmol/l	3.5 - 5.0 mmol/l
Urea	21 mmol/l	2.5 - 6.5 mmol/l
Creatinine (Cr)	341 µmol/l	50 - 120 µmol/l
Bilirubin	18 µmol/l	1 - 22 µmol/l
Aspartate aminotransferase (AST)	31 u/l	10 - 40 u/l
Alkaline phosphatase (ALP)	191 u/l	30 - 150 u/l

Albumin	34 g/l	35 - 55 g/l

Spirometry:

FEV1	72% predicted
FVC	77% predicted
KCO	122% predicted

His chest X-ray is shown below:



Which investigation would help make a diagnosis?

Your answer was correct

A c-ANCA (anti-proteinase 3)

B Anti-GBM antibodies

C p-ANCA (anti-myeloperoxidase)

D *Legionella* antigen

E Echocardiogram

Explanation

A c-ANCA (anti-proteinase 3)

This is a question about pulmonary–renal syndromes (for a list see below).
The raised KCO and the CXR should make one think of pulmonary haemorrhage. This should narrow the diagnosis down to a vasculitis, Goodpasture's syndrome and SLE.

This patient has granulomatosis with polyangiitis. The clinical diagnosis is made by the triad of upper respiratory tract (nasal polyps and sinusitis), lower respiratory tract and renal involvement. It is a small-vessel vasculitis associated with granulomas.

The diagnosis of granulomatosis with polyangiitisis is suggested by a positive c-ANCA with the presence of an antibody to proteinase-3. The chest radiograph usually shows single or multiple nodules, fixed infiltrates or cavities. Pleural effusions and pulmonary infiltrates may also been seen. Diagnosis is usually confirmed with a biopsy. If untreated the median survival is just five months. Treatment is with intravenous cyclophosphamide and prednisolone, which reduces mortality to approximately 10%. Plasmapheresis would be first-line treatment if the disease was primarily renal.

Causes of pulmonary–renal syndromes:

- Systemic diseases:
  Granulomatosis with polyangiitis
  Microscopic polyangiitis
  Goodpasture syndrome
  Systemic lupus erythematosus
  Polyarteritis nodosa
  Henoch–Schönlein purpura
  Churg–Strauss syndrome (renal involvement less common)
- Primary pulmonary disease:
  Legionella pneumonia and interstitial nephritis
  Bacterial pneumonia with renal compromise secondary to sepsis
- Others:
  Pulmonary oedema with acute kidney disease
  Uraemic pneumonitis
  Right-sided bacterial endocarditis – may cause pulmonary embolic lesions and glomerulonephritis
  Iatrogenic glomerulonephritis with ciprofloxacin, e.g. in patients given ciprofloxacin for cystic fibrosis.

Image source: https://upload.wikimedia.org/wikipedia/commons/7/77/Wegeners_Granulomatosis_CXR.jpg (https://upload.wikimedia.org/wikipedia/commons/7/77/Wegeners_Granulomatosis_CXR.jpg)

B Anti-GBM antibodies

Anti-GBM antibodies are seen in Goodpasture syndrome, which isn’t associated with sinusitis; this drives us away from anti-GBM as the cause of this patient’s symptoms.

C p-ANCA (anti-myeloperoxidase)

p-ANCA is associated with microscopic polyangitis, which isn’t typically associated with upper respiratory tract disease. Churg–Strauss can be associated with positive p-ANCA, although it wouldn’t fit here as it’s associated with eosinophilia and asthma symptoms.

D *Legionella* antigen

This may be positive in *Legionella* pneumonia, but the subacute course of this patient’s symptoms counts against Legionnaire’s disease as the underlying diagnosis.

E Echocardiogram

An echocardiogram would not show diagnostic features to confirm diagnosis of granulomatosis with polyangiitis. It would be of most value if left ventricular failure, rather than pulmonary infiltration, was suspected.

70089

Rate this question:

End Session

Previous Question

Tag Question

Feedback

End Review

Difficulty: Easy

Peer Responses %



Show More Questions Like This

Session Progress

Responses Correct:	82
Responses Incorrect:	219
Responses Total:	301
Responses - % Correct:	27%

Granulomatosis with Polyangiitis

Media

Expanded explanations

Flashcards

M. Alam

Created by Mahbub Alam
Chhondo
From Bangladesh